Derivatives

A *derivative* (slope of a line) is measuring the sensitivity of a function's output with respect to it's input. In other words, it measures the *steepness* of the graph of a function.

If we have a linear function like below where for every input x it returns a y value of 3 times the value of x:

$$f(x) = 3x$$

If we have a point where x=2 then y=6

$$f(2) = 3 \cdot 2 = 6$$

Therefore, our first point is at:

If we say x = 6.001, then:

$$f(6.001) = 3 \cdot 6.001 = 18.003$$

Therefore, our second point is at:

If we want to know what the derivative is we take the change in y and divide by the change in x:

$$\frac{\Delta y}{\Delta x} = \frac{6.001 - 2}{18.003 - 6} = \frac{4.001}{12.003} = .\overline{33}$$

Therefore, we say the *derivative of our function* f(x) *with respect to* x is $.\overline{33}$.

$$\frac{df(x)}{dx} = \overline{.33}$$

Or, more precisely, we would use the delta δ term which says: The delta (or rate of change), which is the derivative for the function f(x) with respect to x is . $\overline{33}$.

$$\delta = \frac{df(x)}{dx} = .\overline{33}$$

Derivatives and Quadratics

For getting the derivative of a quadratic function we will need to calculate it with respect to the tangent line as shown below. Also, since quadratics are complex function and don't yield a simple straight line we only know the partial derivative ∂ at any given point since it is constantly changing.

$$\frac{\partial f(x)}{\partial x}$$

