Several Variable Calculus and Differential Equations Review

Harry Han

Dec 2022

Contents

1 Vectors and Geometry **3**

2 Partial Derivatives **4**
2.1 Limits 4
2.2 Partial Derivatives 4
2.3 Maximum and Minimum 6
2.3.1 Function of Three Variables 7
2.3.2 Lagrange Multiplier 7

3 Multiple Integrals **7**
3.1 Double And Triple Integrals 7
3.2 Change of Variables 8
3.2.1 Double Integrals: To Polar Coordinates . . . 8
3.2.2 Triple Integrals: To Cylindrical Coordinates 8
3.2.3 Triple Integrals: To Spherical Coordinates . 9
3.2.4 Change of Variable in General: Jacobian . . . 9

4 Vector Calculus **10**
4.1 Vector Fields 10
4.1.1 Conservative Vector Fields 10
4.2 Curl and Divergence 10
4.3 Line Integral 11
4.4 Surface Integral 11
4.5 Theoremae Trinitatis 13

5 Differential Equations **13**
5.1 Classification 13
5.2 First Order Ordinary Differential Equation 14
5.3 Second Order Ordinary Differential Equation . . . 15
5.3.1 General Solutions to Second Order Homogeneous ODE 15
5.3.2 Solutions to Certain Second Order Non-Homogeneous ODE 16

6 Quaedam Formulae Utillimae **17**
6.1 Trignometric Formulae 17
6.2 Common Integrals 17
6.3 Other Common Computations 17

1 Vectors and Geometry

Theorem 1.0.1 (Vector Computation)**.** *Basics computation rules for vectors:*

1. $\underline{a} \cdot \underline{b} = \cos\theta |\underline{a}||\underline{b}|$
2. $|\underline{a} \times \underline{b}| = \sin\theta |\underline{a}||\underline{b}|$
3. $\underline{a} \times \underline{b} = -\underline{b} \times \underline{a}$
4. $\underline{a} \cdot (\underline{b} \times \underline{c}) = (\underline{a} \times \underline{b}) \cdot \underline{c}$
5. $\underline{a} \times (\underline{b} \times \underline{c}) = (\underline{a} \cdot \underline{c})\underline{b} - (\underline{a} \cdot \underline{b})\underline{c}$
6. $\underline{a} \cdot (\underline{b} \times \underline{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$, *moreover, this equal to the area of the parallelepiped specified by the vectors* $\underline{a}, \underline{b}, \underline{c}$

Definition 1.1 (Line and Plane in 2D and 3D space)**.**

Definition 1.2 (Vector Function)**.** We consider the vector function $\underline{\mathrm{v}}(\mathrm{t})$ whose domain is $\mathbb{R}$ and whose range is $\mathbb{R}^3$. It has three components $f(t), g(t) and h(t)$ and can be written in the form:

$$\underline{v}(t) = \begin{pmatrix} f(t) \\ g(t) \\ h(t) \end{pmatrix} = f(t)\underline{i} + g(t)\underline{j} + h(t)\underline{k}$$

Vector function obey most of the integrating and differentiation rules of a normal function.

Theorem 1.0.2 (Arc Length)**.** *The length of the curve traced out by the postion vector function* $\underline{r}(t) = x(t)\underline{i} + y(t)\underline{j} + z(t)\underline{k}$ *over the interval* $a \leq t \leq b$ *is*

$$s(t) = \int_a^b \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} dt = \int_a^b |\underline{r}'(t)| dt$$

Definition 1.3 (Unit Tangent Vector)**.** For vector function $\underline{r}(t)$, its unit tangent vector is $\underline{T} = \dfrac{\underline{r}(t)}{|\underline{r}(t)|}$

Definition 1.4 (Curvature)**.** For vector function $\underline{r}(t)$, its curvature is

$$\kappa = \left| \frac{\underline{T}'(t)}{s'(t)} \right| = \frac{|\underline{T}'(t)|}{|\underline{r}'(t)|} = \frac{|\underline{r}'(t) \times \underline{r}''(t)|}{|\underline{r}'(t)|^3}$$

Definition 1.5 (Normal and binormal Vector)**.** For vector function $\underline{r}(t)$, its unit tangent vector is $\underline{T} = \dfrac{\underline{r}(t)}{|\underline{r}(t)|}$, and its unit normal vector is $\underline{N}(t) = \dfrac{\underline{T}'(t)}{|\underline{T}'(t)|}$, and its binormal vector is $\underline{T}(t) \times \underline{N}(t)$

2 Partial Derivatives

2.1 Limits

Definition 2.1 (Limits of function of two variables)**.** Let $f(x, y)$ be a function of two variables. The limit of $f(x, y)$ as (x, y) approaches (a, b) is defined as

$$\lim_{(x,y)\to(a,b)} f(x, y) = L$$

if for every $\epsilon > 0$, there exists a $\delta > 0$ such that $\sqrt{(x-a)^2 + (y-a)^2} < \delta$ implies $|f(x, y) - L| < \epsilon$. Significantly, the limit of $f(x, y)$ as $(x, y) \to (a, b)$ is independent of the path of x and y. i.e. if assuming the relationship $y = Y_1(x)$ and $y = Y_2(x)$, provided $Y(a) = b$, we arrives to different value of the limite of $f(x, Y(x))$ as $x \to a$, then limit of $f(x, y)$ as (x, y) approach (a, b) does not exist.

It is sometimes easier to evaluate the limit using polar coordinates (or other kinds of transformation). Let $(x, y) = (r\cos\theta, r\sin\theta)$, then $(x, y) \to 0 \iff r \to 0$ All limit law for function of one variable also apply to function of multiple variables.

Definition 2.2 (Continuity)**.** Let $f(x, y)$ be a function of two variables. $f(x, y)$ is continuous at $(a, b) \iff \lim\limits_{(x,y)\to(a,b)} = f(a, b)$

2.2 Partial Derivatives

Definition 2.3 (Partial Derivative)**.**

$$\frac{\partial f}{\partial x} = \lim_{\Delta x\to 0} \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

More generally, for n variables, we have

$$f_{x_1}(x_1, x_2, \ldots, x_n) = \frac{\partial f}{\partial x_1} = \lim_{\Delta x_1\to 0} \frac{f(x_1 + \Delta x_1, x_2, \ldots, x_n) - f(x_1, x_2, \ldots, x_n)}{\Delta x_1}$$

NB: the notation of partial derivative:

$$f_x(x,y) = \frac{\partial f}{\partial x} = f_x = \frac{\partial}{\partial x} f(x,y)$$

Theorem 2.2.1 (Clairaut's Theorem)**.** *Let $f(x,y)$ be a function of two variables. Then*

$$f_{xy} = f_{yx}$$

More generally the order of the partial derivative does not matter.

Remark 2.1. NB: Tangent plane and linear approximation.

Definition 2.4 (Differentiability)**.** Let $z = f(x,y), \Delta z = f(a+\Delta x, b+\Delta y) - f(x,y)$, then the function is differentialbe if and only if $(\epsilon_1, \epsilon_2) \to (0,0)$ as $(\Delta x, \Delta y) \to (0,0)$, where ϵ_1, ϵ_2 are defined as

$$\Delta z = f_x(a,b)\Delta x + f_y(a,b)\Delta y + \epsilon_1 \Delta x + \epsilon_2 \Delta y$$

Moreover, it can be shown that z is differentiable at (a,b) if and only if the partial derivative f_x, f_y exisits and are continuous near (a,b)

Definition 2.5 (Differentials)**.** $w = f(x,y,z)$ the total differential dw is defined as

$$dw = \frac{\partial w}{\partial x} dx + \frac{\partial w}{\partial y} dy + \frac{\partial w}{\partial z} dz$$

Theorem 2.2.2 (Chain Rule)**.** *Suppose $z = f(x,y)$ and $x = g(t)$, $y = h(t)$ then*

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t}$$

Suppose $z = f(x,y)$ and $x = g(t,u)$, $y = h(t,u)$ then

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial t} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial t}, \frac{\partial z}{\partial u} = \frac{\partial z}{\partial x}\frac{\partial x}{\partial u} + \frac{\partial z}{\partial y}\frac{\partial y}{\partial u}$$

Theorem 2.2.3 (Implicit Differentiation)**.** *Suppose $F(x,y,z) = 0$, and z is a implicit funtion of x, then:*

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z},$$

If z is a implicit function of y, then:

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z}$$

Definition 2.6 (Gradient)**.** Let $f(x, y)$ be a function of two variables, then the gradient of f is defined as a vector function:

$$\nabla f(x, y) = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}\right)$$

Definition 2.7 (Directional Derivative)**.** For funtion $f(x, y)$, the directional derivative at (a, b) at the direction of the unit vector $\underline{u} = (u_1, u_2)$ in $x - y$ plane is:

$$\nabla f \cdot \underline{u} = f_x(a, b)u_1 + f_y(a, b)u_2$$

Note it is a constant.

Theorem 2.2.4 (Maximun Directional Derivative)**.** *Suppose $f(x, y)$ is differentiable at (a, b), then the directional derivative of f at (a, b) in the direction of u is maximized if and only if u is parallel to the gradient of f at (a, b).*

Proof. Notice the directional derivative equals to

$$\nabla f \cdot \underline{u} = \cos\theta |\nabla f||\underline{u}|$$

Which is maximized when $\theta = 0$. □

Remark 2.2. Tangent plain to level surface ans normal line: see lecture notes page 27.

2.3 Maximum and Minimum

Definition 2.8 (Critical Point)**.** Let $f(x, y)$ be a function of two variables, then (a, b) is a critical point of f if $f_x(a) = 0 and f_y = (b)0$, or one of its partialderivative does not exist.

Theorem 2.3.1. *If (a, b) is a local maximum or minimum of $f(x, y)$, then (a, b) is a critical point of f*

Theorem 2.3.2 (Second Derivative Test)**.** *Suppose the secound partial derivative of $f(x, y)$ exists at (a, b) and is continous around it, and $f_x(a, b) = f_y(a, b) = 0$ Let*

$$D(a, b) = f_{xx}(a, b)f_{yy}(a, b) - f_{xy}(a, b)^2 = \begin{vmatrix} f_{xx}(a, b) & f_{xy}(a, b) \\ f_{xy}(a, b) & f_{yy}(a, b) \end{vmatrix}$$

Then

1. *If $D(a, b) > 0$ and $f_{xx} > 0$, then (a, b) is a local minimum of $f(x, y)$*
2. *If $D(a, b) > 0$ and $f_{xx} < 0$, then (a, b) is a local maximum of $f(x, y)$*
3. *If $D(a, b) < 0$, then (a, b) is a saddle point of $f(x, y)$*
4. *If $D(a, b) = 0$, the test is inconclusive*

2.3.1 Function of Three Variables

Theorem 2.3.3 (Second Derivative Test)**.** *Suppose the secound partial derivative of $f(x, y, z)$ exists at (a, b, c) and is continous around it, and $f_x(a, b, c) = f_y(a, b, c) = f_z(a, b, c) = 0$ Let*

$$D_1(a, b, c) = f_{xx}(a, b, c)$$

$$D_2(a, b, c) = \begin{vmatrix} f_{xx}(a, b, c) & f_{xy}(a, b, c) \\ f_{xy}(a, b, c) & f_{yy}(a, b, c) \end{vmatrix}$$

$$D_3(a, b, c) = \begin{vmatrix} f_{xx}(a, b, c) & f_{xy}(a, b, c) & f_{xz}(a, b, c) \\ f_{xy}(a, b, c) & f_{yy}(a, b, c) & f_{yz}(a, b, c) \\ f_{xz}(a, b, c) & f_{yz}(a, b, c) & f_{zz}(a, b, c) \end{vmatrix}$$

Then

1. *If $D_1(a, b, c) > 0$ and $D_2(a, b, c) > 0$ and $D_3(a, b, c) > 0$, then (a, b, c) is a local minimum of $f(x, y, z)$*
2. *If $D_1(a, b, c) < 0$ and $D_2(a, b, c) > 0$ and $D_3(a, b, c) < 0$, then (a, b, c) is a local maximum of $f(x, y, z)$*
3. *If $D_3 \neq 0$ and $(a, b, c) is neitheramaxiumnoraminimum$, then (a, b, c) is a saddle point of $f(x, y, z)$*
4. *If $D_3(a, b, c) = 0$, the test is inconclusive*

2.3.2 Lagrange Multiplier

Theorem 2.3.4 (Lagrange Multiplier)**.** *Let $f(x, y)$ be a function of two variables and differential along the smooth curve $g(x, y) = 0$ where $\nabla g(x, y) \neq 0$. if f attains minimum value along $g = 0$ at (a, b), then there exists $\lambda \in \mathbb{R}$ such that*

$$\nabla f(a, b) = -\lambda \nabla g(a, b)$$

3 Multiple Integrals

3.1 Double And Triple Integrals

Definition 3.1 (Double Integral)**.** Ommited.

Definition 3.2 (Average Value)**.** The average value of a function $f(x, y)$ over a region D is defined as

$$f_{avg} = \frac{1}{A(D)} \iint_D f(x, y) dA$$

Theorem 3.1.1 (Fubini's Theorem)**.** *If $f(x, y)$ is continuous on the rectangle $R = (x, y)\mathbb{R}^2, a \leq x \leq b, c \leq y \leq d$, then*

$$\iint_R f(x, y) dA = \int_a^b \int_c^d f(x, y) dy dx = \int_c^d \int_a^b f(x, y) dx dy$$

Corollary 3.1.2. *Suppose $f(x, y) = g(x)h(y)$ on $R = [a, b] \times [c, d]$, then*

$$\iint_R f(x, y) dA = \int_a^b g(x) dx \int_c^d h(y) dy$$

Remark 3.1. Triple integral, see example 50 page 47.

3.2 Change of Variables

3.2.1 Double Integrals: To Polar Coordinates

Theorem 3.2.1 (Change of Variables)**.** *For (x, y) in Cartesian coordinates, recall the polar coordinate (r, θ) is defined by: $r^2 = x^2 + y^2, x = r\cos\theta, y = r\sin\theta$*

If the area R is given by $r_0 \leq r \leq r_1, \theta_0 \leq \theta \leq \theta_1$, then:

$$\iint_R f(x, y) dA = \iint_D f(r\cos\theta, r\sin\theta) r dr d\theta = \int_{r_0}^{r_1} \int_{\theta_0}^{\theta_1} f(r\cos\theta, r\sin\theta) r dr d\theta$$

Notice the extra r in the integrand.

3.2.2 Triple Integrals: To Cylindrical Coordinates

Theorem 3.2.2 (Change of Variables To Cylindrical Coordinates)**.** *Recall the cylindrical coordinate (r, θ, z) is defined by: $r^2 = x^2 + y^2, x = r\cos\theta, y = r\sin\theta$ If the volume V is given by $r_0 \leq r \leq r_1, \theta_0 \leq \theta \leq \theta_1, z_0 \leq z \leq z_1$, then:*

$$\iiint_V f(x, y, z) dV = \iiint_D f(r\cos\theta, r\sin\theta, z) r dz dr d\theta = \int_{r_0}^{r_1} \int_{\theta_0}^{\theta_1} \int_{z_0}^{z_1} f(r\cos\theta, r\sin\theta, z) r dz dr d\theta$$

Notice the extra r in the integrand.

3.2.3 Triple Integrals: To Spherical Coordinates

Theorem 3.2.3 (Change of Variables To Spherical Coordinates)**.** *Recall the spherical coordinate* (r, θ, ϕ) *is defined by:* $r^2 = x^2+y^2+z^2, x = r\sin\phi\cos\theta, y = r\sin\theta\sin\phi, z = r\cos\phi$

If the volume V *is given by* $r_0 \le r \le r_1, \theta_0 \le \theta \le \theta_1, \phi_0 \le \phi \le \phi_1$*, then:*

$$\iiint_V f(x,y,z)dV = \iiint_D f(r\sin\phi\cos\theta, r\sin\theta\sin\phi, r\cos\phi)r^2\sin\phi dr d\theta d\phi$$

Notice the extra $r^2\sin\phi$ *in the integrand.*

Remark 3.2. More excercises needed.

3.2.4 Change of Variable in General: Jacobian

Definition 3.3 (Jacobian)**.** Consider the transformation from u, v plane to x, y plane, where $x = x(u,v), y = y(u,v)$ For bijective transformation, the jacobian is defined as:

$$J_2 = \frac{\partial x \partial y}{\partial u \partial v} = \frac{\partial x}{\partial u}\frac{\partial y}{\partial v} - \frac{\partial x}{\partial v}\frac{\partial y}{\partial u} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$

The jacobian for transformation from u, v, w plane to x, y, z plane, where $x = x(u,v,w), y = y(u,v,w), z = z(u,v,w)$ is defined as:

$$J_3 = \frac{\partial x \partial y \partial z}{\partial u \partial v \partial w} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{vmatrix}$$

Theorem 3.2.4 (Change of Variable in General)**.** *If the transformation from* u, v *plane to* x, y *plane is bijective, then:*

$$\iint_D f(x,y)dA = \iint_D g(u,v)|J_2|dudv$$

Where $g(u,v) = f(x(u,v), y(u,v))$*.*

If the transformation from u, v, w *plane to* x, y, z *plane is bijective, then:*

$$\iiint_V f(x,y,z)dV = \iiint_V g(u,v,w)|J_3|dudvdw$$

Where $g(u,v,w) = f(x(u,v,w), y(u,v,w), z(u,v,w))$*.*

Notice the Jacobian is taken as absolute value.

Remark 3.3. More excercises needed. See Example 54, lecture note page 51.

4 Vector Calculus

4.1 Vector Fields

Definition 4.1 (Vector Field)**.** A vector field is a function $\underline{F} : \mathbb{R}^n \to \mathbb{R}^n$. The input and output are n dimensional vectors.

4.1.1 Conservative Vector Fields

Theorem 4.1.1 (Conservative Vector Fields)**.** *Vector field $\underline{F}(x, y)$ is conservative, provided that it is defined on all $\mathbb{R}^2$, if and only if*

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$$

For vector field $\underline{G}(x, y, z)$, provided it is defined on all $\mathbb{R}^3$, it is conservative if and only if:

$$\underline{\nabla} \times \underline{G} = 0$$

Definition 4.2 (Conservative Vector Field)**.** A vector field $\underline{F}$ is conservative if there exists a function $f : \mathbb{R}^n \to \mathbb{R}$ such that $\underline{\nabla} f = \underline{F}$.

4.2 Curl and Divergence

Definition 4.3 (Curl)**.** The curl of a vector field $\underline{F} = F_x\underline{i} + F_y\underline{j} + F_z\underline{k}$ is defined as:

$$\underline{\nabla} \times \underline{F} = \underline{i}\left(\frac{\partial F_z}{\partial y} - \frac{\partial F_y}{\partial z}\right) + \underline{j}\left(\frac{\partial F_x}{\partial z} - \frac{\partial F_z}{\partial x}\right) + \underline{k}\left(\frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y}\right)$$

Moreover, curl of $\underline{F}$ can be rewritten as below:

$$\underline{\nabla} \times \underline{F} = \begin{vmatrix} \underline{i} & \underline{j} & \underline{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{vmatrix}$$

Definition 4.4 (Divergence)**.** The divergence of a vector field $\underline{F} = F_x\underline{i} + F_y\underline{j} + F_z\underline{k}$ is defined as:

$$\underline{\nabla} \cdot \underline{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z}$$

Corollary 4.2.1. *Two facts about curl and divergence:*

1. *div (curl $\underline{F}$) = 0*
2. *curl (∇f) = 0*

4.3 Line Integral

Definition 4.5 (Line Integral)**.** If f is a function of three variable, the line integral along the curve γ is defined as:

$$\int_\gamma f(x,y,z)ds = \int_{t_0}^{t_1} f(x(t),y(t),z(t))\sqrt{\frac{dx}{dt}^2 + \frac{dy}{dt}^2 + \frac{dz}{dt}^2}\,dt$$

Where γ is parametrized by $x = x(t), y = y(t), z = z(t)$, from t_0 to t_1.

Moreover

$$\int_\gamma F dx = \int_{t_0}^{t_1} F(x(t),y(t),z(t))\frac{dx}{dt}dt$$

Line integral alone y and z are defined similarly.

Definition 4.6 (Line Integral of a Vector Field)**.** The line integral of a vector field $\underline{F}$ along a curve γ is defined as:

$$\int_\gamma \underline{F}\cdot d\underline{r} = \int_\gamma \underline{F}(\underline{r})\cdot \underline{r}'dt = \int_\gamma P_x dx + P_y dy + P_z dz$$

Where $\underline{F} = \underline{i}P_x + \underline{j}P_y + \underline{k}P_z$

Theorem 4.3.1 (Fundamental Theorem of Line Integral)**.** *If $\underline{F}$ is conservative, and $\underline{\nabla} f = \underline{F}$. Let γ be a curve which starts from (x_0, y_0) and ends at (x_1, y_1). Then*

$$\int_\gamma \underline{F}\cdot d\underline{r} = f(x_1,y_1) - f(x_0,y_0)$$

4.4 Surface Integral

Theorem 4.4.1 (Surface Area)**.** *The surface area of a parametrized surface $\underline{r}(u,v) = \underline{i}x(u,v) + \underline{j}y(u,v) + \underline{k}z(u,v)$, for $(u,v) \in \mathbb{D}$ is:*

$$\iint_{\mathbb{D}} |\underline{r}_u \times \underline{r}_v| dA$$

Where $\underline{r}_u = \dfrac{\partial \underline{r}}{\partial u}$ and $\underline{r}_v = \dfrac{\partial \underline{r}}{\partial v}$.

Notice $\underline{r}_u$ and $\underline{r}_v$ are tangent vectors to the surface, and $||$ operator returns their magnitude.

Corollary 4.4.2 (When z is a function of x and y)**.** *For surface paramatrized as $\underline{r} = \underline{i}x + \underline{j}y + \underline{k}f(x,y)$, the surface area of the surface for $(x,y) \in \mathbb{D}$ is*

$$\iint_{\mathbb{D}} \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2}\,dA$$

Definition 4.7 (Surface Integral)**.** Consider function $f(x, y, z)$ defined on the surface $\mathbb{S}$ which is described by the vector function:

$$\underline{r}(u, v) = \underline{i}x(u, v) + \underline{j}y(u, v) + \underline{k}z(u, v), (u, v) \in \mathbb{D}$$

Then the surface integral of f over the surface $\mathbb{S}$ is defined as:

$$\iint_{\mathbb{S}} f(x, y, z)dS = \iint_{\mathbb{D}} f(\underline{r}(u, v))|\underline{r}_u \times \underline{r}_v|dA$$

Where $\underline{r}_u = \dfrac{\partial \underline{r}}{\partial u}$ $\underline{r}_v = \dfrac{\partial \underline{r}}{\partial v}$, and$f(\underline{r}(u, v)) = f(x(u, v), y(u, v), z(u, v))$

Remark 4.1. More excercises needed.

Definition 4.8 (Oriented Surface)**.** Let $\underline{r}(u, v)$ be a vector function on $\mathbb{D}$ that describes the surface S. $\dfrac{\underline{r}_u \times \underline{r}_v}{|\underline{r}_u \times \underline{r}_v|}$ is its unit normal vector to S.

Moreover, for surface $z = f(x, y)$ the unit normal vector is:

$$\frac{-\underline{i}\dfrac{\partial f}{\partial x} - \underline{j}\dfrac{\partial f}{\partial y} + \underline{k}}{\sqrt{1 + \left(\dfrac{\partial f}{\partial x}\right)^2 + \left(\dfrac{\partial f}{\partial y}\right)^2}}$$

Definition 4.9 (Surface Integral Over Vector Field)**.** Let $\underline{F}(x, y, z)$ be a vector field defined on the surface S. Then the surface integral of $\underline{F}$ over S is defined as:

$$\iint_S \underline{F} \cdot d\underline{S} = \iint_S \underline{F} \cdot \underline{\hat{n}}dS = \iint_{\mathbb{D}} \underline{F}(\underline{r}(u, v)) \cdot \frac{\underline{r}_u \times \underline{r}_v}{|\underline{r}_u \times \underline{r}_v|} \cdot |\underline{r}_u \times \underline{r}_v|dA$$

Which equals to:

$$\iint_{\mathbb{D}} \underline{F}(\underline{r}(u, v)) \cdot (\underline{r}_u \times \underline{r}_v)\, dA$$

Moreover, for surface $z = g(x, y)$, we have:

$$\underline{F} \cdot |\underline{r}_u \times \underline{r}_v| = (F_x\underline{i} + F_y\underline{j} + F_z\underline{k}) \cdot \left(-\frac{\partial g}{\partial x}\underline{i} - \frac{\partial g}{\partial y}\underline{j} + \underline{k}\right)$$

And the surface integral is:

$$\iint_S \underline{F} \cdot d\underline{S} = \iint_{\mathbb{D}} -F_x\frac{\partial g}{\partial x} - F_y\frac{\partial g}{\partial y} + F_z dA$$

4.5 Theoremae Trinitatis

Theorem 4.5.1 (Green's Theorem)**.** *Let γ be positively oriented curve, simply close, and piecewise smooth curve in x, y plane. Let Γ be the region bounded by γ. If $P(x, y)$ and $Q(x, y)$ are continuous functions on Γ, then*

$$\int_\gamma Pdx + Qdy = \iint_\Gamma \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dA$$

Theorem 4.5.2 (Stokes' Theorem)**.** *Let γ be positively oriented, simply close, and piecewise smooth curve in x, y plane. Let Γ be the region bounded by γ. If $\underline{F}(x, y, z)$ is a continuous vector field on Γ, then*

$$\iint_\Gamma \underline{\nabla} \times \underline{F} \cdot d\underline{S} = \int_\gamma \underline{F} \cdot d\underline{r}$$

Theorem 4.5.3 (Divergence Theorem)**.** *Let E be a simple solid region bounded by a positively oriented surface (outward) $\mathbb{S}$. Let $\underline{F} : \mathbb{R}^3 \to \mathbb{R}^3$ be a vector field that has continuous partial derivative in E, then:*

$$\iint_\mathbb{S} \underline{F} \cdot d\underline{S} = \iiint_E \underline{\nabla} \cdot \underline{F} dV$$

5 Differential Equations

5.1 Classification

Definition 5.1 (Classification of Diferential Equation)**.**

1. ordinary differential equation (ODE) involves only differentiation of one variable
2. partial differential equation (PDE) involves differentiation of more than one variable.
3. Order of differential equation is the highest order of differentiation in the equation.

The nth order linear differential equation has the general form:

$$\frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \cdots + a_{n-1} \frac{dy}{dx} + a_n y = a_0(x)$$

Where a_i asre functions of x. The equation is called homogeneous if $a_0(x) = 0$.

If $S_1(x), S_2(x) \ldots$ are solution to the homogeneous equation, then any of its linear combinations are also a solution.

Theorem 5.1.1 (Uniqueness of the solution to Inital Value Problem)**.** *Consider the general n^{th} order linear differential equation:*

$$f(y^n, y^{n-1}, \ldots, y, x) = 0$$

Provided with inital values

$$y(x_0) = a_0, y^1(x_1) = a_1 \ldots y^n(x_n) = a_n$$

There is one unique solution to this Initial Value Problem.

5.2 First Order Ordinary Differential Equation

Theorem 5.2.1 (Solutions to Linear First Order ODE)**.** *The linear ODE of the form:*

$$y' + P(x)y = Q(x)$$

has the general solution:

$$y(x) = e^{-\int P(x)dx}\left(\int Q(x)e^{\int P(x)dx}dx + C\right)$$

Where C is an arbitrary constant.

Theorem 5.2.2 (Separable ODE)**.** *ODE are separarble if it can be written in the form:*

$$\frac{dy}{dx}h(y) = g(x)$$

And the solution is given by simply integration:

$$\int h(y)dy = \int g(x)dx$$

Theorem 5.2.3 (Homogeneous ODE (Another Homogeneous))**.** *A ODE is called homogeneous if it can be written in the form:*

$$\frac{dy}{dx} = g(\frac{y}{x})$$

It can be solved by introducing a new variable $u = \frac{y}{x}$. Notice $\frac{dy}{dx} = u + x\frac{du}{dx}$.

Theorem 5.2.4 (Exact ODE)**.** *The ODE is exact if it can be written in the form:*

$$\frac{\partial \psi}{\partial x} + \frac{\partial \psi}{\partial y}\frac{dy}{dx} = 0$$

Where ψ is a function of x and y.

And its general the solution is $\psi(x, y) = C$.

Moreover, a equation in the form of $M(x, y) + \frac{dy}{dx}N(x, y) = 0$ is if and only if $\frac{\partial M}{y} = \frac{\partial N}{x}$. c.f. the determination of conservative vector field.

Remark 5.1 (On Integration Factor). The process of finding the solution of a differential equation many times involves heavily upon the creative application of algebraic manipulation and the computation capabilities of the problem solver. For a particular kind of equation whose general solution is unknown, there are many times tricks that may be useful. One of them is integration factor.

For example, consider the following equation:

$$3xy + y^2 + (x^2 + xy)y' = 0$$

This equation can be turned into an exact equation by multiplying both sides by x, and the ψ function can be found to be: $x^3y + \frac{1}{2}x^2y^2$.

5.3 Second Order Ordinary Differential Equation

5.3.1 General Solutions to Second Order Homogeneous ODE

Theorem 5.3.1 (General Solutions to Second Order Homogeneous ODE)**.** *Let $y'' + a_1y' + a_2y = 0$ be a second order homogeneous ODE. Substitute the function $T(x) = e^{rx}$, where r is a constant:*

$$r^2e^{rx} + a_1re^{rx} + a_2e^{rx} = 0 \iff r^2 + a_1r + a_2 = 0 \tag{1}$$

Here we have three cases.

***Case 1**: equation 1 has two distinct real roots r_1 and r_2.*

Then the general solution is:

$$y(x) = C_1e^{r_1x} + C_2e^{r_2x}$$

***Case 2**: equation 1 has two repeated real roots r_1.*

Then the general solution is:

$$y(x) = C_1e^{r_1x} + C_2xe^{r_1x}$$

Case 3: *equation* 1 *has two complex roots* $r_1 + im_1$ *and* $r_1 - im_1$, *where* i *is the imaginary unit.*

Recall Euler's formula: $e^{ix} = \cos x + i \sin x$. *Then the general solution is:*

$$y(x) = C_1 e^{r_1 x} e^{im_1 x} + C_2 e^{r_1 x} e^{-im_1 x}$$

Which is equivalent to:

$$y(x) = C_3 \cos(m_1 x) e^{r_1 x} + C_4 \sin(m_1 x) e^{r_1 x}$$

5.3.2 Solutions to Certain Second Order Non-Homogeneous ODE

Theorem 5.3.2 (Method of Variation of Parameters)**.** *Consider the linear non-homogeneous second order ODE:*

$$ay'' + by' + cy = g(t) \tag{2}$$

Suppose the homogeneous equation ($ayy'' + by' + cy = 0$) *has the general solution:*

$$y_c(t) = C_1 y_1(t) + C_2 y_2(t)$$

Then the particular solution of the non-homogeneous equation 2 *is:*

$$y_p(t) = -y_1(t) \int \frac{y_2(t) g(t)}{aW(y_1, y_2)} dt + y_2(t) \int \frac{y_1(t) g(t)}{aW(y_1, y_2)} dt$$

Where W *is Wronskian operator and* $W(y_1, y_2) = y_1 y_2' - y_2 y_1'$.

6 Quaedam Formulae Utillimae

6.1 Trignometric Formulae

1. $\sin^2\theta + \cos^2\theta = 1$
2. $\cos(2\theta) = \cos^2\theta - \sin^2\theta = 2\cos^2\theta - 1$

6.2 Common Integrals

1. $\int \cos^2\theta = \frac{\theta}{2} + \frac{\sin 2\theta}{4}, \int_0^\pi \cos^2\theta = \frac{\pi}{2}$
2. $\int \sin^2\theta = \frac{\theta}{2} - \frac{\sin 2\theta}{4}, \int_0^\pi \sin^2\theta = \frac{\pi}{2}$

6.3 Other Common Computations

1. For $\underline{r}(\phi,\theta) = \sin\phi\cos\theta\underline{i} + \sin\phi\sin\theta\underline{j} + \cos\phi\underline{k}, |\underline{r}_\phi \times \underline{r}_\theta| = \sin\phi$