

# The Square Tile Problem


Harry Richman

University of Michigan

15 November 2018


# The Problem

Floor:



# The Problem

Floor:




Problem:

- square tiles of any size, \$1 each
- what is minimal cost  $C(11, 13)$ ?

# The Problem

Floor:




Theorem (Dehn):

A square tiling of a  $\mathbb{Q}$ -rectangle must have  $\mathbb{Q}$ -side lengths.

# The Problem

Floor:




Problem:

- square tiles of any **integer** size, \$1 each
- what is minimal cost  $C(11, 13)$ ?


# The Problem

What tilings work?



# The Problem


What tilings work?



“dumb” tiling:  $C(11, 13) \leq 11 \cdot 13 = 143$

# The Problem

What tilings work?




“greedy” tiling:  $C(11, 13) \leq 1 + 5 + 2 = 8$



# The Problem


What tilings work?



better tiling:  $C(11, 13) \leq 6$

# The Problem

What tilings work?



better (best??) tiling:  $C(11, 13) \leq 6$

# The Problem: general case

Problem: floor of dimensions  $m$  by  $n$

- square tiles of integer size, \$1 each
- what is minimal cost  $C(m, n)$ ?

## The Problem: general case

Problem: floor of dimensions  $m$  by  $n$

- square tiles of integer size, \$1 each
- what is minimal cost  $C(m, n)$ ?

Upper bound: “dumb” tiling

$$C(m, n) \leq mn.$$

# The Problem: general case

Problem: floor of dimensions  $m$  by  $n$

- square tiles of integer size, \$1 each
- what is minimal cost  $C(m, n)$ ?

Upper bound: “dumb” tiling

$$C(m, n) \leq mn.$$

Lower bound: assume  $m \leq n$ ; largest tile is  $\leq m$  by  $m$

$$\Rightarrow C(m, n) = (\# \text{ squares}) \geq \frac{\text{total area}}{\text{max. tile area}} = \frac{mn}{m^2} = n/m$$

## The Problem: general case

OPEN Problem: floor of dimensions  $m$  by  $n$

- square tiles of integer size, \$1 each
- what is minimal cost  $C(m, n)$ ?

Upper bound: “dumb” tiling


$$C(m, n) \leq mn.$$

Lower bound: assume  $m \leq n$ ; largest tile is  $\leq m$  by  $m$

$$\Rightarrow C(m, n) = (\# \text{ squares}) \geq \frac{\text{total area}}{\text{max. tile area}} = \frac{mn}{m^2} = n/m$$

# The Problem: better bounds


Upper bound: “greedy” tiling



# The Problem: better bounds

Upper bound: “greedy” tiling

$$C(11, 13) \leq 1 + 5 + 2 = 8$$






# The Problem: better bounds

Upper bound: “greedy” tiling


$$C(11, 13) \leq 1 + 5 + 2 = 8$$



Continued fraction:  $\frac{13}{11} = 1 + \frac{2}{11} = 1 + \frac{1}{5 + \frac{1}{2}}$

# The Problem: better bounds

Upper bound: “greedy” tiling       $C(11, 13) \leq 1 + 5 + 2 = 8$



Continued fraction:  $\frac{13}{11} = 1 + \frac{2}{11} = 1 + \frac{1}{5 + \frac{1}{2}}$

$\Rightarrow C(m, n) \leq (\text{sum of continued-fraction terms})$

# The Problem: better bounds

Lower bound:  $C(m, n) \geq n/m$

Can this be improved?

# The Problem: better bounds

Lower bound:  $C(m, n) \geq n/m$

Can this be improved?



$$C(1, 13) = 13 = n/m$$

# The Problem: better bounds

Lower bound:  $C(m, n) \geq n/m$

Can this be improved?



$$C(1, 13) = 13 = n/m$$

If “aspect ratio”  $n/m$  is bounded, lower bound CAN be improved

# Electrical networks

Ohm's Law:


$$(\text{voltage drop}) = (\text{current}) \times (\text{resistance})$$

# Electrical networks

Ohm's Law:

$$(\text{voltage drop}) = (\text{current}) \times (\text{resistance})$$

Consider “current-voltage space”:




# Electrical networks

Ohm's Law:

$$(\text{voltage drop}) = (\text{current}) \times (\text{resistance})$$

Consider “current-voltage space”:






# Electrical networks

Ohm's Law:

$$(\text{voltage drop}) = (\text{current}) \times (\text{resistance})$$

Consider “current-voltage space”:




# Electrical networks

Ohm's Law:

$$(\text{voltage drop}) = (\text{current}) \times (\text{resistance})$$

Consider “current-voltage space”:




# Electrical networks

Ohm's Law:

$$(\text{voltage drop}) = (\text{current}) \times (\text{resistance})$$

Consider “current-voltage space”:



# Electrical networks

Ohm's Law:

$$(\text{voltage drop}) = (\text{current}) \times (\text{resistance})$$

Consider “current-voltage space”:

Square tiles  $\leftrightarrow$  unit resistors in IV-space!

# A Lower Bound

## Theorem (Kenyon)

*Let  $C(m, n)$  be the minimal cost of square-tiling a floor of size  $m$  by  $n$ , where  $m$  and  $n$  have no common factor. Then*

$$C(m, n) \geq \log_2(n)$$

# References



R. L. Brooks, C. A. B. Smith, A. H. Stone, and W. T. Tutte (1940)

The Dissection of Rectangles into Squares

*Duke Math. J.*, **7**, no. 1, pp. 312–340.



Richard Kenyon (1996)

Tiling a Rectangle with the Fewest Squares

*J. Combin. Theory Ser. A*, **76**, pp. 272–291.




FIG. 1

Thank you!