



# Dilated floor functions and their commutators

Harry Richman

joint w/ Jeff Lagarias and Takumi Murayama  
University of Michigan

December 5, 2018



# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

Examples:

①  $\lfloor 20.18 \rfloor =$

②  $\lfloor 1.99 \rfloor =$

③  $\lfloor -1.2 \rfloor =$

④  $\lfloor 5 \rfloor =$

# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

Examples:

①  $\lfloor 20.18 \rfloor = 20$

②  $\lfloor 1.99 \rfloor =$

③  $\lfloor -1.2 \rfloor =$

④  $\lfloor 5 \rfloor =$

# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

Examples:

①  $\lfloor 20.18 \rfloor = 20$

②  $\lfloor 1.99 \rfloor = 1$

③  $\lfloor -1.2 \rfloor =$

④  $\lfloor 5 \rfloor =$

# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

Examples:

①  $\lfloor 20.18 \rfloor = 20$

②  $\lfloor 1.99 \rfloor = 1$

③  $\lfloor -1.2 \rfloor = -2$

④  $\lfloor 5 \rfloor =$

# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

Examples:

①  $\lfloor 20.18 \rfloor = 20$

②  $\lfloor 1.99 \rfloor = 1$

③  $\lfloor -1.2 \rfloor = -2$

④  $\lfloor 5 \rfloor = 5$

# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

Examples:

①  $\lfloor 20.18 \rfloor = 20$

②  $\lfloor 1.99 \rfloor = 1$

③  $\lfloor -1.2 \rfloor = -2$

④  $\lfloor 5 \rfloor = 5$

$\lfloor x \rfloor$  is also known as the “greatest integer function”:

$$\lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\}$$



# Floor functions

The **floor function** rounds a real number down to the nearest integer

$$\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$




Figure: Graph of  $f(x) = \lfloor x \rfloor$

# Floor functions

A **dilated floor function** rounds down after rescaling by parameter  $\alpha$

$$f_{\alpha}(x) := \lfloor \alpha x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

# Floor functions

A **dilated floor function** rounds down after rescaling by parameter  $\alpha$

$$f_\alpha(x) := \lfloor \alpha x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$




Figure: Graph of  $f_\varphi(x) = \lfloor \varphi x \rfloor$ , where  $\varphi = \frac{1+\sqrt{5}}{2}$

# Floor functions

A **dilated floor function** rounds down after rescaling by parameter  $\alpha$

$$f_\alpha(x) := \lfloor \alpha x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$$

$\rightsquigarrow f_\alpha$  discretizes  $\mathbb{R}$   
“at length scale  $\alpha^{-1}$ ”




Figure: Graph of  $f_\varphi(x) = \lfloor \varphi x \rfloor$ , where  $\varphi = \frac{1+\sqrt{5}}{2}$

# Dilated floor functions: Why care?

Elementary number theory:

$$\text{val}_p(n!) = \left\lfloor \frac{1}{p} n \right\rfloor + \left\lfloor \frac{1}{p^2} n \right\rfloor + \left\lfloor \frac{1}{p^3} n \right\rfloor + \cdots$$

# Dilated floor functions: Why care?

Elementary number theory:

$$\text{val}_p(n!) = \left\lfloor \frac{1}{p}n \right\rfloor + \left\lfloor \frac{1}{p^2}n \right\rfloor + \left\lfloor \frac{1}{p^3}n \right\rfloor + \cdots$$

Note:  $\text{val}_p(M)$  = largest exponent  $e$  such that  $p^e$  divides  $M$

# Dilated floor functions: Why care?

Elementary number theory:

$$\text{val}_p(n!) = \left\lfloor \frac{1}{p}n \right\rfloor + \left\lfloor \frac{1}{p^2}n \right\rfloor + \left\lfloor \frac{1}{p^3}n \right\rfloor + \cdots$$

Riemann zeta function:

$$\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s} \quad \text{satisfies} \quad \int_0^\infty \lfloor x \rfloor x^{-s} \frac{dx}{x} = \frac{1}{s} \zeta(s)$$

# Dilated floor functions: Why care?

Elementary number theory:

$$\text{val}_p(n!) = \left\lfloor \frac{1}{p}n \right\rfloor + \left\lfloor \frac{1}{p^2}n \right\rfloor + \left\lfloor \frac{1}{p^3}n \right\rfloor + \cdots$$

Riemann zeta function:

$$\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s} \quad \text{satisfies} \quad \int_0^\infty \lfloor \alpha x \rfloor x^{-s} \frac{dx}{x} = \frac{\alpha^s}{s} \zeta(s)$$



# Dilated floor functions: Why care?

Elementary number theory:

$$\text{val}_p(n!) = \left\lfloor \frac{1}{p}n \right\rfloor + \left\lfloor \frac{1}{p^2}n \right\rfloor + \left\lfloor \frac{1}{p^3}n \right\rfloor + \cdots$$

Riemann zeta function:

$$\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s} \quad \text{satisfies} \quad \int_0^\infty \lfloor \alpha x \rfloor x^{-s} \frac{dx}{x} = \frac{\alpha^s}{s} \zeta(s)$$

i.e.  $-\frac{\alpha^{-s}}{s} \zeta(-s)$  is the Mellin transform of  $f_\alpha(x) = \lfloor \alpha x \rfloor$

# Dilated floor functions: Why care?

Elementary number theory:

$$\mathrm{val}_p(n!) = \left\lfloor \frac{1}{p}n \right\rfloor + \left\lfloor \frac{1}{p^2}n \right\rfloor + \left\lfloor \frac{1}{p^3}n \right\rfloor + \cdots$$

Riemann zeta function:

$$\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s} \quad \text{satisfies} \quad \int_0^\infty \lfloor \alpha x \rfloor x^{-s} \frac{dx}{x} = \frac{\alpha^s}{s} \zeta(s)$$

i.e.  $-\frac{\alpha^{-s}}{s} \zeta(-s)$  is the Mellin transform of  $f_\alpha(x) = \lfloor \alpha x \rfloor$

Algebraic geometry: measuring singularities, minimal model program...

# Dilated floor functions: Why care?

Elementary number theory:

$$\begin{aligned} \text{val}_p(n!) &= \left\lfloor \frac{1}{p}n \right\rfloor + \left\lfloor \frac{1}{p^2}n \right\rfloor + \left\lfloor \frac{1}{p^3}n \right\rfloor + \cdots \\ &\stackrel{?}{=} \left\lfloor \frac{1}{p}n \right\rfloor + \left\lfloor \frac{1}{p} \left\lfloor \frac{1}{p}n \right\rfloor \right\rfloor + \left\lfloor \frac{1}{p} \left\lfloor \frac{1}{p} \left\lfloor \frac{1}{p}n \right\rfloor \right\rfloor \right\rfloor + \cdots \end{aligned}$$

Riemann zeta function:

$$\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s} \quad \text{satisfies} \quad \int_0^\infty \lfloor \alpha x \rfloor x^{-s} \frac{dx}{x} = \frac{\alpha^s}{s} \zeta(s)$$

i.e.  $-\frac{\alpha^{-s}}{s} \zeta(-s)$  is the Mellin transform of  $f_\alpha(x) = \lfloor \alpha x \rfloor$

Algebraic geometry: measuring singularities, minimal model program...

# Dilated floor functions: Why care?

Economics: currency exchange

one US dollar = 0.80 UK pound (approximately)

= 1.33 Can. dollar

= 0.88 Eur. euro

# Dilated floor functions: Why care?

Economics: currency exchange

one US dollar =  $\frac{4}{5}$  UK pound (approximately)

=  $\frac{4}{3}$  Can. dollar

=  $\frac{8}{9}$  Eur. euro

# Dilated floor functions: Why care?

Economics: currency exchange

one US dollar  $= \frac{4}{5}$  UK pound (approximately)

$= \frac{4}{3}$  Can. dollar

$= \frac{8}{9}$  Eur. euro

To exchange money from US to UK

$$x \mapsto \left\lfloor \frac{4}{5}x \right\rfloor$$

(bank does not give back change)


# Dilated floor functions: Why care?

Economics: currency exchange

one US dollar =  $\frac{4}{5}$  UK pound (approximately)


=  $\frac{4}{3}$  Can. dollar

=  $\frac{8}{9}$  Eur. euro



# Dilated floor functions: Why care?


Economics: currency exchange





# Dilated floor functions: Why care?


Economics: currency exchange



Problem: Would you rather exchange money through Canada or Europe?

# Dilated floor functions: Why care?

Economics: currency exchange



Problem: Would you rather exchange money through Canada or Europe?

$$\left[ \frac{3}{5} \left[ \frac{4}{3} x \right] \right] \quad \text{vs.} \quad \left[ \frac{9}{10} \left[ \frac{8}{9} x \right] \right] ?$$

# Composing floor functions

## Vague Question

What happens when we compose  $f_\alpha$  and  $f_\beta$ ?

# Composing floor functions

## Vague Question

What happens when we compose  $f_\alpha$  and  $f_\beta$ ?




Figure: Graph of  $f_1 \circ f_\varphi = \lfloor \lfloor \varphi x \rfloor \rfloor$  where  $\varphi = \frac{1+\sqrt{5}}{2}$

# Composing floor functions

## Vague Question

What happens when we compose  $f_\alpha$  and  $f_\beta$ ?






Figure: Graph of  $f_\varphi \circ f_1 = \lfloor \varphi \lfloor x \rfloor \rfloor$  where  $\varphi = \frac{1+\sqrt{5}}{2}$

# Composing floor functions

## Vague Question

What happens when we compose  $f_\alpha$  and  $f_\beta$ ?

Example:  $f_1 \circ f_\varphi$  vs  $f_\varphi \circ f_1$




Observations:

# Composing floor functions

## Vague Question

What happens when we compose  $f_\alpha$  and  $f_\beta$ ?

Example:  $f_1 \circ f_\varphi$  vs  $f_\varphi \circ f_1$




Observations:

# Composing floor functions

## Vague Question

What happens when we compose  $f_\alpha$  and  $f_\beta$ ?

Example:  $f_1 \circ f_\varphi$  vs  $f_\varphi \circ f_1$



Observations:  $f_1 \circ f_\varphi \neq f_\varphi \circ f_1$




# Composing floor functions

## Vague Question

What happens when we compose  $f_\alpha$  and  $f_\beta$ ?

Example:  $f_1 \circ f_\varphi$  vs  $f_\varphi \circ f_1$




Observations:

•  $f_1 \circ f_\varphi \neq f_\varphi \circ f_1$

•  $f_1 \circ f_\varphi \geq f_\varphi \circ f_1$

# Composing floor functions

Example:  $f_1 \circ f_\varphi$  vs  $f_\varphi \circ f_1$



Observations:

- $f_1 \circ f_\varphi \neq f_\varphi \circ f_1$

- $f_1 \circ f_\varphi \geq f_\varphi \circ f_1$


## Problem A

For which  $(\alpha, \beta)$  do we have

$$f_\alpha \circ f_\beta = f_\beta \circ f_\alpha?$$

# Composing floor functions

Example:  $f_1 \circ f_\varphi$  vs  $f_\varphi \circ f_1$



Observations:  $\bullet f_1 \circ f_\varphi \neq f_\varphi \circ f_1$

$\bullet f_1 \circ f_\varphi \geq f_\varphi \circ f_1$

## Problem A

For which  $(\alpha, \beta)$  do we have

$$f_\alpha \circ f_\beta = f_\beta \circ f_\alpha?$$

## Problem B

For which  $(\alpha, \beta)$  do we have

$$f_\alpha \circ f_\beta \geq f_\beta \circ f_\alpha?$$

# Composing floor functions: results

## Problem A

For which  $(\alpha, \beta)$  do we have  $f_\alpha \circ f_\beta = f_\beta \circ f_\alpha$ ?

# Composing floor functions: results

## Problem A

For which  $(\alpha, \beta)$  do we have  $\lfloor \alpha \lfloor \beta x \rfloor \rfloor = \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ ?


# Composing floor functions: results

## Problem A

For which  $(\alpha, \beta)$  do we have  $\lfloor \alpha \lfloor \beta x \rfloor \rfloor = \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ ?

## Theorem (Lagarias–Murayama–R)

*All solutions to (A) are:*



# Composing floor functions: results


## Problem A

For which  $(\alpha, \beta)$  do we have  $\lfloor \alpha \lfloor \beta x \rfloor \rfloor = \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ ?

## Theorem (Lagarias–Murayama–R)

All solutions to (A) are:

$$\alpha = \frac{1}{m}, \beta = \frac{1}{n}$$



In fact:  $\lfloor \frac{1}{m} \lfloor \frac{1}{n} x \rfloor \rfloor = \lfloor \frac{1}{mn} x \rfloor$

# Composing floor functions: results


## Problem A

For which  $(\alpha, \beta)$  do we have  $\lfloor \alpha \lfloor \beta x \rfloor \rfloor = \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ ?

## Theorem (Lagarias–Murayama–R)

All solutions to (A) are:

$$\alpha = \frac{1}{m}, \beta = \frac{1}{n}$$



In fact:  $\left\lfloor \frac{1}{m} \left\lfloor \frac{1}{n} x \right\rfloor \right\rfloor = \left\lfloor \frac{1}{mn} x \right\rfloor$

$$\left\lfloor \frac{1}{p^2} n \right\rfloor = \left\lfloor \frac{1}{p} \left\lfloor \frac{1}{p} n \right\rfloor \right\rfloor !$$



# Composing floor functions: results

## Problem B

For which  $(\alpha, \beta)$  do we have  $f_\alpha \circ f_\beta \geq f_\beta \circ f_\alpha$ ?

# Composing floor functions: results

## Problem B

For which  $(\alpha, \beta)$  do we have  $\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ ?


# Composing floor functions: results

## Problem B


For which  $(\alpha, \beta)$  do we have  $\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ ?

## Theorem (Lagarias–R)

*All solutions to (B) are:*




# Composing floor functions: results



$\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ : positive-dilation results


## Theorem (Lagarias–R)

*All positive solutions to (B) are:*



# $\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ : positive-dilation results


Coordinate change:




# $\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ : positive-dilation results

Coordinate change:

$$\mu = \frac{1}{\alpha}, \nu = \frac{\beta}{\alpha}$$




$\rightsquigarrow$




# $\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ : positive-dilation results

Coordinate change:


$$\mu = \frac{1}{\alpha}, \nu = \frac{\beta}{\alpha}$$



$\rightsquigarrow$



Symmetries:







# $\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ : positive-dilation results

Coordinate change:


$$\mu = \frac{1}{\alpha}, \nu = \frac{\beta}{\alpha}$$




$\rightsquigarrow$



Symmetries:




$\rightsquigarrow$



$\lfloor \alpha \lfloor \beta x \rfloor \rfloor \geq \lfloor \beta \lfloor \alpha x \rfloor \rfloor$ : positive-dilation results

Where do **green** solution curves come from?



# Proof ingredient: Beatty sequences

Parameter  $\mu \geq 1$ ,

$$\mathcal{B}(\mu) = \{\lfloor \mu \rfloor, \lfloor 2\mu \rfloor, \lfloor 3\mu \rfloor, \dots\} \subset \mathbb{N}$$

# Proof ingredient: Beatty sequences

Parameter  $\mu \geq 1$ ,

$$\mathcal{B}(\mu) = \{\lfloor \mu \rfloor, \lfloor 2\mu \rfloor, \lfloor 3\mu \rfloor, \dots\} \subset \mathbb{N}$$

Note:

$$\mathcal{B}(\mu) = \text{output values of } f_\mu \circ f_1(x) = \lfloor \mu \lfloor x \rfloor \rfloor$$

# Proof ingredient: Beatty sequences

Parameter  $\mu \geq 1$ ,

$$\mathcal{B}(\mu) = \{\lfloor \mu \rfloor, \lfloor 2\mu \rfloor, \lfloor 3\mu \rfloor, \dots\} \subset \mathbb{N}$$

Note:

$$\mathcal{B}(\mu) = \text{output values of } f_\mu \circ f_1(x) = \lfloor \mu \lfloor x \rfloor \rfloor$$

Example:  $\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618$ ,  $\mathcal{B}(\varphi) = \{1, 3, 4, 6, 8, 9, 11, 12, \dots\}$

# Proof ingredient: Beatty sequences

Parameter  $\mu \geq 1$ ,

$$\mathcal{B}(\mu) = \{\lfloor \mu \rfloor, \lfloor 2\mu \rfloor, \lfloor 3\mu \rfloor, \dots\} \subset \mathbb{N}$$

Note:

$$\mathcal{B}(\mu) = \text{output values of } f_\mu \circ f_1(x) = \lfloor \mu \lfloor x \rfloor \rfloor$$

Example:  $\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618, \quad \mathcal{B}(\varphi) = \{1, 3, 4, 6, 8, 9, 11, 12, \dots\}$   
 $\varphi^2 = \frac{3+\sqrt{5}}{2} \approx 2.618, \quad \mathcal{B}(\varphi^2) = \{2, 5, 7, 10, 13, 15, 18, \dots\}$

# Proof ingredient: Beatty sequences

Parameter  $\mu \geq 1$ ,

$$\mathcal{B}(\mu) = \{\lfloor \mu \rfloor, \lfloor 2\mu \rfloor, \lfloor 3\mu \rfloor, \dots\} \subset \mathbb{N}$$

Note:

$$\mathcal{B}(\mu) = \text{output values of } f_\mu \circ f_1(x) = \lfloor \mu \lfloor x \rfloor \rfloor$$

Example:  $\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618, \quad \mathcal{B}(\varphi) = \{1, 3, 4, 6, 8, 9, 11, 12, \dots\}$   
 $\varphi^2 = \frac{3+\sqrt{5}}{2} \approx 2.618, \quad \mathcal{B}(\varphi^2) = \{2, 5, 7, 10, 13, 15, 18, \dots\}$

What is “special” about  $\mathcal{B}(\varphi)$  and  $\mathcal{B}(\varphi^2)$ ?

# Proof ingredient: Beatty sequences

Parameter  $\mu \geq 1$ ,

$$\mathcal{B}(\mu) = \{\lfloor \mu \rfloor, \lfloor 2\mu \rfloor, \lfloor 3\mu \rfloor, \dots\} \subset \mathbb{N}$$

Note:

$$\mathcal{B}(\mu) = \text{output values of } f_\mu \circ f_1(x) = \lfloor \mu \lfloor x \rfloor \rfloor$$

Example:  $\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618$ ,  $\mathcal{B}(\varphi) = \{1, 3, 4, 6, 8, 9, 11, 12, \dots\}$   
 $\varphi^2 = \frac{3+\sqrt{5}}{2} \approx 2.618$ ,  $\mathcal{B}(\varphi^2) = \{2, 5, 7, 10, 13, 15, 18, \dots\}$

Theorem (“Beatty’s Theorem,” Ostrowski, Hyslop, Aitken, ..)

If  $\mu$  and  $\nu$  are irrational and satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ , then

$$\mathcal{B}(\mu) \cap \mathcal{B}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}(\nu) = \mathbb{N}$$

i.e. their Beatty sequences **partition**  $\mathbb{N}$ .



# Proof ingredient: Beatty sequences

## Theorem (“Beatty’s Theorem” 1926)

*Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}(\nu)$  partition  $\mathbb{N}$ , i.e.*

$$\mathcal{B}(\mu) \cap \mathcal{B}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}(\nu) = \mathbb{N}$$

**iff**  $\mu$  and  $\nu$  are irrational and satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

# Proof ingredient: Beatty sequences

## Theorem (“Beatty’s Theorem” 1926)

*Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}(\nu)$  partition  $\mathbb{N}$ , i.e.*

$$\mathcal{B}(\mu) \cap \mathcal{B}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}(\nu) = \mathbb{N}$$

**iff**  $\mu$  and  $\nu$  are *irrational* and satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

# Proof ingredient: Beatty sequences

## Theorem (“Beatty’s Theorem” 1926)

Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}(\nu) = \mathbb{N}$$

iff  $\mu$  and  $\nu$  are *irrational* and satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

## Proposition 1 (Lagarias–R)

Modified\* Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}^{<}(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}^{<}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}^{<}(\nu) = \mathbb{N}$$

iff  $\mu$  and  $\nu$  satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

# Proof ingredient: Beatty sequences

## Theorem (“Beatty’s Theorem” 1926)

Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}(\nu) = \mathbb{N}$$

iff  $\mu$  and  $\nu$  are *irrational* and satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

## Proposition 1 (Lagarias–R)

Modified\* Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}^{<}(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}^{<}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}^{<}(\nu) = \mathbb{N}$$

iff  $\mu$  and  $\nu$  satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

Idea: “break ties” between  $\mu\mathbb{N}$  and  $\nu\mathbb{N}$

# Proof ingredient: Beatty sequences

## Theorem (“Beatty’s Theorem” 1926)

Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}(\nu) = \mathbb{N}$$

iff  $\mu$  and  $\nu$  are *irrational* and satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

## Proposition 1 (Lagarias–R)

Modified\* Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}^{<}(\nu)$  partition  $\mathbb{Z}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}^{<}(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}^{<}(\nu) = \mathbb{Z}$$

iff  $\mu$  and  $\nu$  satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

Idea: “break ties” between  $\mu\mathbb{Z}$  and  $\nu\mathbb{Z}$

# Proof ingredient: Beatty sequences

## Proposition 1 (Lagarias–R)

Modified\* Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}^<(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}^<(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}^<(\nu) = \mathbb{N}$$

**iff**  $\mu$  and  $\nu$  satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

Idea: “break ties” between  $\mu\mathbb{N}$  and  $\nu\mathbb{N}$

# Proof ingredient: Beatty sequences

## Proposition 1 (Lagarias–R)

Modified\* Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}^<(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}^<(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}^<(\nu) = \mathbb{N}$$

**iff**  $\mu$  and  $\nu$  satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

Idea: “break ties” between  $\mu\mathbb{N}$  and  $\nu\mathbb{N}$

## Proposition 2 (Lagarias–R)

For parameters  $(\alpha, \beta) > 0$ ,

$$f_\alpha \circ f_\beta \geq f_\beta \circ f_\alpha \quad \text{iff} \quad \mathcal{B}(\mu) \cap \mathcal{B}^<(\nu) = \emptyset$$

where  $\mu = \frac{1}{\alpha}$  and  $\nu = \frac{\beta}{\alpha}$ .

# Proof ingredient: Beatty sequences

## Proposition 1 (Lagarias–R)

Modified\* Beatty sequences  $\mathcal{B}(\mu)$ ,  $\mathcal{B}^<(\nu)$  partition  $\mathbb{N}$ , i.e.

$$\mathcal{B}(\mu) \cap \mathcal{B}^<(\nu) = \emptyset \quad \text{and} \quad \mathcal{B}(\mu) \cup \mathcal{B}^<(\nu) = \mathbb{N}$$

iff  $\mu$  and  $\nu$  satisfy  $\frac{1}{\mu} + \frac{1}{\nu} = 1$ .

Idea: “break ties” between  $\mu\mathbb{N}$  and  $\nu\mathbb{N}$

## Proposition 2 (Lagarias–R)

For parameters  $(\alpha, \beta) > 0$ ,


$$f_\alpha \circ f_\beta \geq f_\beta \circ f_\alpha \quad \text{iff} \quad \mathcal{B}(\mu) \cap \mathcal{B}^<(\nu) = \emptyset$$

where  $\mu = \frac{1}{\alpha}$  and  $\nu = \frac{\beta}{\alpha}$ .




# Proof ingredient: Beatty sequences

$\rightsquigarrow$  **Green** solution curves come from Beatty sequences



# Proof ingredient: Beatty sequences

↪ **Green** solution curves come from Beatty sequences



How do we know there are **no more** solutions?

# Proof ingredient: Torus subgroups

Torus surface  $\mathbb{T} = \mathbb{R}^2 / \mathbb{Z}^2$


A point  $(\sigma, \tau) \in \mathbb{T}$  generates a **cyclic subgroup** of  $\mathbb{T}$

# Proof ingredient: Torus subgroups

Torus surface  $\mathbb{T} = \mathbb{R}^2 / \mathbb{Z}^2$

A point  $(\sigma, \tau) \in \mathbb{T}$  generates a **cyclic subgroup** of  $\mathbb{T}$

Example:




# Proof ingredient: Torus subgroups

Torus surface  $\mathbb{T} = \mathbb{R}^2 / \mathbb{Z}^2$

A point  $(\sigma, \tau) \in \mathbb{T}$  generates a **cyclic subgroup** of  $\mathbb{T}$

Example:



## Vague Question


When is  $(\sigma, \tau) \in \mathbb{T}$  a “minimal” generator for its subgroup?

# Proof ingredient: Torus subgroups

Torus surface  $\mathbb{T} = \mathbb{R}^2 / \mathbb{Z}^2$

A point  $(\sigma, \tau) \in \mathbb{T}$  generates a **cyclic subgroup** of  $\mathbb{T}$

Example:



## Vague Question


When is  $(\sigma, \tau) \in \mathbb{T}$  a “minimal” generator for its subgroup?

# Proof ingredient: Torus subgroups

Torus surface  $\mathbb{T} = \mathbb{R}^2 / \mathbb{Z}^2$

A point  $(\sigma, \tau) \in \mathbb{T}$  generates a **cyclic subgroup** of  $\mathbb{T}$

Def.  $(\sigma, \tau)$  is **weakly minimal** if



## Vague Question


When is  $(\sigma, \tau) \in \mathbb{T}$  a “minimal” generator for its subgroup?

# Proof ingredient: Torus subgroups

Torus surface  $\mathbb{T} = \mathbb{R}^2 / \mathbb{Z}^2$

A point  $(\sigma, \tau) \in \mathbb{T}$  generates a **cyclic subgroup** of  $\mathbb{T}$

Def.  $(\sigma, \tau)$  is **strongly minimal** if



## Vague Question

When is  $(\sigma, \tau) \in \mathbb{T}$  a “minimal” generator for its subgroup?




# Proof ingredient: Torus subgroups


## Vague Question

When is  $(\sigma, \tau) \in \mathbb{T}$  a “minimal” generator for its subgroup?

$(\sigma, \tau)$  **weakly** minimal



$(\sigma, \tau)$  **strongly** minimal




# Proof ingredient: Torus subgroups


## Vague Question

When is  $(\sigma, \tau) \in \mathbb{T}$  a “minimal” generator for its subgroup?

$(\sigma, \tau)$  **weakly** minimal



$(\sigma, \tau)$  **strongly** minimal



## Proposition 3 (Lagarias–R)


If  $(\sigma, \tau)$  is a weakly minimal generator, it is also strongly minimal.

# Proof ingredient: Torus subgroups

## Proposition 3 (Lagarias–R)

If  $(\sigma, \tau)$  is a weakly minimal generator, it is also strongly minimal.

All minimal generators of cyclic subgroups, in  $\mathbb{T}$ :



# Composing floor functions: Why **really** care?

# Composing floor functions: Why **really** care?

(Recall: P.N.T. says  $\pi(x) = \#\{\text{prime } p \leq x\} \approx \frac{x}{\log x}$  )

**Riemann hypothesis**:  $\Pi(x) := \sum_{p \leq x} \log p$

$$\text{R. H.} \quad \Leftrightarrow \quad \Pi(x) = x + O(x^{1/2+\epsilon})$$

# Composing floor functions: Why **really** care?

(Recall: P.N.T. says  $\pi(x) = \#\{\text{prime } p \leq x\} \approx \frac{x}{\log x}$  )

Riemann hypothesis:  $\Pi(x) := \sum_{p \leq x} \log p$

$$\text{R. H.} \quad \Leftrightarrow \quad \Pi(x) = x + O(x^{1/2+\epsilon})$$

Mertens function:

$$M(x) := \sum_{n \leq x} \mu(n) \quad \text{where } \mu(n) \in \{\pm 1, 0\} \text{ is the Möbius function}$$

$$\text{R. H.} \quad \Leftrightarrow \quad M(x) = O(x^{1/2+\epsilon})$$

# Composing floor functions: Why **really** care?

(Recall: P.N.T. says  $\pi(x) = \#\{\text{prime } p \leq x\} \approx \frac{x}{\log x}$  )

Riemann hypothesis:  $\Pi(x) := \sum_{p \leq x} \log p$

$$\text{R. H.} \quad \Leftrightarrow \quad \Pi(x) = x + O(x^{1/2+\epsilon})$$

Mertens function:

$$M(x) := \sum_{n \leq x} \mu(n) \quad \text{where } \mu(n) \in \{\pm 1, 0\} \text{ is the Möbius function}$$

$$\text{R. H.} \quad \Leftrightarrow \quad M(x) = O(x^{1/2+\epsilon})$$

Jean-Paul Cardinal (2010) defined a “2-dimensional analogue” of the Mertens function

# Composing floor functions: Why **really** care?

Let  $\{d_i\} = \{n, \lfloor \frac{1}{2}n \rfloor, \lfloor \frac{1}{3}n \rfloor, \lfloor \frac{1}{4}n \rfloor, \dots, 1\}$  be the “almost divisors” of  $n$ .

In Cardinal’s matrix  $\mathcal{M}_n$ , the entry in position  $i, j$  is

$$\mathcal{M}_n(i, j) = M\left(\left\lfloor \frac{1}{d_i d_j} n \right\rfloor\right) = M\left(\left\lfloor \frac{1}{d_i} \left\lfloor \frac{1}{d_j} n \right\rfloor \right\rfloor\right) = M\left(\left\lfloor \frac{1}{d_j} \left\lfloor \frac{1}{d_i} n \right\rfloor \right\rfloor\right)$$

## Theorem (Cardinal 2010)

*Riemann hypothesis is equivalent to*

$$\|\mathcal{M}_n\| = O(n^{1/2+\epsilon}) \quad \text{as } n \rightarrow \infty.$$



# Composing floor functions: Why **really** care?

Let  $\{d_i\} = \{n, \lfloor \frac{1}{2}n \rfloor, \lfloor \frac{1}{3}n \rfloor, \lfloor \frac{1}{4}n \rfloor, \dots, 1\}$  be the “almost divisors” of  $n$ .

In Cardinal's matrix  $\mathcal{M}_n$ , the entry in position  $i, j$  is

$$\mathcal{M}_n(i, j) = M\left(\left\lfloor \frac{1}{d_i d_j} n \right\rfloor\right) = M\left(\left\lfloor \frac{1}{d_i} \left\lfloor \frac{1}{d_j} n \right\rfloor \right\rfloor\right) = M\left(\left\lfloor \frac{1}{d_j} \left\lfloor \frac{1}{d_i} n \right\rfloor \right\rfloor\right)$$

## Theorem (Cardinal 2010)

*Riemann hypothesis is equivalent to*

$$\|\mathcal{M}_n\| = O(n^{1/2+\epsilon}) \quad \text{as } n \rightarrow \infty.$$

Computational evidence suggests that  $\|\mathcal{M}_n\|$  is better behaved than Mertens function  $M(n)$  as  $n \rightarrow \infty \dots$

# Composing floor functions: Why **really** care?

Let  $\{d_i\} = \{n, \lfloor \frac{1}{2}n \rfloor, \lfloor \frac{1}{3}n \rfloor, \lfloor \frac{1}{4}n \rfloor, \dots, 1\}$  be the “almost divisors” of  $n$ .

In Cardinal's matrix  $\mathcal{M}_n$ , the entry in position  $i, j$  is

$$\mathcal{M}_n(i, j) = M\left(\left\lfloor \frac{1}{d_i d_j} n \right\rfloor\right) = M\left(\left\lfloor \frac{1}{d_i} \left\lfloor \frac{1}{d_j} n \right\rfloor \right\rfloor\right) = M\left(\left\lfloor \frac{1}{d_j} \left\lfloor \frac{1}{d_i} n \right\rfloor \right\rfloor\right)$$

( Note: “almost divisors of almost divisors are almost divisors”! )

Theorem (Cardinal 2010)

*Riemann hypothesis is equivalent to*

$$\|\mathcal{M}_n\| = O(n^{1/2+\epsilon}) \quad \text{as } n \rightarrow \infty.$$

Computational evidence suggests that  $\|\mathcal{M}_n\|$  is better behaved than Mertens function  $M(n)$  as  $n \rightarrow \infty \dots$

# References



S. Beatty (1926)

Problem 3173

*Amer. Math. Monthly* **33**(3) 159.



J.-P. Cardinal (2010)

Symmetric matrices related to the Mertens function

*Lin. Alg. Appl.* **432**(1), 161–172.



J. C. Lagarias, T. Murayama, D. H. Richman (2016)

Dilated floor functions that commute


*Amer. Math. Monthly* **163**(10), [arXiv:1611.05513](#).



J. C. Lagarias and D. H. Richman (2018)

Dilated floor functions with nonnegative commutator I  
to appear in *Acta Arith.*, [arXiv:1806.00579](#).

# Dilated floor function commutators



Thank you!