

Weierstrass points on a tropical curve

Harry Richman

University of Michigan

hrichman@umich.edu

University of Washington AAG Seminar

29 October 2019

What is a Weierstrass point?

Definition: X a smooth algebraic curve, D_N a divisor of degree N
 \rightsquigarrow projective embedding $\phi : X \rightarrow \mathbb{P}^r$.

What is a Weierstrass point?

Definition: X a smooth algebraic curve, D_N a divisor of degree N
 \rightsquigarrow projective embedding $\phi : X \rightarrow \mathbb{P}^r$.


$$\begin{aligned} W(D_N) &= \{x \in X : \exists H \subset \mathbb{P}^r \text{ s.t. } m_x(H \cap X) \geq r + 1\} \\ &= \left\{ x \in X : \begin{array}{l} \text{“higher-than-expected” tangency with} \\ \text{some hyperplane } H \text{ at } x \end{array} \right\} \end{aligned}$$

What is a Weierstrass point?

Definition:

$$W(D_N) = \left\{ x \in X : \begin{array}{l} \text{"higher-than-expected" tangency with} \\ \text{some hyperplane } H \text{ at } x \end{array} \right\}$$

Example: $X = \{xyz + x^3 + y^3 + z^3 = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$



$N = 3$


What is a Weierstrass point?

Definition:

$$W(D_N) = \left\{ x \in X : \begin{array}{l} \text{“higher-than-expected” tangency with} \\ \text{some hyperplane } H \text{ at } x \end{array} \right\}$$

Example: $X = \{xyz + x^3 + y^3 + z^3 = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$

$N = 3$




What is a Weierstrass point?

Definition:

$$W(D_N) = \left\{ x \in X : \begin{array}{l} \text{“higher-than-expected” tangency with} \\ \text{some hyperplane } H \text{ at } x \end{array} \right\}$$

Example: $X = \{xyz + x^3 + y^3 + z^3 = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$

$N = 3$




What is a Weierstrass point?

Definition:

$$W(D_N) = \left\{ x \in X : \begin{array}{l} \text{“higher-than-expected” tangency with} \\ \text{some hyperplane } H \text{ at } x \end{array} \right\}$$

Example: $X = \{xyz + x^3 + y^3 + z^3 = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$



$$N = 3, \quad \#W(D) = 9$$

What is a Weierstrass point?

Intuition (Mumford):

N -torsion points
on an elliptic curve \leftrightarrow Weierstrass points of D_N
on a higher-genus curve

What is a Weierstrass point?

Intuition (Mumford):

N -torsion points
on an elliptic curve \leftrightarrow Weierstrass points of D_N
on a higher-genus curve

Numerical “evidence”: as N grows,

$$\#(\text{Weierstrass points of } D_N) = gN^2 + O(N)$$

What is a Weierstrass point?

Intuition (Mumford):

N -torsion points
on an elliptic curve \leftrightarrow Weierstrass points of D_N
on a higher-genus curve

Numerical “evidence”: as N grows,

$$\#(\text{Weierstrass points of } D_N) = gN^2 + O(N)$$

Problem

How are Weierstrass points distributed on an algebraic curve?

Weierstrass points: genus 1, complex case


Problem

How are Weierstrass points distributed on genus 1 curve X/\mathbb{C} ?

Weierstrass points: genus 1, complex case

Problem


How are Weierstrass points distributed on genus 1 curve X/\mathbb{C} ?



Weierstrass points: genus 1, complex case

Problem


How are Weierstrass points distributed on genus 1 curve X/\mathbb{C} ?



Weierstrass points: genus 1, complex case

Problem


How are Weierstrass points distributed on genus 1 curve X/\mathbb{C} ?



Weierstrass points: genus 1, complex case

Problem

How are Weierstrass points distributed on genus 1 curve X/\mathbb{C} ?



\rightsquigarrow Weierstrass points distribute **uniformly**, w.r.t. $\mathbb{C} \rightarrow \mathbb{C}/\Lambda$

Weierstrass points: genus ≥ 2 , complex case


Problem

How are Weierstrass points distributed on **higher** genus curve X/\mathbb{C} ?

Weierstrass points: genus ≥ 2 , complex case

Problem


How are Weierstrass points distributed on higher genus curve X/\mathbb{C} ?



Weierstrass points: genus ≥ 2 , complex case

Problem


How are Weierstrass points distributed on higher genus curve X/\mathbb{C} ?



Weierstrass points: genus ≥ 2 , complex case

Problem


How are Weierstrass points distributed on higher genus curve X/\mathbb{C} ?



Weierstrass points: genus ≥ 2 , complex case

Problem

How are Weierstrass points distributed on higher genus curve X/\mathbb{C} ?




Theorem (Neeman, 1984)

Suppose X is a complex algebraic curve of genus $g \geq 2$. Then $W(D_N)$ distributes according to the Bergman measure as $N \rightarrow \infty$.

Weierstrass points: non-Archimedean case

Problem

How are Weierstrass points distributed on X/\mathbb{K} , $val : \mathbb{K}^\times \rightarrow \mathbb{R}$?




Source: Matt Baker's math blog

Weierstrass points: non-Archimedean case

Problem

How are Weierstrass points distributed on X/\mathbb{K} X^{an} ?




Source: Matt Baker's math blog

Weierstrass points: non-Archimedean case

Problem

How are Weierstrass points distributed on X/\mathbb{K} X^{an} ?



Source: Matt Baker's math blog


Theorem (Amini, 2014)

Suppose X^{an} is a Berkovich curve of genus $g \geq 2$. Then $W(D_N)$ distributes according to the Zhang measure as $N \rightarrow \infty$.

Weierstrass points: non-Archimedean case

Problem

How are Weierstrass points distributed on X/\mathbb{K} X^{an} ?



Source: Matt Baker's math blog


Problem (Amini, 2014)

Does the distribution follow from considering only the skeleton $\Gamma \subset X^{\text{an}}$?

Weierstrass points: non-Archimedean case

Problem

How are Weierstrass points distributed on X/\mathbb{K} X^{an} ?




Source: Matt Baker's math blog

Problem (Amini, 2014)

Does the distribution follow from considering only the skeleton $\Gamma \subset X^{\text{an}}$?

What is a tropical curve?


Tropical curve (= a skeleton of X^{an})



What is a tropical curve?

Tropical curve (= a skeleton of X^{an})

(combinatorics) = finite graph with edge lengths




What is a tropical curve?

Tropical curve (= a skeleton of X^{an})

(combinatorics) = finite graph with edge lengths

(alg. geometry) = model for a degenerating algebraic curve




What is a tropical curve?

Tropical curve (= a skeleton of X^{an})

(combinatorics) = finite graph with edge lengths

(alg. geometry) = model for a degenerating algebraic curve

Example: $X_t = \{xyz - tx^3 + t^2y^3 + t^5z^3 = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$




What is a tropical curve?

Tropical curve (= a skeleton of X^{an})


(combinatorics) = finite graph with edge lengths

(alg. geometry) = model for a degenerating algebraic curve

Example: $X_t = \{xyz - t^1x^3 + t^2y^3 + t^5z^3 = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$



\rightsquigarrow
dual graph
of X_0




What is a tropical curve?

Tropical curve (= a skeleton of X^{an})


(combinatorics) = ~~finite graph with edge lengths~~ metric graph

(alg. geometry) = model for a degenerating algebraic curve

Example: $X_t = \{xyz - t^1x^3 + t^2y^3 + t^5z^3 = 0\} \subset \mathbb{P}_{\mathbb{C}}^2$



\rightsquigarrow
dual graph
of X_0



Tropical curves: divisor theory

Tropical curve = metric graph

alg. curve X		tropical curve Γ
divisors $\text{Div}(X)$	\rightsquigarrow	divisors $\text{Div}(\Gamma)$
meromorphic functions	\rightsquigarrow	piecewise \mathbb{Z} -linear functions
linear system $ D $	\rightsquigarrow	linear system $ D $
$= \mathbb{P}^r$		$=$ polyhedral complex of $\dim \geq r$
rank $r = \dim D $	\rightsquigarrow	rank $r =$ Baker-Norine rank

Tropical curves: divisor theory

Tropical curve = metric graph

alg. curve X		tropical curve Γ
divisors $\text{Div}(X)$	\rightsquigarrow	divisors $\text{Div}(\Gamma)$
meromorphic functions	\rightsquigarrow	piecewise \mathbb{Z} -linear functions
linear system $ D $	\rightsquigarrow	linear system $ D $
$= \mathbb{P}^r$		$=$ polyhedral complex of $\dim \geq r$
rank $r = \dim D $	\rightsquigarrow	rank $r =$ Baker-Norine rank

Intuition: linear equivalence on Γ = “discrete current flow”
 $|D| = \{E \text{ lin. equiv. to } D, E \geq 0\}$

Tropical curves: divisor theory

Tropical curve = metric graph

alg. curve X		tropical curve Γ
divisors $\text{Div}(X)$	\rightsquigarrow	divisors $\text{Div}(\Gamma)$
meromorphic functions	\rightsquigarrow	piecewise \mathbb{Z} -linear functions
linear system $ D $	\rightsquigarrow	linear system $ D $
$= \mathbb{P}^r$		$=$ polyhedral complex of $\dim \geq r$
rank $r = \dim D $	\rightsquigarrow	rank $r =$ Baker-Norine rank

Intuition: linear equivalence on Γ = “discrete current flow”

$$|D| = \{E \text{ lin. equiv. to } D, E \geq 0\}$$


q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




What happens as q varies?

Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:




Tropical curves: reduced divisors

Tropical curve = metric graph

Intuition: q -reduced divisor $\text{red}_q[D] =$ “energy-minimizing” divisor in $|D|$

Example:



Tropical curves: Weierstrass points

Problem

How are Weierstrass points distributed on a tropical curve?

Definition: Γ = metric graph, D_N divisor of degree N
 \rightsquigarrow Baker–Norine rank $r = r(D_N)$

$$W(D_N) = \{x \in X : \text{red}_x[D_N] \geq (r+1)x\}$$

Tropical curves: Weierstrass points

Problem

How are Weierstrass points distributed on a tropical curve?

Definition: Γ = metric graph, D_N divisor of degree N

\rightsquigarrow Baker–Norine rank $r = N - g$ when $N \gg 0$

$$W(D_N) = \{x \in X : \text{red}_x[D_N] \geq (r + 1)x\}$$

Tropical curves: Weierstrass points

Problem

How are Weierstrass points distributed on a tropical curve?

Definition: Γ = metric graph, D_N divisor of degree N

\rightsquigarrow Baker–Norine rank $r = N - g$ when $N \gg 0$

$$W(D_N) = \{x \in X : \text{red}_x[D_N] \geq (r + 1)x\}$$

Observation: as N grows,

$$\#(\text{Weierstrass points of } D_N) = gN + O(1) \quad \dots$$

Tropical curves: Weierstrass points

Problem

How are Weierstrass points distributed on a tropical curve?

Definition: Γ = metric graph, D_N divisor of degree N

\rightsquigarrow Baker–Norine rank $r = N - g$ when $N \gg 0$

$$W(D_N) = \{x \in X : \text{red}_x[D_N] \geq (r + 1)x\}$$


Observation: as N grows,

$$\#(\text{Weierstrass points of } D_N) = gN + O(1) \quad \dots$$

EXCEPT sometimes $\#(\text{Weierstrass points}) = \infty$

Weierstrass points: tropical case


Example: Genus $g(\Gamma) = 1$:



degree $D = 6$,

Weierstrass points: tropical case


Example: Genus $g(\Gamma) = 1$:



$$\text{degree } D = 6, \quad \rightsquigarrow \quad \#(W(D)) = 6$$

Weierstrass points: tropical case


Example: Genus $g(\Gamma) = 3$:



degree $D = 4$,

Weierstrass points: tropical case


Example: Genus $g(\Gamma) = 3$:



$$\text{degree } D = 4, \quad \rightsquigarrow \quad \#(W(D)) = 8$$

Weierstrass points: tropical case


Example: Genus $g(\Gamma) = 3$:



degree $D = 4$,

Weierstrass points: tropical case


Example: Genus $g(\Gamma) = 3$:



degree $D = 4$, \rightsquigarrow $\#(W(D)) = \infty!$

Weierstrass points: tropical case

Example: Genus $g(\Gamma) = 3$:



degree $D = 4$, \rightsquigarrow $\#(\mathbf{W}(D)) = \infty!$

Weierstrass points: tropical case

In general, this problem doesn't happen!

Theorem (R)

For a generic divisor class $[D]$, the Weierstrass locus $W(D)$ is finite.

Weierstrass points: tropical case

In general, this problem doesn't happen!

Theorem (R)

For a generic divisor class $[D]$, the Weierstrass locus $W(D)$ is finite.

So, we can still ask

Problem

How are Weierstrass points distributed **supposing** $W(D)$ is finite?

Weierstrass points: tropical case

In general, this problem doesn't happen!

Theorem (R)

For a generic divisor class $[D]$, the Weierstrass locus $W(D)$ is finite.

So, we can still ask

Problem

How are Weierstrass points distributed ~~supposing $W(D)$ is finite~~
for **generic** $[D]$?

Weierstrass points: tropical case

In general, this problem doesn't happen!

Theorem (R)

For a generic divisor class $[D]$, the Weierstrass locus $W(D)$ is finite.

So, we can still ask

Problem

How are Weierstrass points distributed ~~supposing $W(D)$ is finite~~
for **generic** $[D]$?

Theorem (R)

For a sequence of generic divisor classes $[D_N]$ on Γ , the Weierstrass locus $W(D_N)$ distributes according to Zhang's canonical measure μ .

Electrical networks

Γ = electrical network by replacing each edge \rightsquigarrow resistor

Electrical networks

Γ = electrical network by replacing each edge \rightsquigarrow resistor

Given $y, z \in \Gamma$, let

$$j_z^y = \left(\begin{array}{l} \text{voltage on } \Gamma \text{ when 1 unit of} \\ \text{current is sent from } y \text{ to } z \end{array} \right)$$

By Ohm's law, **current** = $\frac{\text{voltage}}{\text{resistance}}$ = **slope** of j_z^y

Electrical networks


Γ = electrical network by replacing each edge \rightsquigarrow resistor

Given $y, z \in \Gamma$, let

$$j_z^y = \left(\begin{array}{l} \text{voltage on } \Gamma \text{ when 1 unit of} \\ \text{current is sent from } y \text{ to } z \end{array} \right)$$


By Ohm's law, **current** = $\frac{\text{voltage}}{\text{resistance}} = \mathbf{slope}$ of j_z^y

Example: current = $(j_z^y)'$




Electrical networks

Example: current = $(j_z^y)'$



satisfies Laplacian $\Delta(j_z^y) = z - y$



Electrical networks: Canonical measure

Γ = metric graph

Definition (“electrical” version, Chinburg–Rumely–Baker–Faber)

Zhang’s **canonical measure** μ on an edge is the “current defect”

$$\begin{aligned}\mu(e) &= \text{current bypassing } e \text{ when 1 unit sent from } e^- \text{ to } e^+ \\ &= 1 - (\text{current through } e \text{ when ...})\end{aligned}$$

Electrical networks: Canonical measure


Γ = metric graph

Definition (“electrical” version, Chinburg–Rumely–Baker–Faber)

Zhang’s **canonical measure** μ on an edge is the “current defect”

$$\begin{aligned}\mu(e) &= \text{current bypassing } e \text{ when 1 unit sent from } e^- \text{ to } e^+ \\ &= 1 - (\text{current through } e \text{ when } \dots)\end{aligned}$$

Example:



Electrical networks: Canonical measure

Γ = metric graph

Definition (“electrical” version, Chinburg–Rumely–Baker–Faber)

Zhang’s **canonical measure** μ on an edge is the “current defect”

$$\begin{aligned}\mu(e) &= \text{current bypassing } e \text{ when 1 unit sent from } e^- \text{ to } e^+ \\ &= 1 - (\text{current through } e \text{ when ...})\end{aligned}$$

Generally:

- $0 \leq \mu(e) \leq 1$
- $\mu(e) = 0 \Leftrightarrow e$ a bridge
- $\mu(e) = 1 \Leftrightarrow e$ a loop

Foster’s Theorem: $\mu(\Gamma) = \sum_{e \in E} \mu(e) = g$

Tropical Weierstrass distribution: proof idea

Theorem (R)

For a sequence of generic divisor classes $[D_N]$ on Γ , the Weierstrass locus $W(D_N)$ distributes according to Zhang's canonical measure μ .

Namely, for any edge e

$$\frac{\#(W(D_N) \cap e)}{N} \rightarrow \mu(e) \quad \text{as} \quad N \rightarrow \infty.$$

Tropical Weierstrass distribution: proof idea

Theorem (R)

For a sequence of generic divisor classes $[D_N]$ on Γ , the Weierstrass locus $W(D_N)$ distributes according to Zhang's canonical measure μ .

Namely, for any edge e

$$\frac{\#(W(D_N) \cap e)}{N} \rightarrow \mu(e) \quad \text{as} \quad N \rightarrow \infty.$$

Idea:

(discrete current flow) $\xrightarrow{N \rightarrow \infty}$ (continuous current flow)

Tropical Weierstrass distribution: proof idea

Theorem (R)

For a sequence of generic divisor classes $[D_N]$ on Γ , the Weierstrass locus $W(D_N)$ distributes according to Zhang's canonical measure μ .

Namely, for any edge e

$$\frac{\#(W(D_N) \cap e)}{N} \rightarrow \mu(e) \quad \text{as} \quad N \rightarrow \infty.$$

Idea:

$$\begin{array}{ccc} \text{(discrete current flow)} & \xrightarrow{N \rightarrow \infty} & \text{(continuous current flow)} \\ & & \updownarrow \\ & & \text{canonical measure } \mu(e) \end{array}$$

Tropical Weierstrass distribution: proof idea

Theorem (R)

For a sequence of generic divisor classes $[D_N]$ on Γ , the Weierstrass locus $W(D_N)$ distributes according to Zhang's canonical measure μ .


Namely, for any edge e


$$\frac{\#(W(D_N) \cap e)}{N} \rightarrow \mu(e) \quad \text{as} \quad N \rightarrow \infty.$$


Idea:


$$\begin{array}{ccc} \text{(discrete current flow)} & \xrightarrow{N \rightarrow \infty} & \text{(continuous current flow)} \\ \updownarrow & & \updownarrow \\ \#(\text{Weierstrass points on } e) & & \text{canonical measure } \mu(e) \end{array}$$


References

 David Mumford (1977)
Curves and their Jacobians
The University of Michigan Press, Ann Arbor, MI.


 Amnon Neeman (1984)
The distribution of Weierstrass points on a compact Riemann surface
Ann. of Math. **120** 317–328.

 Shouwu Zhang (1993)
Admissable pairing on a curve
Invent. Math. **112** 171–193.

 Matt Baker and Xander Faber (2006)
Metrized graphs, Laplacian operators, and electrical networks
Amer. Math. Soc., Providence, RI.

 Omid Amini (2014)
Equidistribution of Weierstrass points on curves over non-Archimedean fields
preprint, [arXiv:1412.0926v1](https://arxiv.org/abs/1412.0926v1).

Weierstrass points on a tropical curve



Thank you!