

Clinical Trial Data

CANXIOL® EVIDENCE SUMMARY

A prospective, randomized, double-blind, multicenter comparative study to Evaluate the Efficacy, Safety and Pharmacokinetics of Cannabidiol oral solution versus matching Placebo for treatment of mild to moderate anxiety disorders

PHASE III · RCT · DOUBLE-BLIND · PARALLEL-GROUP

10

sites, 4 regions of India

178 randomized

89 CBD : 89 Placebo

Significant response, all clinical endpoints

Anxiety, clinician global impression, depressive symptoms, and sleep quality each improved by roughly half to two-thirds from baseline with cannabidiol.

Well-tolerated

41 of 178 participants reported at least one adverse event; none were serious, and none led to treatment discontinuation.

Endpoints

Primary: GAD-7, HAM-A


Secondary: CGI-I, CGI-S, PHQ-9, PSQI

Assessed: Baseline and Visit 9 (end of treatment)

DOSING REGIMEN

150 mg BID in week 1, titrated to 225 mg BID in week 2 and 300 mg BID in week 3, continued for 8 weeks (Visit 9), then tapered to 150 mg QD for 1 week. The dose is added to about 150 mL of water, mixed, and consumed 30 minutes after food.


SECONDARY ENDPOINTS – CLINICIAN-RATED SCALES



Both co-primary endpoints met

GAD-7 – 7.02 pts • HAM-A – 11.9 pts vs. placebo, both $p < 0.0001$

PRIMARY ENDPOINTS – ANXIETY SCORES IMPROVED SIGNIFICANTLY



SECONDARY ENDPOINTS – MOOD & SLEEP IMPROVED ALONGSIDE ANXIETY

