

# The Pisticci Farm Project

The Pisticci Farm Project is a project of the Upper Manhattan restaurant [Pisticci](#). The project has three elements:

- *engaging* the **restaurant** staff and patrons
- creating *zero waste* through **composting** within the city
- *producing* high-quality vegetables for Pisticci on the **multi-site farm**

This document outlines the technologies and processes involved in the composting and farming elements of the project.

## Compost

The discarded organic material from Pisticci restaurant are separated at source (ie. the kitchen, bar, etc.) from other ‘waste’. These materials are composted using an oxygenated hot composting method to ensure rapid decomposition, no unpleasant odours and the creation of a high quality fertiliser and biological inoculum for the soil at the **farm**.

## Aerated Static Pile (ASP) Composting

Pisticci is using the aerated static pile (ASP) method of composting as developed by Peter Moon of [O2 Compost](#).

The Pisticci ASP system using includes three enclosed boxes which are bottom-fed pressurised air through *100 mm (4 in)* from a blower mounted above the bins.

## Organic Material Collection Process

- Organic materials are collected in white plastic bags by the restaurant staff and deposited in wheelie-bins next to the compost bins.

## Composting Processes   Establishing a new Pile

1. Ensure that all pipes in the manifold in the bottom of the bin are present and properly fitted together.
2. Add woodchips to just cover the pipes and thoroughly moisten the woodchips.
3. Open the valve above the bin to ensure that air will flow into the pile while composting proceeds.

Organic materials may now be added.

### **Ongoing Addition of Organic Materials**

1. Use the composting fork to mix through any previously added materials and add water from the hose to bring the moisture levels up to approximately 50% moisture.
2. Empty two or three white bags of organic materials into the bin currently in use. Attempt to mix bags which are mostly vegetable waste (high nitrogen materials) and bags which are mostly napkins / dry 'high carbon' materials.
3. Mix these materials together in the bin using the composting fork.
4. Use the bucket to carry several buckets of woodchips from the woodchip storage bin to the compost bin to cover the material from the bags.
5. Mix the woodchips into the other materials using the composting fork.
6. Allow the materials to sit in the bin until the next addition to absorb moisture before adding any additional water (see point 1 above).

### **When a Bin is Full**

1. When a bin is full, make sure the moisture levels are appropriate (ie. 50% moisture), add a layer of woodchip over the top layer and cover with geotextile fabric / weed matting.
2. ensure the valve for the air is in the fully open position.
3. Leave to mature for as long as possible before the compost is taken up to the farm
4. Compost will be taken to the farm when two bins are full and one bin is one quarter to half-way full.

### **Transporting Compost to the Farm**

When two bins are full and one bin is one quarter to half-way full, the most mature pile should be transported to the farm.

1. Position the truck somewhere near to the front of the restaurant which will allow for wheelbarrow access via the gate at the side of the building.
2. Lift the front panel of the bin to be removed so the the bolts which secure it are lifted from their holes and set the front panel aside.
3. Using a wheelbarrow, transport the compost to the truck until the bin is empty.
4. Clean all areas and close the bin by returning the front panel to its place.

### **System Specifications**

- **The Bins** are 5' long, 4' high and 3' wide to fit within the small alleyway behind the restaurant.
- **The Blower**
- **The Timer** is set to be on for 2 minutes every 30 minutes to ensure the piles remain oxygenated.

## Farm

The Pisticci Farm Project is a multi-site micro farm. At the time of writing the farm consists of two sites:

- 43 Old Post Rd South, Croton-on-Hudson, NY (OPRS); and
- 716 Kitchawan Rd, Ossining, NY (KITCH).

The **Old Post Rd South** site was developed and farmed in 2015. The **Kitchawan** site was secured at the end of 2015 and will be developed and farmed in addition to the OPRS site in 2016.

## Old Post Rd South

The Old Post Rd South site is the flagship site of the Pisticci Farm Project. It consists of indoor and outdoor growing space. There are two large old glass greenhouses which have been restored and developed for all season production of vegetables. The 2015 growing season was used for diverse crop production with a focus on fast growing, high value crops, variety trials and winter greens production. In 2016, this site will be used primarily for all season greens production and greenhouse production of tomatoes, peppers, eggplant and cucumbers.

**The Outdoor Growing Area** The outdoor growing area consists of 27 beds of varying sizes in three blocks (Lower, Middle, Top). The beds are all 30 inches wide with 18 inch paths between them and vary in size from 10 to 40 feet with a total of 620 linear feet of bed space (1550 ft<sup>2</sup>).

**The Greenhouses** In 2015, the indoor space consisted of 12 beds across two greenhouses. The beds ranged in size from 29 to 60 feet with a total of 517 linear feet of bed space (1292 ft<sup>2</sup>). In early 2016, these indoor beds were delineated with 6" x 2" boards to create a total of 14 beds 29 inches wide with 11 inch intermediate paths. This new configuration has a total of 550 bed feet (1630 ft<sup>2</sup>).

**Hydronic Heating** is used in the greenhouses to allow for all season production. The original soil is separated from the imported soil and compost by 2 inches of foam insulation. PEX tubing runs over the top of the insulation at 12 inch

spacing. The PEX tubing is covered with between 12 and 18 inches of topsoil mixed with compost. This soil is kept at a minimum of 50°F (10°C) to allow for ongoing plant growth during the cooler month.

## Kitchawan

Kitchawan Farm is a diversified working farm with a focus on rough housing of horses. The Pisticci Farm project leases a small area for vegetable production. In Spring of 2016, this area will be developed into sets of 30 inch wide beds. This site will be used for crops n

## Crops

Type	Crop	Varieties	DTM	Location
green	<i>arugula</i>	astro	21-30	OPR
green	<i>kale</i>	toscana	35-40	KITCH
green	<i>lettuce</i>	salanova	55	KITCH
green	<i>mesclun mix</i>	5 star; all star	28-30	OPR
green	<i>dandelion</i>	clio, garnet stem	35-48	OPR
green	<i>escarole</i>	eros	45	OPR
green	<i>beet greens</i>	bull's blood; early wonder; red devil	35	KITCH
cucurbit	<i>summer squash</i>	zephyr; goldmine; safari; slik pik	50-54	KITCH
cucurbit	<i>pattypan squash</i>	sunburst; g-start	50-52	KITCH
cucurbit	<i>cucumber</i>	katrina	48	OPR
solanum	<i>pepper</i>	carmen; sprinter; sympathy; moonset	60	GH; KITCH
solanum	<i>tomato</i>	taxi; rebelski; manero; beorange	70-75	GH
solanum	<i>tomato (cherry)</i>	santorango; favorita; yellow pear	58-70	GH
solanum	<i>eggplant</i>	nadia; orient charm; jaylo	62-67	GH; KITCH
herb	<i>basil</i>	nufar; neopolitano	77	GH; KITCH
root	<i>beet</i>	baby beet; boulder; red ace	40-50	KITCH
root	<i>radish</i>	d'avignon	21	KITCH

## Priority Crops for 2016

## Crop Descriptions

The following includes cultivation notes on each of the above crops if they were successfully grown in 2015. If crops were not grown in 2015 or the cultivation description is well detailed elsewhere then references are given.

### arugula

**Propagation:** Direct Sown

**Seeder:** Glasser (small bore)

**Rows:** 7-9 (more rows makes plants 'stemmy')

**Harvest:** Use Serrated Greens Knife (blue handle)

**Store:** Coolroom

### basil

**Propagation:** 72 cell tray

**Rows:** 3

**Spacing:** 12"

**Harvest:** Cut stems down to node with secateurs

**Store:** Coolroom

**beets** Beets were not grown as a root crop in 2015. Refer to *The Market Gardener* or other reference for cultural details.

### beet greens

**Propagation:** Direct Sown

**Seeder:** Earthway (chard plates)

**Rows:** 7

**Harvest:** Use Serrated Greens Knife (blue handle)

**Store:** Coolroom

**cucumber** Cucumbers were not grown in 2015. Refer to *The Market Gardener* or other reference for cultural details.

**dandelion**

**Propagation:** 72 cell tray  
**Rows:** 5  
**Spacing:** 6"  
**Harvest:** Use Serrated Greens Knife (blue handle)  
**Store:** Coolroom

**eggplant**

**Propagation:** 72 cell tray transplanted to 4" pot  
**Rows:** 1  
**Spacing:** 18"  
**Harvest:** Cut stems above fruit with **secateurs**  
**Store:** Basement

**escarole**

**Propagation:** 72 cell tray  
**Rows:** 4  
**Spacing:** 8"  
**Harvest:** Use Serrated Greens Knife (blue handle)  
**Store:** Coolroom

**kale** *baby leaf*

**Propagation:** Direct Sown  
**Seeder:** Glasser (small bore)  
**Rows:** 7  
**Harvest:** Use Serrated Greens Knife (blue handle)  
**Store:** Coolroom

*large leaf*

**Propagation:** 72 cell tray  
**Rows:** 4  
**Spacing:** 8"  
**Harvest:** twist off mature leaves  
**Store:** Coolroom

**lettuce (salanova)**

**Propagation:** 72 cell tray

**Rows:** 4

**Spacing:** 8"

**Harvest:** Cut off outside leaves with lettuce field knife  
(yellow handle)

**Store:** Coolroom

**mesclun mix**

**Propagation:** Direct Sown

**Seeder:** Glasser (large bore)

**Rows:** 9

**Harvest:** Use Serrated Greens Knife (blue handle)

**Store:** Coolroom

**pattypan squash**

**Propagation:** 72 cell tray transplanted to 4" pot

**Rows:** 1

**Spacing:** 18"

**Harvest:** twist off or cut off fruits at base of stem

**Store:** Basement

**pepper**

**Propagation:** 72 cell tray transplanted to 4" pot

**Rows:** 1

**Spacing:** 18"

**Harvest:** twist off or cut off fruits at base of stem

**Store:** Basement

**radish** Radishes were not grown in 2015. Refer to The Market Gardener or other reference for cultural details.

## summer squash

**Propagation:** 72 cell tray transplanted to 4" pot

**Rows:** 1

**Spacing:** 18"

**Harvest:** twist off or cut off fruits at base of stem

**Store:** Basement

## tomato

**Propagation:** 72 cell tray transplanted to 4" pot

**Rows:** 1

**Spacing:** 18"

**Harvest:** twist off or cut off fruits at base of stem

**Store:** Basement

Refer to **The Market Gardener** or other reference for details of trellising system and offset training.

## Nursery

Pisticci Full Circle Farm produces all of its own seedlings. All seed to date has been purchased from Johnny's Selected Seed. Potting medium is made at the farm.

**Potting Medium** Potting soil is usually made up in batches of one wheelbarrow with the major ingredients being measured using a 5 gallon bucket. The **potting soil** consists of: \* 2 buckets sifted compost \* 2 buckets sifted coir \* 1 bucket sifted bed soil \* 3 tbsp fish hydrolysate \* 1 tsp solu-kelp \* 3 tbsp EM \* 2 buckets water

### *Method*

1. mix fish hydrolysate, solu-kelp and EM in water
2. add coir to water and let stand until expanded; add extra water if required
3. sift all other ingredients and mix together thoroughly
4. store in galvanised bin, preferably for a week or more before use.



## Fertility

Soil is at the heart of any vegetable growing system with integrity. The Pisticci Farm project utilises a number of strategies for developing and maintaining a healthy soil for the production of healthy nutritious plants. These include:

- composting
- soil testing and amending for mineral balance
- biological inocula / biofertiliser
- minimal tillage
- fertigation

**Soil Testing and Amending for Mineral Balance** Soil testing has been carried out to determine how to amend soil to achieve a balance of minerals for ideal plant health. The principal tool used to determine the ‘ideal’ is the set of worksheets created by Steve Solomon and Erica Reinheimer (available at <http://www.newsociety.com/var/storage/blurbs/IntelligentGardener-Worksheets.pdf>). This process is well described in “The Intelligent Gardener” by Steve Solomon.

## Recommended Soil Treatments for 2016    TODO TABLE

**Biological Inocula** The principal form of biological activation / inoculation of soil and leaf surfaces is through the use of EM; effective microorganisms.

**Activated EM** is created through the following process:

### *Ingredients and Equipment*

1. EM • 1® — 1 gallon
2. Unsulfured Blackstrap Molasses — 1 gallon
3. water (preferably warm / ~ 110°F (43°C)).
4. pH test papers with range to 3.5
5. airtight container and airlock

### *Method*

1. Mix ingredients in the container.
2. Check the initial pH with pH paper.
3. Put on lid and airlock
4. Ferment at room temperature for 3-5 days.

5. Some time between days 3 and 5, remove the lid and check the pH of the liquid using pH paper. If the pH is 3.8 or below, allow the fermentation to complete for an additional 5-7 days. Toward the end of the fermentation, check the smell of the product. It should have some alcohol smell, some white flakes on it and look and smell similar to the original EM • 1®. If all these are true, it is ready to use.

**Activated EM** is used in a range of applications at Pisticci Full Circle Farm:

- **Foliar Spray** — 3 tbsp (1.5 oz) of *activated EM* to 1.5 gallons of water — apply to leaf surfaces with sprayer
  1. other soluble fertilisers (solukelp, fish hydrolysate, blood meal) may be added
- **Potting Media** – add 3 tbsp to the water used to hydrate the coir block or add to existing potting mix
- **Soil Drench** — 3 tbsp (1.5 oz) of *activated EM* to 1.5 gallons of water — apply to soil with sprayer or through venturi.

**Fertigation** A Mazzei venturi injection system has been installed in the water line that preceeds the solenoid valves which control each of the irrigation stations. This allows soluble nutrient and biological innocula to be applied through the irrigation water given to crops.

The ideal use of this system is still being worked out. The current regime is as follows:

*Application Rate*

- EM — ( 40 gallons / acre / year) **0.45 oz / 100 ft<sup>2</sup> / 14 days**
- Solukelp — (312 oz / acre / year) **0.03 oz / 100 ft<sup>2</sup> / 14 days**
- Neptune's Harvest Fish Fertilizer — (12 gallons / acre / year) **0.14 oz / 100 ft<sup>2</sup> / 14 days**

*Western Greenhouse* (400 sq ft bed area)

- EM — 6 tbsp / 14 days
- Solu-Kelp — 1 tsp / 14 days
- Neptune's Harvest Fish Fertilizer — 2 tbsp / 14 days

*Eastern Greenhouse* (900 sq ft bed area)

- EM — 9 tbsp / 14 days
- Solu-Kelp — 1.5 tsp / 14 days
- Neptune's Harvest Fish Fertilizer — 2.5 tbsp / 14 days

## Suppliers

### Aloha Medicinals

**url:** <http://www.alohamedicinals.com/>

**details:** inoculated grain, mushroom growing supplies

### Arbico Organics

**url:** <http://www.arbico-organics.com/>

**details:** EM • 1®

### Central Irrigation (Elmsford)

**url:** <http://www.centralirrigationsupply.com/>

**details:** general irrigation supplies

### Compostwerks

**url:** <http://www.compostwerks.com/>

**details:** soil ammendments, biofertiliser ingredients

**contacts:** *Gregg Twehues* ([gregg@compostwerks.com](mailto:gregg@compostwerks.com)); *Peter Schmidt* ([peter@compostwerks.com](mailto:peter@compostwerks.com))

### Farmtek

**url:** <http://www.farmtek.com/>

**details:** general equipment

**contact:** *Virginia Daly* ([vdaly@farmtek.com](mailto:vdaly@farmtek.com))

### Growers Supply

**url:** <http://www.growerssupply.com/>

**details:** general equipment

### Johnny's Selected Seeds

**url:** <http://www.johnnyseeds.com/>

**details:** seed, tools

## Logan Labs

**url:** <http://www.loganlabs.com/>

**details:** soil testing

## Nolts Produce

**url:** <http://www.noltsproducesupplies.net/>

**details:** row cover, irrigation supplies, tools

## O2 Compost

**url:** <http://www.o2compost.com/>

**details:** composting support

**contact:** *Peter Moon* ([peter@o2compost.com](mailto:peter@o2compost.com))

## Teraganix

**url:** <http://www.teraganix.com/>

**details:** EM • 1®, pH test papers

## The Green Growler (Croton)

**url:** <http://thegreengrowler.com/>

**details:** airlocks

## Glossary

**50% moisture** - Compost or composting materials with approximately 50% moisture will make your hand moist when squeezed or produce a drop or two of water.

**high carbon materials** - organic materials such as paper, napkins, cardboard, woodchips, straw etc. which have relatively much more elemental carbon (C) than elemental nitrogen (N).

**high nitrogen materials** - organic materials such as manure, vegetable scraps, grass clippings etc. which have high levels of elemental nitrogen (N) relative to elemental carbon (C).