

CHAPTER

5

Mathematical Induction
and Binomial Theorem

Section-A

JEE Advanced/ IIT-JEE

A Fill in the Blanks

- The larger of $99^{50} + 100^{50}$ and 101^{50} is
(1982 - 2 Marks)
- The sum of the coefficients of the polynomial $(1 + x - 3x^2)^{2163}$ is
(1982 - 2 Marks)
- If $(1 + ax)^n = 1 + 8x + 24x^2 + \dots$ then $a = \dots$ and $n = \dots$
(1983 - 2 Marks)
- Let n be positive integer. If the coefficients of 2nd, 3rd, and 4th terms in the expansion of $(1 + x)^n$ are in A.P., then the value of n is
(1994 - 2 Marks)
- The sum of the rational terms in the expansion of $(\sqrt{2} + 3^{1/5})^{10}$ is
(1997 - 2 Marks)

C MCQs with One Correct Answer

- Given positive integers $r > 1, n > 2$ and that the coefficient of $(3r)$ th and $(r + 2)$ th terms in the binomial expansion of $(1 + x)^{2n}$ are equal. Then
(1983 - 1 Mark)
(a) $n = 2r$ (c) $n = 2r + 1$
(d) none of these
- The coefficient of x^4 in $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$ is (1983 - 1 Mark)
(a) $\frac{405}{256}$ (b) $\frac{504}{259}$
(c) $\frac{450}{263}$ (d) none of these
- The expression $\left(x + (x^3 - 1)^{\frac{1}{2}}\right)^5 + \left(x - (x^3 - 1)^{\frac{1}{2}}\right)^5$ is a polynomial of degree (1992 - 2 Marks)
(a) 5 (b) 6 (c) 7 (d) 8
- If in the expansion of $(1 + x)^m (1 - x)^n$, the coefficients of x and x^2 are 3 and -6 respectively, then m is (1999 - 2 Marks)
(a) 6 (b) 9 (c) 12 (d) 24
- For $2 \leq r \leq n$, $\binom{n}{r} + 2\binom{n}{r-1} + \binom{n}{r-2} =$ (2000S)
(a) $\binom{n+1}{r-1}$ (b) $2\binom{n+1}{r+1}$ (c) $2\binom{n+2}{r}$ (d) $\binom{n+2}{r}$

- In the binomial expansion of $(a - b)^n, n \geq 5$, the sum of the 5th and 6th terms is zero. Then a/b equals (2001S)
(a) $(n-5)/6$ (b) $(n-4)/5$
(c) $5/(n-4)$ (d) $6/(n-5)$

- The sum $\sum_{i=0}^m \binom{10}{i} \binom{20}{m-i}$, (where $\binom{p}{q} = 0$ if $p < q$) is maximum when m is (2002S)
(a) 5 (b) 10 (c) 15 (d) 20
- Coefficient of t^{24} in $(1+t^2)^{12} (1+t^{12}) (1+t^{24})$ is (2003S)
(a) ${}^{12}C_6 + 3$ (b) ${}^{12}C_6 + 1$ (c) ${}^{12}C_6$ (d) ${}^{12}C_6 + 2$
- If ${}^{n-1}C_r = (k^2 - 3) {}^nC_{r+1}$, then $k \in$ (2004S)
(a) $(-\infty, -2]$ (b) $[2, \infty)$ (c) $[-\sqrt{3}, \sqrt{3}]$ (d) $(\sqrt{3}, 2]$
- The value of

$$\binom{30}{0}\binom{30}{10} - \binom{30}{1}\binom{30}{11} + \binom{30}{2}\binom{30}{12} - \dots + \binom{30}{20}\binom{30}{30}$$

is where

$$\binom{n}{r} = {}^nC_r \quad (2005S)$$

- (a) $\binom{30}{10}$ (b) $\binom{30}{15}$ (c) $\binom{60}{30}$ (d) $\binom{31}{10}$
- For $r = 0, 1, \dots, 10$, let A_r, B_r and C_r denote, respectively, the coefficient of x^r in the expansions of $(1+x)^{10}, (2010)$
 $(1+x)^{20}$ and $(1+x)^{30}$. Then $\sum_{r=1}^{10} A_r(B_{10}B_r - C_{10}A_r)$ is equal to
(a) $B_{10} - C_{10}$ (b) $A_{10}(B_{10}^2 - C_{10}A_{10})$
(c) 0 (d) $C_{10} - B_{10}$
- Coefficient of x^{11} in the expansion of $(1+x^2)^4(1+x^3)^7(1+x^4)^{12}$ is (JEE Adv. 2014)
(a) 1051 (b) 1106 (c) 1113 (d) 1120

D MCQs with One or More than One Correct

- If C_r stands for nC_r , then the sum of the series
$$\frac{2\left(\frac{n}{2}\right)! \left(\frac{n}{2}\right)!}{n!} [C_0^2 - 2C_1^2 + 3C_2^2 - \dots + (-1)^n (n+1)C_n^2],$$

where n is an even positive integer, is equal to (1986 - 2 Marks)

(a) 0 (b) $(-1)^{n/2}(n+1)$

(c) $(-1)^{n/2}(n+2)$ (d) $(-1)^n n$

(e) none of these.

2. If $a_n = \sum_{r=0}^n \frac{1}{n C_r}$, then $\sum_{r=0}^n \frac{r}{n C_r}$ equals (1998 - 2 Marks)

(a) $(n-1)a_n$

(b) na_n

(c) $\frac{1}{2}na_n$

(d) None of the above

E Subjective Problems

1. Given that (1979)
 $C_1 + 2C_2x + 3C_3x^2 + \dots + 2nC_{2n}x^{2n-1} = 2n(1+x)^{2n-1}$

where $C_r = \frac{(2n)!}{r!(2n-r)!}$ $r=0, 1, 2, \dots, 2n$

Prove that

$C_1^2 - 2C_2^2 + 3C_3^2 - \dots - 2nC_{2n}^2 = (-1)^n C_n$

2. Prove that $7^{2n} + (2^{3n-3})(3^{n-1})$ is divisible by 25 for any natural number n . (1982 - 5 Marks)

3. If $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$ then show that the sum of the products of the C_i 's taken two at a time,

represented by $\sum_{0 \leq i < j \leq n} C_i C_j$ is equal to $2^{2n-1} - \frac{(2n)!}{2(n!)^2}$

(1983 - 3 Marks)

4. Use mathematical Induction to prove : If n is any odd positive integer, then $n(n^2 - 1)$ is divisible by 24. (1983 - 2 Marks)

5. If p be a natural number then prove that $p^{n+1} + (p+1)^{2n-1}$ is divisible by $p^2 + p + 1$ for every positive integer n . (1984 - 4 Marks)

6. Given $s_n = 1 + q + q^2 + \dots + q^n$;

$S_n = 1 + \frac{q+1}{2} + \left(\frac{q+1}{2}\right)^2 + \dots + \left(\frac{q+1}{2}\right)^n, q \neq 1$ Prove that

$n^{+1}C_1 + n^{+1}C_2s_1 + n^{+1}C_3s_2 + \dots + n^{+1}C_ns_n = 2^n S_n$ (1984 - 4 Marks)

7. Use method of mathematical induction $2 \cdot 7^n + 3 \cdot 5^n - 5$ is divisible by 24 for all $n > 0$ (1985 - 5 Marks)

8. Prove by mathematical induction that - (1987 - 3 Marks)

$\frac{(2n)!}{2^{2n}(n!)^2} \leq \frac{1}{(3n+1)^{1/2}}$ for all positive Integers n .

9. Let $R = (5\sqrt{5} + 11)^{2n+1}$ and $f = R - [R]$, where $[]$ denotes the greatest integer function. Prove that $Rf = 4^{2n+4}$. (1988 - 5 Marks)

10. Using mathematical induction, prove that (1989 - 3 Marks)

${}^m C_0 {}^n C_k + {}^m C_1 {}^n C_{k-1} + \dots + {}^m C_k {}^n C_0 = ({}^{m+n} C_k)$,

where m, n, k are positive integers, and ${}^p C_q = 0$ for $p < q$.

11. Prove that (1989 - 5 Marks)

$C_0 - 2^2 C_1 + 3^2 C_2 - \dots + (-1)^n (n+1)^2 C_n = 0,$

 $n > 2$, where $C_r = {}^n C_r$.

12. Prove that $\frac{n^7}{7} + \frac{n^5}{5} + \frac{2n^3}{3} - \frac{n}{105}$ is an integer for every positive integer n . (1990 - 2 Marks)

13. Using induction or otherwise, prove that for any non-negative integers m, n, r and k , (1991 - 4 Marks)

$$\sum_{m=0}^k (n-m) \frac{(r+m)!}{m!} = \frac{(r+k+1)!}{k!} \left[\frac{n}{r+1} - \frac{k}{r+2} \right]$$

14. If $\sum_{r=0}^{2n} a_r (x-2)^r = \sum_{r=0}^{2n} b_r (x-3)^r$ and $a_k = 1$ for all

$k \geq n$, then show that $b_n = {}^{2n+1} C_{n+1}$ (1992 - 6 Marks)

15. Let $p \geq 3$ be an integer and α, β be the roots of $x^2 - (p+1)x + 1 = 0$ using mathematical induction show that $\alpha^n + \beta^n$.

(i) is an integer and (ii) is not divisible by p (1992 - 6 Marks)

16. Using mathematical induction, prove that

$\tan^{-1}(1/3) + \tan^{-1}(1/7) + \dots + \tan^{-1}\{1/(n^2 + n + 1)\}$
 $= \tan^{-1}\{n/(n+2)\}$ (1993 - 5 Marks)

17. Prove that $\sum_{r=1}^k (-3)^{r-1} {}^{3n} C_{2r-1} = 0$, where $k = (3n)/2$ and n is an even positive integer. (1993 - 5 Marks)

18. If x is not an integral multiple of 2π use mathematical induction to prove that : (1994 - 4 Marks)

$\cos x + \cos 2x + \dots + \cos nx = \cos \frac{n+1}{2} x \sin \frac{nx}{2} \operatorname{cosec} \frac{x}{2}$

19. Let n be a positive integer and (1994 - 5 Marks)
 $(1+x+x^2)^n = a_0 + a_1x + \dots + a_{2n}x^{2n}$
Show that $a_0^2 - a_1^2 + a_2^2 - \dots + a_{2n}^2 = a_n$

20. Using mathematical induction prove that for every integer $n \geq 1$, $(3^{2n}-1)$ is divisible by 2^{n+2} but not by 2^{n+3} . (1996 - 3 Marks)

21. Let $0 < A_i < \pi$ for $i = 1, 2, \dots, n$. Use mathematical induction to prove that

$\sin A_1 + \sin A_2 + \dots + \sin A_n \leq n \sin \left(\frac{A_1 + A_2 + \dots + A_n}{n} \right)$

where ≥ 1 is a natural number.

{You may use the fact that

$p \sin x + (1-p) \sin y \leq \sin [px + (1-p)y],$
where $0 \leq p \leq 1$ and $0 \leq x, y \leq \pi$.} (1997 - 5 Marks)

22. Let p be a prime and m a positive integer. By mathematical induction on m , or otherwise, prove that whenever r is an integer such that p does not divide r , p divides ${}^m p C_r$. (1998 - 8 Marks)

[Hint: You may use the fact that $(1+x)^{(m+1)p} = (1+x)^p (1+x)^{mp}$]

Mathematical Induction and Binomial Theorem

23. Let n be any positive integer. Prove that
(1999 - 10 Marks)

$$\sum_{k=0}^m \frac{\binom{2n-k}{k}}{\binom{2n-k}{n}} \cdot \frac{(2n-4k+1)}{(2n-2k+1)} 2^{n-2k} = \frac{\binom{n}{m}}{\binom{2n-2m}{n-m}} 2^{n-2m}$$

for each non-negative integer $m \leq n$. (Here $\binom{p}{q} = {}^pC_q$).

24. For any positive integer m, n (with $n \geq m$), let $\binom{n}{m} = {}^nC_m$.

$$\text{Prove that } \binom{n}{m} + \binom{n-1}{m} + \binom{n-2}{m} + \dots + \binom{m}{m} = \binom{n+1}{m+1}.$$

Hence or otherwise, prove that

$$\binom{n}{m} + 2\binom{n-1}{m} + 3\binom{n-2}{m} + \dots + (n-m+1)\binom{m}{m} = \binom{n+2}{m+2}.$$

(2000 - 6 Marks)

25. For every positive integer n , prove that

$$\sqrt{4n+1} < \sqrt{n} + \sqrt{n+1} < \sqrt{4n+2}. \text{ Hence or otherwise,}$$

prove that $[\sqrt{n} + \sqrt{n+1}] = [\sqrt{4n+1}]$, where $[x]$ denotes the greatest integer not exceeding x . (2000 - 6 Marks)

26. Let a, b, c be positive real numbers such that $b^2 - 4ac > 0$ and let $\alpha_1 = c$. Prove by induction that

$$\alpha_{n+1} = \frac{a\alpha_n^2}{(b^2 - 2a(\alpha_1 + \alpha_2 + \dots + \alpha_n))} \text{ is well-defined and}$$

$$\alpha_{n+1} < \frac{\alpha_n}{2} \text{ for all } n = 1, 2, \dots \text{ (Here, 'well-defined' means}$$

that the denominator in the expression for α_{n+1} is not zero.)
(2001 - 5 Marks)

27. Use mathematical induction to show that
(25)ⁿ⁺¹ - 24n + 5735 is divisible by (24)² for all $n = 1, 2, \dots$
(2002 - 5 Marks)
28. Prove that
(2003 - 2 Marks)

$$2^k \binom{n}{0} \binom{n}{k} - 2^{k-1} \binom{n}{2} \binom{n}{k-1} \binom{n-1}{1} \binom{n-1}{k-1} + 2^{k-2} \binom{n-2}{k-2} - \dots - (-1)^k \binom{n}{k} \binom{n-k}{0} = \binom{n}{k}.$$

29. A coin has probability p of showing head when tossed. It is tossed n times. Let p_n denote the probability that no two (or more) consecutive heads occur. Prove that $p_1 = 1$, $p_2 = 1 - p^2$ and $p_n = (1 - p) \cdot p_{n-1} + p(1 - p)p_{n-2}$ for all $n \geq 3$.

Prove by induction on n , that $p_n = A\alpha^n + B\beta^n$ for all $n \geq 1$, where α and β are the roots of quadratic equation

$$x^2 - (1-p)x - p(1-p) = 0 \text{ and } A = \frac{p^2 + \beta - 1}{\alpha\beta - \alpha^2}, B = \frac{p^2 + \alpha - 1}{\alpha\beta - \beta^2}.$$

(2000 - 5 Marks)

I

Integer Value Correct Type

1. The coefficients of three consecutive terms of $(1+x)^{n+5}$ are in the ratio 5 : 10 : 14. Then $n =$ (JEE Adv. 2013)
2. Let m be the smallest positive integer such that the coefficient of x^2 in the expansion of $(1+x)^2 + (1+x)^3 + \dots + (1+x)^{49} + (1+mx)^{50}$ is $(3n+1) {}^{51}C_3$ for some positive integer n . Then the value of n is (JEE Adv. 2016)

Section-B

JEE Main / AIEEE

1. The coefficients of x^p and x^q in the expansion of $(1+x)^{p+q}$ are [2002]
(a) equal
(b) equal with opposite signs
(c) reciprocals of each other
(d) none of these
2. If the sum of the coefficients in the expansion of $(a+b)^n$ is 4096, then the greatest coefficient in the expansion is [2002]
(a) 1594 (b) 792 (c) 924 (d) 2924
3. The positive integer just greater than $(1+0.0001)^{10000}$ is [2002]
(a) 4 (b) 5 (c) 2 (d) 3
4. r and n are positive integers $r > 1$, $n > 2$ and coefficient of $(r+2)^{\text{th}}$ term and $3r^{\text{th}}$ term in the expansion of $(1+x)^{2n}$ are equal, then n equals [2002]
(a) $3r$ (b) $3r+1$ (c) $2r$ (d) $2r+1$
5. If $a_n = \sqrt{7 + \sqrt{7 + \sqrt{7 + \dots}}}$ having n radical signs then by methods of mathematical induction which is true [2002]
(a) $a_n > 7 \forall n \geq 1$ (b) $a_n < 7 \forall n \geq 1$
(c) $a_n < 4 \forall n \geq 1$ (d) $a_n < 3 \forall n \geq 1$
6. If x is positive, the first negative term in the expansion of $(1+x)^{27/5}$ is [2003]
(a) 6th term (b) 7th term (c) 5th term (d) 8th term.
7. The number of integral terms in the expansion of $(\sqrt{3} + \sqrt[5]{5})^{256}$ is [2003]
(a) 35 (b) 32 (c) 33 (d) 34
8. Let $S(K) = 1 + 3 + 5 + \dots + (2K-1) = 3 + K^2$. Then which of the following is true [2004]
(a) Principle of mathematical induction can be used to prove the formula
(b) $S(K) \Rightarrow S(K+1)$
(c) $S(K) \not\Rightarrow S(K+1)$
(d) $S(1)$ is correct
9. The coefficient of the middle term in the binomial expansion in powers of x of $(1+\alpha x)^4$ and of $(1-\alpha x)^6$ is the same if α equals [2004]
(a) $\frac{3}{5}$ (b) $\frac{10}{3}$ (c) $-\frac{3}{10}$ (d) $-\frac{5}{3}$

10. The coefficient of x^n in expansion of $(1+x)(1-x)^n$ is
 (a) $(-1)^{n-1}n$ (b) $(-1)^n(1-n)$ [2004]
 (c) $(-1)^{n-1}(n-1)^2$ (d) $(n-1)$
11. The value of ${}^{50}C_4 + \sum_{r=1}^6 {}^{56-r}C_3$ is [2005]
 (a) ${}^{55}C_4$ (b) ${}^{55}C_3$ (c) ${}^{56}C_3$ (d) ${}^{56}C_4$
12. If $A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then which one of the following holds for all $n \geq 1$, by the principle of mathematical induction [2005]
 (a) $A^n = nA - (n-1)I$ (b) $A^n = 2^{n-1}A - (n-1)I$
 (c) $A^n = nA + (n-1)I$ (d) $A^n = 2^{n-1}A + (n-1)I$
13. If the coefficient of x^7 in $\left[ax^2 + \left(\frac{1}{bx}\right)\right]^{11}$ equals the coefficient of x^{-7} in $\left[ax - \left(\frac{1}{bx^2}\right)\right]^{11}$, then a and b satisfy the relation [2005]
 (a) $a-b=1$ (b) $a+b=1$
 (c) $\frac{a}{b}=1$ (d) $ab=1$
14. If x is so small that x^3 and higher powers of x may be neglected, then $\frac{(1+x)^{\frac{3}{2}} - \left(1 + \frac{1}{2}x\right)^3}{(1-x)^{\frac{1}{2}}}$ may be approximated as [2005]
 (a) $1 - \frac{3}{8}x^2$ (b) $3x + \frac{3}{8}x^2$
 (c) $-\frac{3}{8}x^2$ (d) $\frac{x}{2} - \frac{3}{8}x^2$
15. If the expansion in powers of x of the function $\frac{1}{(1-ax)(1-bx)}$ is $a_0 + a_1x + a_2x^2 + a_3x^3 + \dots$, then a_n is [2006]
 (a) $\frac{b^n - a^n}{b-a}$ (b) $\frac{a^n - b^n}{b-a}$
 (c) $\frac{a^{n+1} - b^{n+1}}{b-a}$ (d) $\frac{b^{n+1} - a^{n+1}}{b-a}$
16. For natural numbers m, n if $(1-y)^m(1+y)^n = 1 + a_1y + a_2y^2 + \dots$ and $a_1 = a_2 = 10$, then (m, n) is [2006]
 (a) (20, 45) (b) (35, 20)
 (c) (45, 35) (d) (35, 45)
17. In the binomial expansion of $(a-b)^n$, $n \geq 5$, the sum of 5th and 6th terms is zero, then a/b equals [2007]
 (a) $\frac{n-5}{6}$ (b) $\frac{n-4}{5}$ (c) $\frac{5}{n-4}$ (d) $\frac{6}{n-5}$
18. The sum of the series ${}^{20}C_0 - {}^{20}C_1 + {}^{20}C_2 - {}^{20}C_3 + \dots - \dots + {}^{20}C_{10}$ is [2007]
 (a) 0 (b) ${}^{20}C_{10}$ (c) $-{}^{20}C_{10}$ (d) $\frac{1}{2} {}^{20}C_{10}$
19. Statement -1 : $\sum_{r=0}^n (r+1) {}^nC_r = (n+2)2^{n-1}$. [2008]
 Statement-2 : $\sum_{r=0}^n (r+1) {}^nC_r x^r = (1+x)^n + nx(1+x)^{n-1}$.
 (a) Statement -1 is false, Statement-2 is true
 (b) Statement -1 is true, Statement-2 is true; Statement -2 is a correct explanation for Statement-1
 (c) Statement -1 is true, Statement-2 is true; Statement -2 is not a correct explanation for Statement-1
 (d) Statement -1 is true, Statement-2 is false
20. The remainder left out when $8^{2n} - (62)^{2n+1}$ is divided by 9 is : [2009]
 (a) 2 (b) 7 (c) 8 (d) 0
21. Let $S_1 = \sum_{j=1}^{10} j(j-1) {}^{10}C_j$, $S_2 = \sum_{j=1}^{10} j {}^{10}C_j$ and $S_3 = \sum_{j=1}^{10} j^2 {}^{10}C_j$.
 Statement-1 : $S_3 = 55 \times 2^9$.
 Statement-2 : $S_1 = 90 \times 2^8$ and $S_2 = 10 \times 2^8$. [2010]
 (a) Statement -1 is true, Statement-2 is true ; Statement -2 is not a correct explanation for Statement -1.
 (b) Statement -1 is true, Statement -2 is false.
 (c) Statement -1 is false, Statement -2 is true .
 (d) Statement -1 is true, Statement 2 is true ; Statement -2 is a correct explanation for Statement -1.
22. The coefficient of x^7 in the expansion of $(1-x-x^2+x^3)^6$ is [2011]
 (a) -132 (b) -144 (c) 132 (d) 144
23. If n is a positive integer, then $(\sqrt{3}+1)^{2n} - (\sqrt{3}-1)^{2n}$ is : [2012]
 (a) an irrational number
 (b) an odd positive integer
 (c) an even positive integer
 (d) a rational number other than positive integers
24. The term independent of x in expansion of $\left(\frac{x+1}{x^{2/3}-x^{1/3}+1} - \frac{x-1}{x-x^{1/2}}\right)^{10}$ is [JEE M 2013]
 (a) 4 (b) 120 (c) 210 (d) 310
25. If the coefficients of x^3 and x^4 in the expansion of $(1+ax+bx^2)(1-2x)^{18}$ in powers of x are both zero, then (a, b) is equal to: [JEE M 2014]
 (a) $\left(14, \frac{272}{3}\right)$ (b) $\left(16, \frac{272}{3}\right)$ (c) $\left(16, \frac{251}{3}\right)$ (d) $\left(14, \frac{251}{3}\right)$
26. The sum of coefficients of integral power of x in the binomial expansion $(1-2\sqrt{x})^{50}$ is : [JEE M 2015]
 (a) $\frac{1}{2}(3^{50}-1)$ (b) $\frac{1}{2}(2^{50}+1)$
 (c) $\frac{1}{2}(3^{50}+1)$ (d) $\frac{1}{2}(3^{50})$
27. If the number of terms in the expansion of $\left(1 - \frac{2}{x} + \frac{4}{x^2}\right)^n$, $x \neq 0$, is 28, then the sum of the coefficients of all the terms in this expansion, is : [JEE M 2016]
 (a) 243 (b) 729 (c) 64 (d) 2187