

Assignment 16

Sri Harsha CH

Abstract—This document explains the conditions for two vector spaces to be isomorphic.

Download all python codes from

<https://github.com/harshachinta/EE5609-Matrix-Theory/tree/master/Assignments/Assignment16/code>

and latex-tikz codes from

<https://github.com/harshachinta/EE5609-Matrix-Theory/tree/master/Assignments/Assignment16>

1 PROBLEM

Let \mathbf{V} and \mathbf{W} be finite-dimensional vector spaces over the field \mathbf{F} . Prove that \mathbf{V} and \mathbf{W} are isomorphic if and only if $\dim \mathbf{V} = \dim \mathbf{W}$

2 EXPLANATION

If \mathbf{V} and \mathbf{W} are vector spaces over the field \mathbf{F} , any one to one linear transformation T of \mathbf{V} onto \mathbf{W} is called an isomorphism of \mathbf{V} onto \mathbf{W} .

Let $T : \mathbf{V} \rightarrow \mathbf{W}$ be an isomorphism and let $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis for \mathbf{V} . Since it is isomorphic we know that T is one to one and onto. We need to show that $\{T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_n)\}$ is a basis for \mathbf{W} so that $\dim \mathbf{V} = n = \dim \mathbf{W}$.

1. To see that if set of vectors in \mathbf{W} are linearly independent we use the property that T is one to one. Let us consider a linear combination.

$$\alpha_1 T(\mathbf{v}_1) + \alpha_2 T(\mathbf{v}_2) + \dots + \alpha_n T(\mathbf{v}_n) = \mathbf{0} \quad (2.0.1)$$

$$T(\alpha_1 \mathbf{v}_1) + T(\alpha_2 \mathbf{v}_2) + \dots + T(\alpha_n \mathbf{v}_n) = \mathbf{0} \quad (2.0.2)$$

$$\implies \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n = \mathbf{0} \quad (2.0.3)$$

$$\implies \alpha_1 = \alpha_2 = \dots = \alpha_n = 0 \quad (2.0.4)$$

From equation (2.0.1) and (2.0.4), the set of vectors $\{T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_n)\}$ are linearly independent.

2. To check if the set of vectors span \mathbf{W} . Since T is onto, for any $\mathbf{y} \in \mathbf{W}$ there exists an $\mathbf{x} \in \mathbf{V}$ such that $T(\mathbf{x}) = \mathbf{y}$.

$$\mathbf{x} = \alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n \quad (2.0.5)$$

$$T(\mathbf{x}) = T(\alpha_1 \mathbf{v}_1 + \alpha_2 \mathbf{v}_2 + \dots + \alpha_n \mathbf{v}_n) = \mathbf{y} \quad (2.0.6)$$

$$\alpha_1 T(\mathbf{v}_1) + \alpha_2 T(\mathbf{v}_2) + \dots + \alpha_n T(\mathbf{v}_n) = \mathbf{y} \quad (2.0.7)$$

From equation (2.0.7), any vector in \mathbf{W} can be represented as linear combination of $\{T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_n)\}$.

Therefore from equation (2.0.4) and (2.0.7), $\{T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_n)\}$ are linearly independent and span \mathbf{W} .

$$\implies \dim \mathbf{W} = n = \dim \mathbf{V}$$

3 SOLUTION

\mathbf{V} and \mathbf{W} are isomorphic if and only if $\dim \mathbf{V} = \dim \mathbf{W}$