

Comprehensive Study Notes on Causality Assessment in Pharmacovigilance: Principles, Standardized Algorithms, and Regulatory Applications

Foundations of Causality Assessment in Pharmacovigilance

The Conceptual Shift from Adverse Event to Adverse Drug Reaction

Pharmacovigilance is rooted in the continuous scientific evaluation of the causal link between pharmaceutical interventions and adverse clinical outcomes. In clinical safety operations, an Adverse Event (AE) is defined as any untoward medical occurrence in a patient or clinical investigation subject administered a pharmaceutical product, which does not necessarily possess a confirmed causal relationship with the treatment. An AE can present as an unfavorable or unintended sign, symptom, disease state, or abnormal laboratory metric that is temporally associated with the use of a medicinal compound, regardless of whether it is considered product-related.

Causality assessment, formally designated as relatedness assessment, constitutes the structured evaluation of the likelihood that a suspect medicinal agent is etiologically responsible for the observed adverse event. When scientific, medical, and pharmacological analysis determines that there is a reasonable possibility of a causal link, the event undergoes a regulatory transition from an AE to an Adverse Drug Reaction (ADR). This operational differentiation governs global pharmacovigilance obligations. While every ADR originates as an AE, only confirmed or suspected ADRs trigger targeted risk-benefit re-evaluations, revisions of the Reference Safety Information (RSI), updates to the Company Core Data Sheet (CCDS), and expedited notifications to international regulatory authorities.

Safety Phenomenon	Regulatory Definition	Evidentiary Threshold	Lifecycle Regulatory Impact
Adverse Event (AE)	Any untoward medical occurrence in a patient administered a medicinal product, lacking mandatory causal relatedness.	Mere coincidence in time; requires temporal overlap without confirmed mechanism.	Logged within internal safety databases; tracked for potential aggregate statistical disproportionality.
Adverse Drug Reaction (ADR)	A noxious and unintended response to a medicinal product where a causal	Plausible pharmacology, consistent latency, and exclusion of obvious	Incorporated into safety aggregate reports; drives updates to Prescribing Information

Safety Phenomenon	Regulatory Definition	Evidentiary Threshold	Lifecycle Regulatory Impact
	connection is a reasonable possibility.	alternative causes.	and product warnings.

Operational Purpose in Drug Safety Lifecycle Management

Causality assessment operates as a critical gatekeeper within the pharmaceutical safety lifecycle, functioning across individual case safety report (ICSR) intake, ongoing clinical trials, and post-marketing surveillance. At the individual case processing level, assigning relatedness dictates whether a safety narrative fulfills the criteria for expedited 7-day or 15-day regulatory reporting. On an aggregate population level, systematic causality determination reduces diagnostic noise, allowing drug safety systems to identify genuine biological risks. Rigorous relatedness assessment provides the core qualitative foundation for signal detection algorithms. Automated disproportionality metrics—such as the Proportional Reporting Ratio (PRR) and Information Component (IC)—rely on accurate primary case categorization to distinguish true pharmacological toxicities from background clinical noise. By systematically evaluating every safety case, marketing authorization holders and health agencies can quantify evolving benefit-risk balances, design targeted Risk Evaluation and Mitigation Strategies (REMS), implement mandatory risk minimization measures, and refine clinical dosology to safeguard public health.

Epidemiological Roots: The Bradford Hill Causation Framework

Algorithmic and qualitative causality frameworks in modern pharmacovigilance derive their foundational logic from the epidemiological criteria established by Sir Austin Bradford Hill in 1965. Rather than relying on subjective clinical intuition, standardized causality adjudication translates Hill's population-level criteria into individual patient assessments. Temporality serves as an absolute prerequisite, requiring that drug exposure precedes the onset of the adverse medical occurrence. The biological gradient, or dose-response relationship, provides mechanistic confirmation when an adverse reaction intensifies following dose escalation or diminishes after dosage reduction. Biological plausibility evaluates whether the observed pathology aligns with known receptor affinities, enzymatic metabolism, or toxicological dynamics. Coherence and consistency assess whether the event mirrors findings from preclinical animal models, randomized clinical trials, or documented class-effect toxicities. Finally, experimental evidence—represented in individual pharmacovigilance by the clinical course following drug cessation (dechallenge) and controlled or accidental re-exposure (rechallenge)—provides direct verification of drug-induced pathogenesis.

Core Determinants and Clinical Variables of Relatedness

Temporal Sequence and Biokinetic Plausibility

The establishment of a plausible temporal sequence requires that the interval between drug administration and event manifestation aligns with the underlying pharmacokinetic and pathophysiological processes. Evaluators must assess drug absorption rates, peak plasma

concentrations ($C_{\max}$), elimination half-life ($t_{1/2}$), and the time required to achieve steady-state concentration (C_{ss}) alongside the biological latency of the specific pathology. Latency intervals vary significantly across clinical reaction types. Immediate Type I IgE-mediated hypersensitivity reactions, such as anaphylaxis or urticaria, typically manifest within minutes to a few hours following drug administration. Conversely, delayed-type cell-mediated reactions, including severe cutaneous adverse reactions like Stevens-Johnson Syndrome (SJS) and Toxic Epidermal Necrolysis (TEN), characteristically exhibit an onset latency ranging from 1 to 4 weeks. Drug-Induced Liver Injury (DILI) often follows an idiosyncratic latency spanning weeks or months, whereas drug-induced malignancies or fibrotic disorders can remain clinically occult for years. Consequently, an adverse event occurring outside the biologically viable latency window weakens suspected relatedness, pointing instead toward alternative clinical factors.

Dechallenge and Rechallenge Dynamics

The dechallenge and rechallenge observation sequence serves as the primary clinical tool for evaluating individual case causality. Dechallenge refers to the clinical trajectory observed when the suspect medicinal product is discontinued or its dosage substantially decreased.

Rechallenge assesses the clinical response following subsequent re-administration of the suspect compound to the same individual.

Clinical Observation Axis	Operational Definition	Representative Clinical Presentation	Impact on Causality Adjudication
Positive Dechallenge	Resolution, significant attenuation, or normalization of the adverse event following drug withdrawal.	Elevated hepatic transaminases return to baseline within two weeks of stopping suspect statin therapy.	Strongly substantiates a causal association; fulfills mandatory criteria for WHO-UMC "Probable" or "Certain".
Negative Dechallenge	Persistence, progression, or deterioration of the adverse manifestation despite complete drug discontinuation.	Acute interstitial nephritis continues to deteriorate unabated following cessation of an NSAID.	Substantially weakens drug etiology; indicates self-sustaining pathology or an independent alternative cause.
Uninformative Dechallenge	Inability to evaluate the independent effect of withdrawal due to confounding variables or irreversible endpoints.	Patient death, concomitant administration of specific antidotes, or events with irreversible tissue destruction.	Precludes definitive dechallenge scoring; limits classification to "Possible" or "Conditional" categories.
Positive Rechallenge	Definite, unambiguous recurrence of the identical clinical reaction following deliberate or inadvertent re-exposure.	Recurrence of fixed drug eruption at the identical anatomical location upon restarting allopurinol.	Provides near-conclusive clinical proof; essential for assigning the WHO-UMC "Certain" classification.
Negative Rechallenge	Absence of event recurrence when the	No recurrence of gastrointestinal distress	Strongly refutes causal relatedness; indicates

Clinical Observation Axis	Operational Definition	Representative Clinical Presentation	Impact on Causality Adjudication
	patient is re-exposed to the drug at therapeutic doses.	upon restarting the medication under identical clinical conditions.	the primary incident was coincidental or linked to other factors.

Deliberate rechallenge entails significant ethical, legal, and clinical risks. It is contraindicated when the initial adverse event involves life-threatening manifestations, such as systemic anaphylaxis, extensive epidermal necrolysis, agranulocytosis, or severe cardiac arrhythmias. Because intentional rechallenge is restricted to non-severe reactions, definitive positive rechallenge documentation is rare in post-marketing spontaneous reporting. As a result, the majority of high-confidence causal assessments in pharmacovigilance remain categorized as "Probable" rather than "Certain".

Confounding Factors: Concomitant Therapy and Host Disease Pathophysiology

Accurate causality assessment requires the methodical identification and systematic exclusion of confounding variables. Confounders typically manifest as underlying patient pathophysiology or concomitant drug exposures.

Underlying patient disease often presents clinical manifestations that mimic genuine drug toxicity. For instance, a reported worsening of renal function in a diabetic patient receiving an angiotensin-converting enzyme (ACE) inhibitor could indicate drug-induced hemodynamic changes, the natural progression of diabetic nephropathy, or acute dehydration from an intercurrent gastrointestinal illness. Differentiating between these causes requires reviewing the complete medical history, baseline laboratory trends, comorbid conditions, and objective diagnostic biomarkers.

Concomitant multi-drug regimens introduce complex diagnostic challenges, particularly in polymorbid geriatric or oncology populations. When an adverse event emerges during polypharmacy, safety evaluators must map the specific administration timeline of each agent, cross-reference their known adverse effect profiles, and analyze potential pharmacokinetic interactions, such as cytochrome P450 inhibition or competitive plasma protein binding. If two or more suspect drugs share a similar temporal onset and toxicity profile, establishing an exclusive causal attribution becomes impossible, requiring both agents to be processed as suspect therapies.

The WHO-UMC Causality Assessment Framework

Conceptual Architecture and Practical Assessment Logic

Developed by the World Health Organization Collaborating Centre for International Drug Monitoring in Uppsala, Sweden, the WHO-UMC causality assessment system is the international standard for evaluating spontaneous adverse event reports. The WHO-UMC framework combines clinical-pharmacological plausibility with the completeness of the case documentation, providing a harmonized qualitative methodology for national pharmacovigilance centers worldwide.

The system organizes clinical judgment through a sequential, criteria-based evaluation rather

than a numerical score. The evaluator reviews the case history across four primary diagnostic axes: temporal plausibility, biological response to drug withdrawal, clinical outcome upon re-administration, and the presence or absence of alternative explanations.

The evaluation process begins with a completeness check of the incoming data. If the case lacks essential clinical details, it is routed into the non-evaluable categories: "Unassessable" when missing or contradictory information cannot be retrieved, or "Conditional" if additional diagnostic data is being gathered. For evaluable reports, the assessor examines the temporal sequence. If the timing is improbable or alternative causes provide a more likely explanation, the case is classified as "Unlikely". When the temporal sequence is reasonable, the assessor reviews dechallenge data and alternative etiologies. Cases with unclear dechallenge outcomes or plausible disease-related explanations are categorized as "Possible". If there is a positive dechallenge and competing causes are unlikely, the report is assigned to "Probable". Finally, if the case includes an unambiguous positive rechallenge and confirms an established pharmacological mechanism with complete documentation, it achieves the highest category of "Certain".

Standardized Diagnostic Category Specifications

| Causality Category | Temporal Sequence Window | Response to Withdrawal (Dechallenge) | Response to Re-exposure (Rechallenge) | Alternative Etiologies and Confounders | Documentation Completeness Requirements | | :--- | :--- | :--- | :--- | :--- | :--- | | **Certain** | Plausible, well-documented interval between drug intake and event onset. | Event abates with a pharmacologically or pathologically plausible recovery profile. | Unambiguous, satisfactory rechallenge confirms recurrence, if clinically appropriate. | Cannot be explained by concurrent medical disease or other pharmaceutical therapies. | Comprehensive; event must be an established pharmacological or phenomenological entity. | | **Probable / Likely** | Reasonable temporal sequence relative to drug administration. | Clinically reasonable improvement or resolution following cessation. | Formal rechallenge information is not required for assignment. | Unlikely to be attributed to the patient's concurrent disease or other therapies. | High; clear clinical chronology without significant missing variables or discrepancies. | | **Possible** | Reasonable temporal sequence relative to drug administration. | Dechallenge data is missing, ambiguous, indeterminate, or uninformative. | Rechallenge information is absent, not attempted, or unevaluated. | Could be equally explained by concurrent medical illness or co-administered drugs. | Moderate; confirms the adverse occurrence, but lacks key clinical diagnostic elements. | | **Unlikely** | Improbable temporal interval between drug use and event manifestation. | Clinical course following withdrawal is non-contributory or discordant. | Re-exposure information is non-contributory or unnecessary. | Underlying disease processes or other drugs provide a more plausible explanation. | Adequate to identify a more convincing alternative pathological cause. | | **Conditional / Unclassified** | Temporal sequence documented, but requires further assessment. | Indeterminate pending acquisition of ongoing patient follow-up. | Indeterminate pending receipt of follow-up clinical documentation. | Cannot be evaluated until supplemental medical tests are reviewed. | Provisional status; essential records are actively sought or currently under review. | | **Unassessable / Unclassifiable** | Report suggests an adverse drug reaction. | Cannot be assessed due to absent or unretrievable documentation. | Cannot be assessed due to lack of verifiable clinical data. | Cannot be determined due to conflicting, contradictory, or absent facts. | Severely defective; documentation is insufficient, inconsistent, and cannot be supplemented. |

Adjudicating Drug-Drug Interactions (DDIs)

The WHO-UMC framework evaluates drug-drug interactions using an actor-victim paradigm. Within this model, the victim drug is the compound whose pharmacokinetic disposition (absorption, distribution, metabolism, elimination) or pharmacodynamic activity is altered, leading to therapeutic failure or toxic accumulation. The actor drug is the newly introduced or dose-adjusted compound that modifies the victim drug's metabolic clearance or receptor sensitivity.

When assessing causality in an interaction-mediated adverse event, the evaluator examines the temporal relationship relative to the introduction or dose change of the actor drug, rather than the baseline victim therapy. For example, when a patient on stable warfarin anticoagulation develops an elevated International Normalized Ratio (INR) and spontaneous hemorrhage shortly after starting amiodarone, amiodarone functions as the pharmacokinetic actor through potent CYP2C9 inhibition. Causality is evaluated based on the timing of the actor drug's introduction, the biological latency of enzymatic inhibition, and the clinical recovery observed upon dechallenge of the actor agent.

Algorithmic Assessment: The Naranjo ADR Probability Scale

Ten-Point Scoring Methodology

Formulated by Naranjo and colleagues in 1981, the Naranjo Algorithm provides an objective, questionnaire-driven scoring system to determine adverse drug reaction probability. The tool uses ten targeted clinical questions, assigning numerical weights ranging from -1 to +2 based on standardized responses ("Yes," "No," or "Do Not Know / Not Done"). This structure minimizes subjective observer bias and improves reproducibility across different safety evaluators.

Query Identifier	Algorithmic Assessment Query	Response Weight: Yes	Response Weight: No	Response Weight: Do Not Know / Not Done
Q1	Are there previous conclusive reports on this reaction?	+1	0	0
Q2	Did the adverse event appear after the suspected drug was administered?	+2	-1	0
Q3	Did the adverse reaction improve when the drug was discontinued or a specific antagonist was administered?	+1	0	0
Q4	Did the adverse reaction reappear	+2	-1	0

Query Identifier	Algorithmic Assessment Query	Response Weight: Yes	Response Weight: No	Response Weight: Do Not Know / Not Done
	when the drug was readministered?			
Q5	Are there alternative causes (other than the drug) that could on their own have caused the reaction?	-1	+2	0
Q6	Did the reaction reappear when a placebo was given?	-1	+1	0
Q7	Was the drug detected in blood (or other fluids) in concentrations known to be toxic?	+1	0	0
Q8	Was the reaction more severe when the dose was increased, or less severe when the dose was decreased?	+1	0	0
Q9	Did the patient have a similar reaction to the same or similar drugs in any previous exposure?	+1	0	0
Q10	Was the adverse event confirmed by any objective evidence?	+1	0	0

Mathematical Stratification and Probability Tiers

The composite Naranjo probability score is derived through the linear summation of the ten scored parameters:

$$\text{Naranjo Score} = \sum_{i=1}^{10} Q_i$$

The resulting score places the adverse event into one of four probability categories:

- **Definite (Score ≥ 9):** The clinical event occurs within a plausible temporal sequence following drug intake, improves upon dechallenge, recurs on rechallenge, is

substantiated by objective laboratory or clinical evidence, and has alternative etiologies ruled out.

- **Probable (Score = 5 to 8):** The event follows a reasonable temporal sequence, shows a clear clinical improvement upon drug cessation, is supported by objective findings, and is unlikely to be driven by alternative etiologies.
- **Possible (Score = 1 to 4):** The event demonstrates an acceptable temporal sequence, but documentation regarding dechallenge is incomplete or confounded, and competing disease states or co-medications could plausibly account for the presentation.
- **Doubtful (Score ≤ 0):** The event fails to align with the expected temporal sequence, does not resolve upon dechallenge, or is more convincingly explained by the patient's underlying clinical condition.

Comparative Analysis: WHO-UMC Framework versus Naranjo Nomogram

Operational Domain	WHO-UMC Causality System	Naranjo Algorithm Scale
Underlying Approach	Qualitative, expert-judgment framework based on clinical reasoning.	Structured, deterministic questionnaire using categorical point scoring.
Classification Strategy	Direct criteria matching against defined category descriptions.	Cumulative numerical scoring mapped to defined probability tiers.
Inter-rater Reliability	Moderate; influenced by individual clinical judgment and experience.	High; objective questions reduce scoring variability between evaluators.
Operational Efficiency	High; streamlined for large-scale post-marketing surveillance.	Moderate; requires evaluating ten specific questions per suspect event.
Primary Environment	National pharmacovigilance centers and global post-marketing databases.	Randomized clinical trials, academic case reports, and controlled studies.
Handling Polypharmacy & DDIs	Flexible; accommodates complex actor-victim drug-drug interactions.	Limited; penalizes co-medications and lacks parameters for multi-drug interactions.
Specialized Populations	Broadly applicable across diverse patient cohorts and disease contexts.	Less reliable in pediatric, critically ill, and multi-morbid populations.

Vaccine Pharmacovigilance: The Revised WHO-AEFI Causality Framework

Distinctions in Vaccine Safety Evaluation

Vaccine pharmacovigilance presents distinct immunological, epidemiological, and operational considerations compared to small-molecule safety surveillance. Vaccines are administered primarily to healthy individuals, often infants, children, and pregnant populations, as preventive

public health measures, creating low societal tolerance for adverse events. Unlike daily maintenance pharmacotherapies, immunization typically involves isolated single doses or spaced prime-boost regimens, making classic dechallenge and rechallenge observations rare. Furthermore, mass immunization programs often coincide with periods in early life when infectious illnesses, developmental disorders, and congenital conditions naturally emerge. This temporal overlap creates substantial potential for coincidental associations. In addition, vaccines are complex biological formulations susceptible to cold-chain excursions, reconstitution mistakes, and injection-site complications. These characteristics require specialized causality assessment frameworks, codified in the World Health Organization User Manual for AEFI Causality Assessment (ISBN 9789241516990).

The Four-Stage AEFI Investigation Lifecycle

Under the revised WHO-AEFI guidelines, causality assessment follows a standardized four-stage investigation lifecycle.

Stage 1 covers Eligibility Screening, which confirms that the safety report meets basic validity standards (identifiable patient, identifiable vaccinator or reporter, documented vaccine, and adverse outcome) and verifies that the adverse event has a confirmed clinical or laboratory diagnosis. If the clinical diagnosis remains unconfirmed, formal causality assessment is deferred.

Stage 2 involves Checklist Evaluation, where evaluators assess the case against four key criteria: identifying known alternative causes for the condition, confirming established biological mechanisms based on literature, verifying that the event onset falls within a plausible temporal window, and assessing potential cold-chain failures, programmatic deviations, or administration errors.

Stage 3 applies Algorithmic Integration, guiding the evaluator through a structured decision logic matrix to reconcile contradictory clinical findings, assess the balance of evidence, and determine primary etiological attribution.

Stage 4 completes the process with Final Category Assignment, classifying the case into standardized categories—Consistent, Indeterminate, Inconsistent, or Unclassifiable—to support domestic and international immunization surveillance.

Classification Typology of the WHO-AEFI System

Primary Category	Sub-Category Identifier	Etiological Description	Representative Clinical Presentation
A. Consistent with causal association to immunization	A1: Vaccine product-related reaction	Caused or precipitated by the intrinsic biological or chemical components of the vaccine.	Extensive local erythema, high post-vaccination fever, or vaccine-strain virus disease.
	A2: Vaccine quality defect-related reaction	Caused by quality defects in the formulation, vial integrity, or delivery device.	Sterile local abscess formation resulting from inadequate dissolution of aluminum adjuvant.
	A3: Immunization	Resulting from	Injection-site sterile

Primary Category	Sub-Category Identifier	Etiological Description	Representative Clinical Presentation
	error-related reaction	inappropriate handling, storage, reconstitution, or administration errors.	abscess or tissue damage caused by subcutaneous instead of intramuscular injection.
	A4: Immunization anxiety-related reaction	Arising from anxiety or psychological stress related to the immunization procedure.	Vasovagal syncope, hyperventilation, or stress-induced fainting in adolescents following injection.
B. Indeterminate	B1: Consistent temporal relationship, insufficient definitive evidence	Plausible temporal latency, but clinical evidence is insufficient to confirm a causal link.	A newly recognized inflammatory or neurological event occurring within a biologically plausible window.
	B2: Conflicting evidence for and against association	Clinical or epidemiological evidence presents contradictory indications of causality.	Post-immunization encephalopathy in an infant who also tests positive for enterovirus infection.
C. Inconsistent with causal association to immunization	Coincidental event	Driven by an underlying hereditary, infectious, or environmental condition.	Bacterial meningitis developing 24 hours after immunization, confirmed via positive CSF bacterial culture.
D. Unclassifiable	Unclassifiable event	Essential clinical, laboratory, or epidemiological information is permanently missing.	Severe post-immunization collapse reported without patient details, clinical outcomes, or follow-up.

Host Predisposition Factors in Vaccine Causality

Accurate vaccine causality adjudication requires evaluating individual host characteristics and underlying medical conditions. Host susceptibility factors can alter immunological responses, predisposing individuals to adverse reactions that would not occur in healthy populations. Primary or acquired immunodeficiencies represent critical host considerations. Patients with congenital T-cell or B-cell defects, or those receiving immunosuppressive therapies, face elevated risks of disseminated disease following live-attenuated viral or bacterial vaccines. For example, administering the live BCG vaccine to an infant with unrecognized Severe Combined Immunodeficiency (SCID) can lead to disseminated BCG-itis, whereas oral poliovirus vaccine (OPV) can trigger vaccine-associated paralytic poliomyelitis (VAPP).

Underlying autoimmune traits can similarly influence post-vaccination outcomes. Individuals with genetic predispositions, such as specific human leukocyte antigen (HLA) alleles, may

experience heightened systemic inflammatory cascades or transient disease flares following the immune stimulation induced by adjuvanted formulations.

Genetic channelopathies also intersect with post-vaccination responses. For instance, children with pathogenic mutations in the *SCN1A* sodium channel gene (associated with Dravet syndrome) frequently present with their first prolonged febrile seizure following standard childhood vaccinations. In these patients, the physiological fever triggered by the vaccine acts as a secondary biological stressor that unmasks an underlying genetic disorder, rather than the vaccine functioning as the primary neurotoxic cause.

Industrial Implementation, Medical Review, and Regulatory Reporting

ICSR Lifecycle Management: From Day Zero to Triage

In clinical safety operations, Individual Case Safety Report (ICSR) processing follows strict regulatory timelines. The regulatory compliance clock begins on Day Zero, defined as the calendar day an employee, vendor, or authorized representative of a Marketing Authorization Holder (MAH) becomes aware of an adverse event containing the four minimum validity criteria. These criteria require an identifiable patient, an identifiable reporter, at least one suspect medicinal product, and at least one documented adverse event.

During the initial triage and case intake phase, reports are evaluated against standard seriousness criteria under ICH E2A and E2D guidelines. An adverse event is classified as serious if it results in death, is life-threatening, necessitates inpatient hospitalization or prolongation of existing hospitalization, results in persistent or significant disability or incapacity, causes a congenital anomaly or birth defect, or is considered an Important Medical Event (IME) that requires intervention to prevent one of these outcomes. Serious adverse reactions require expedited regulatory submission, with fatal or life-threatening Suspected Unexpected Serious Adverse Reactions (SUSARs) mandated within 7 calendar days, and all other serious unexpected reactions required within 15 calendar days.

Multi-Tier to Binary Translation for Regulatory Compliance

Safety surveillance databases—such as Oracle Argus Safety and Veeva Vault Safety—track multi-tier causality classifications, including WHO-UMC categories and numerical Naranjo scores. However, regulatory authorities, including the United States Food and Drug Administration (FDA), the European Medicines Agency (EMA), and the United Kingdom Medicines and Healthcare products Regulatory Agency (MHRA), mandate a binary causality decision (Related vs. Not Related) for individual case processing and expedited submissions.

Internal Granular Causality Tier	Regulatory Binary Adjudication	Expedited Reporting Impact	Aggregate Safety Impact
WHO-UMC: Certain Naranjo: Definite (≥ 9)	Related (Reasonable Causal Possibility).	Mandates expedited 7-day or 15-day submission if serious and unexpected.	Compiled as confirmed reaction in PSUR/PBRER reports.
WHO-UMC: Probable / Likely Naranjo:	Related (Reasonable Causal Possibility).	Mandates expedited 7-day or 15-day	Analyzed in aggregate benefit-risk and signal

Internal Granular Causality Tier	Regulatory Binary Adjudication	Expedited Reporting Impact	Aggregate Safety Impact
Probable (5 - 8)		submission if serious and unexpected.	evaluations.
WHO-UMC: Possible Naranjo: Possible (1 - 4)	Related (Reasonable Causal Possibility).	Mandates expedited 7-day or 15-day submission if serious and unexpected.	Tracked for emerging safety signals and pattern changes.
WHO-UMC: Unlikely Naranjo: Doubtful (1e 0)	Not Related (No Causal Relationship).	Excluded from expedited reporting; does not trigger 7-day or 15-day alert.	Documented in periodic aggregate reports as unrelated AEs.
WHO-UMC: Conditional / Unclassified	Provisional / Conservative Related.	Conservatively reported as Related if serious and unexpected, pending queries.	Targeted for regulatory follow-up to acquire missing clinical data.
WHO-UMC: Unassessable	Unclassifiable.	Non-reportable if core validity criteria cannot be confirmed.	Retained in safety databases; excluded from signal metrics.

A central tenet of international regulatory harmonization is that when a primary healthcare professional reporter attributes causality to a suspect drug, the sponsor cannot downgrade the case to "Not Related" for expedited reporting purposes. Although a pharmaceutical sponsor's safety physician may document an internal assessment of "Not Related" in the company comment section, the primary reporter's positive causality determination governs regulatory submission clocks. This regulatory rule prevents pharmaceutical manufacturers from using internal medical opinions to avoid mandatory 15-day expedited reporting obligations.

The Medical Reviewer and Safety Physician Mandate

Medical reviewers and safety physicians provide specialized clinical evaluation throughout the pharmacovigilance lifecycle. They evaluate complex clinical data, assess multi-drug regimens, interpret laboratory trends, and determine the company's formal position on product causality. A core responsibility of the safety physician is distinguishing between Labeledness, Expectedness, and Listedness. Labeledness reflects whether the adverse event is documented in local, market-specific Prescribing Information, such as the USPI in the United States or the SmPC across the European Union. Expectedness applies to clinical development, determining whether an event matches the character, severity, and frequency described in the clinical study protocol or Investigator's Brochure (IB). Listedness evaluates whether the event appears in the company's global reference safety document, the Company Core Data Sheet (CCDS). These distinctions, combined with the binary causality determination, establish whether an adverse event qualifies as an unexpected adverse reaction requiring expedited regulatory distribution. Safety physicians also draft the final medical narrative, synthesizing patient medical history, drug exposure dates, event latency, and laboratory findings into a clear, cohesive clinical summary. In aggregate safety surveillance, medical reviewers monitor recurring trends across Periodic Benefit-Risk Evaluation Reports (PBRERs) and Development Safety Update Reports (DSURs) to detect emerging safety signals and evaluate the long-term risk profile of the

medicinal product.

Practical Case Scenarios and Worked Adjudications

Scenario 1: Amoxicillin-Associated Cutaneous Eruption

A 32-year-old female without significant past medical history was prescribed amoxicillin 500 mg orally three times daily for acute streptococcal pharyngitis. On day 8 of therapy, she developed an extensive, pruritic, symmetric maculopapular exanthem across her trunk and extremities. She was taking no other medications. Amoxicillin was immediately discontinued, and oral cetirizine was administered. Within 72 hours of cessation, the rash faded significantly, resolving completely by day 5 post-discontinuation. No rechallenge was performed.

Under the WHO-UMC system, this event is adjudicated as **Probable / Likely**. The case demonstrates a reasonable temporal sequence, with the rash emerging on day 8 of antibiotic exposure. The clinical trajectory shows a positive dechallenge, as the rash improved markedly within 72 hours of drug cessation. Streptococcal pharyngitis rarely produces a delayed maculopapular rash on its own, and the absence of concurrent therapies makes alternative causes unlikely. The case cannot be classified as "Certain" because deliberate rechallenge was not attempted and formal immunological testing was not performed.

Evaluating the case with the Naranjo algorithm yields a composite score of +7[span_129](start_span)[span_129](end_span)[span_144](start_span)[span_144](end_span)[span_159](start_span)[span_159](end_span), categorizing the reaction as a **Probable ADR**.

Point allocation across the 10 parameters includes: +1 for Question 1 (amoxicillin exanthem is well documented in published literature);

+2[span_130](start_span)[span_130](end_span)[span_145](start_span)[span_145](end_span)[span_160](start_span)[span_160](end_span) for Question 2 (the eruption appeared after drug administration); +1 for Question 3 (clinical resolution followed drug cessation); 0 for Question 4 (no rechallenge was attempted); +2 for Question 5 (alternative clinical causes were ruled out); 0 for Question 6 (no placebo control was administered); 0 for Question 7 (serum drug levels were not measured); 0 for Question 8 (the dose was not adjusted prior to cessation); 0 for Question 9 (no previous exposure history was recorded); and +1 for Question 10 (the eruption was verified by objective clinical examination).

Scenario 2: Atorvastatin and Clarithromycin Pharmacokinetic Interaction Induced Rhabdomyolysis

A 64-year-old male maintained on atorvastatin 80 mg daily for two years for familial hypercholesterolemia was diagnosed with community-acquired pneumonia and prescribed clarithromycin 500 mg twice daily. Five days later, he presented to the emergency department with severe, diffuse muscle pain, proximal muscle weakness, and tea-colored urine. Diagnostic testing revealed a marked elevation in serum creatine kinase (CK) at 28,500 IU/L (reference range: 30 - 200\text{ IU/L}) and acute kidney injury, with serum creatinine rising to 2.8\text{ mg/dL} from a baseline of 1.0\text{ mg/dL}. Both atorvastatin and clarithromycin were immediately stopped, and aggressive intravenous hydration was initiated. Over the subsequent 10 days, his myalgia resolved, CK concentrations declined to 420\text{ IU/L}, and renal function returned to baseline. Three months later, low-dose atorvastatin (10 mg daily) was reintroduced without clarithromycin, and the patient remained asymptomatic without CK elevation.

In this drug-drug interaction case, the WHO-UMC system is applied using the actor-victim paradigm. Clarithromycin is designated as the pharmacokinetic actor drug, adjudicated as **Probable / Likely**. Atorvastatin represents the victim compound, also adjudicated as **Probable / Likely**. While the patient had tolerated high-dose atorvastatin for two years, the introduction of clarithromycin—a potent CYP3A4 and OATP1B1 inhibitor—significantly reduced atorvastatin clearance, leading to toxic systemic accumulation. The temporal latency of 5 days matches the expected timeframe for enzymatic inhibition and statin accumulation. Discontinuing both agents resulted in a positive dechallenge, and subsequent re-exposure to low-dose atorvastatin in the absence of the CYP3A4 inhibitor produced a negative rechallenge, confirming the interaction-driven etiology. Because the reaction required inpatient hospitalization and involved acute kidney injury, the event meets regulatory seriousness criteria, mandating a 15-day expedited submission to health authorities.

Scenario 3: Post-Immunization Syncope versus Systemic Anaphylaxis

A 15-year-old female received her first dose of the quadrivalent Human Papillomavirus (HPV) recombinant vaccine at a school-based immunization clinic. Four minutes after intramuscular administration, while remaining seated in the post-vaccination observation area, she became pale, experienced diaphoresis, and had a brief 20-second episode of unconsciousness. Clinical assessment immediately following the event identified bradycardia with a heart rate of 52 bpm, blood pressure of 88/54 mmHg, and cool, clammy extremities. The patient exhibited no urticaria, facial angioedema, wheezing, stridor, or respiratory compromise. She was placed in the supine position with her lower extremities elevated. Within 5 minutes, her consciousness returned, blood pressure normalized to 110/70 mmHg, and her heart rate stabilized at 76 bpm. After 45 minutes of observation, she was discharged without ongoing symptoms.

Under the revised WHO-AEFI classification framework, this reaction is categorized as **Category A4 – Immunization Anxiety-Related Reaction**. The rapid onset, transient loss of consciousness, bradycardia, hypotension, and swift recovery upon assume a supine position represent classic vasovagal neurocardiogenic syncope triggered by the stress and anxiety of the injection procedure. The complete absence of cutaneous signs, such as urticaria or angioedema, alongside normal respiratory function, definitively excludes an IgE-mediated vaccine product reaction (Category A1). The adverse event was driven by the physiological response to the injection process rather than the biological contents of the vaccine.

Operational Pitfalls and Professional Assessment Guidelines

Common Analytical and Operational Errors in Case Processing

A common operational error in clinical safety processing involves conflating a successful medical intervention with a positive dechallenge. When an adverse event abates only following the administration of an antidote, systemic corticosteroids, or aggressive clinical therapy, the improvement cannot be attributed solely to the withdrawal of the suspect drug. Scoring this clinical trajectory as an unconfounded dechallenge overstates the certainty of drug-relatedness; it should instead be documented as an assisted dechallenge.

Case processors also frequently default to the "Unassessable" category when secondary clinical

details are missing. If a safety report satisfies the four minimum validity criteria and provides a clear temporal sequence, lacking information about previous literature or dechallenge should not lead to an unassessable designation. The evaluator should assign the case to the "Conditional" category while follow-up information is actively pursued, or classify it as "Possible" if the temporal relationship is reasonable.

In vaccine safety assessments, overlooking individual host risk factors can lead to misattribution. Classifying an adverse event as entirely coincidental without assessing host characteristics, such as underlying immune deficiencies or genetic variants, can obscure important vaccine-host biological interactions.

Finally, applying the Naranjo algorithmic tool in complex polypharmacy settings often introduces systematic scoring distortion. In elderly, oncology, or intensive-care patients receiving complex multi-drug regimens, the Naranjo tool frequently assigns artificially low scores because points are deducted for alternative explanations and co-medications. In these clinical settings, qualitative expert evaluation via the WHO-UMC framework offers greater flexibility and clinical accuracy.

Core Principles for Regulatory Audits and Professional Compliance

Causality conclusions must be supported by verifiable documentation within the safety narrative. Regulatory inspectors regularly audit ICSR files, and vague medical opinions lacking objective clinical data, clear temporal chronologies, or relevant laboratory results will not withstand scrutiny during pharmacovigilance inspections.

Safety departments must maintain strict regulatory adherence regarding primary reporter opinions. An internal company assessment of "Not Related" cannot override a reporting healthcare professional's suspected causality to avoid expedited reporting clocks. When a reporting physician suspects a causal link, the report must be handled under expedited submission guidelines.

Causality assessment is an iterative, dynamic process throughout the case safety lifecycle. Cases initially categorized as "Conditional" must be promptly re-evaluated upon receipt of supplemental medical records, discharge summaries, or laboratory tests. Regular updates to causality assignments ensure that cumulative safety databases reflect accurate clinical information, supporting reliable signal detection and protecting patient safety across the entire pharmaceutical lifecycle.

Works cited

1. Certificate Course in Pharmacovigilance - VigiServe Foundation, <https://www.vigiservefoundation.org/courses/ccp/>
2. What is Causality assessment in PV? | Q17 - YouTube, https://www.youtube.com/watch?v=P03_DdaXjH4
3. WHO Causality Assessment Scale Guide | PDF - Scribd, <https://www.scribd.com/document/896932093/WHO-CAUSALITY-SCALE-1>
4. The use of the WHO-UMC system for standardised case causality, https://lms.thsti.in/wp-content/uploads/2024/04/WHO_UMC-system-for-standardised-case-causality-assessment.pdf
5. Causality Assessment - YouTube, <https://www.youtube.com/watch?v=6gH4AWnwTDI>
6. What is Causality Assessment in Pharmacovigilance?, <https://www.ddregpharma.com/what-is-causality-assessment-in-pharmacovigilance>
7. Naranjo Scale - Causality assessment Method - YouTube,

<https://www.youtube.com/watch?v=sTPLPH4EigA> 8. Pharmacovigilance ICSR (Individual Case Safety Report) - YouTube, <https://www.youtube.com/watch?v=PdxzXNF-xbl> 9. Medical Review – Basics - VigiServe Foundation, <https://www.vigiservefoundation.org/courses/medical-review-basics/> 10. Basic Certification Course in Pharmacovigilance, <https://www.vigiservefoundation.org/courses/pharmacovigilance-basics-industry/> 11. Signal Management Process and WHO UMC Concepts - YouTube, https://www.youtube.com/watch?v=k_e6iKo3l_A 12. What is Naranjo Scale ? | Casualty Assessment | MCQ - YouTube, <https://www.youtube.com/watch?v=ayeywcC3ca0> 13. Causality Assessment | Pharmacovigilance Interview Questions, https://www.youtube.com/watch?v=UGp6_-JYvMM 14. Safety profile following mRNA-1273 COVID-19 primary vaccination, <https://pmc.ncbi.nlm.nih.gov/articles/PMC13255034/> 15. Efficacy and Feasibility of SMS M-Health for the Detection of, <https://www.preprints.org/manuscript/202303.0523> 16. A Two-Year Pharmacovigilance Analysis of Adverse Drug Reactions, <https://www.mdpi.com/2077-0383/15/2/848> 17. evsjv`k †M‡RU - Department of Printing and Publications, https://www.dpp.gov.bd/upload_file/gazettes/39351_86883.pdf 18. DENUNCIA PENALE STRAEPOCALE (file traducibile, <https://salvatoreraino.com/denuncia-penale-straepocale/> 19. Causality assessment of an adverse event following immunization, <https://www.who.int/publications/i/item/9789241516990> 20. Seriousness Assessment in Pharmacovigilance - YouTube, <https://www.youtube.com/watch?v=tb613EiaZ9U> 21. WHO-UMC Causality Assessment Scale - CLINI INDIA, <https://www.cliniindia.com/who-umc-causality-assessment-scale/>