2015. Average Height of Buildings in Each Segment Premium Medium ♥ Topics 🖫 Companies 🗘 Hint A perfectly straight street is represented by a number line. The street has building(s) on it and is represented by a 2D integer array buildings, where buildings[i] = [starti, endi, heighti]. This means that there is a building with height; in the half-closed segment [start; end;). You want to describe the heights of the buildings on the street with the minimum number of non-overlapping segments. The street can be represented by the 2D integer array street where street[j] = [leftj, rightj, averagej] describes a half-closed segment [left_j, right_j) of the road where the average heights of the buildings in the segment is average_j. • For example, if buildings = [[1,5,2],[3,10,4]], the street could be represented by street = [[1,3,2],[3,5,3],[5,10,4]] because: • From 1 to 3, there is only the first building with an average height of 2 / 1 = 2. • From 3 to 5, both the first and the second building are there with an average height of (2+4) / 2 = 3. • From 5 to 10, there is only the second building with an average height of 4 / 1 = 4. Given buildings, return the 2D integer array street as described above (excluding any areas of the street where there are no buldings). You may return the array in any order. The average of n elements is the sum of the n elements divided (integer division) by n. A half-closed segment [a, b) is the section of the number line between points a and b including point a and not including point b. Example 1: **Input:** buildings = [[1,4,2],[3,9,4]]**Output:** [[1,3,2],[3,4,3],[4,9,4]] Explanation: From 1 to 3, there is only the first building with an average height of 2 / 1 = 2. From 3 to 4, both the first and the second building are there with an average height of (2+4) / 2 = 3. From 4 to 9, there is only the second building with an average height of 4 / 1 = 4. Example 2: **Input:** buildings = [[1,3,2],[2,5,3],[2,8,3]] **Output:** [[1,3,2],[3,8,3]] Explanation:

```
From 1 to 2, there is only the first building with an average height of 2 / 1 = 2.
From 2 to 3, all three buildings are there with an average height of (2+3+3) / 3 = 2.
From 3 to 5, both the second and the third building are there with an average height of (3+3) / 2 = 3.
From 5 to 8, there is only the last building with an average height of 3 / 1 = 3.
The average height from 1 to 3 is the same so we can group them into one segment.
The average height from 3 to 8 is the same so we can group them into one segment.
```

Example 3:

```
Output: [[1,2,1],[5,6,1]]
Explanation:
From 1 to 2, there is only the first building with an average height of 1 / 1 = 1.
From 2 to 5, there are no buildings, so it is not included in the output.
From 5 to 6, there is only the second building with an average height of 1 / 1 = 1.
We cannot group the segments together because an empty space with no buildings seperates the segments.
```

- **Constraints:**
- 0 <= start_i < end_i <= 10⁸

Seen this question in a real interview before? 1/5

• 1 <= buildings.length <= 10⁵

Input: buildings = [[1,2,1],[5,6,1]]

• buildings[i].length == 3

• 1 <= height_i <= 10⁵

Yes No Submissions 3.8K Acceptance Rate 59.8% Accepted 2.3K ♥ Topics

Array Greedy Sorting Heap (Priority Queue) Companies 0 - 6 months Microsoft 2

O Hint 1 Try sorting the start and end points of each building.

O Hint 2 The naive solution is to go through each position on the street and keep track of the sum of all the buildings at that position and the number of buildings at that position.

O Hint 3 How could we optimize that solution to pass?

O Hint 4 We don't need to go through every position, just the ones where a building starts or a building ends.

₹ Similar Questions

Average Waiting Time Describe the Painting

Amount of New Area Painted Each Day 🏠 Divide Intervals Into Minimum Number of Groups

Copyright © 2024 LeetCode All rights reserved

Discussion (0)