

339. Nested List Weight Sum Premium

Medium

Topics

Companies

You are given a nested list of integers `nestedList`. Each element is either an integer or a list whose elements may also be integers or other lists.

The **depth** of an integer is the number of lists that it is inside of. For example, the nested list `[1, [2, 2], [[3], 2], 1]` has each integer's value set to its **depth**.

Return the sum of each integer in `nestedList` multiplied by its **depth**.

Example 1:

`nestedList = [[1, 1], 2, [1, 1]]`
`depth = [2, 2, 1, 2, 2]`

Input: `nestedList = [[1,1],2,[1,1]]`

Output: 10

Explanation: Four 1's at depth 2, one 2 at depth 1. $1*2 + 1*2 + 2*1 + 1*2 + 1*2 = 10$.

Example 2:

`nestedList = [1, [4, [6]]]`
`depth = [1, 2, 3]`

Input: `nestedList = [1,[4,[6]]]`

Output: 27

Explanation: One 1 at depth 1, one 4 at depth 2, and one 6 at depth 3. $1*1 + 4*2 + 6*3 = 27$.

Example 3:

Input: `nestedList = [0]`

Output: 0

Constraints:

- $1 \leq \text{nestedList.length} \leq 50$
- The values of the integers in the nested list is in the range $[-100, 100]$.
- The maximum **depth** of any integer is less than or equal to 50.

Seen this question in a real interview before? 1/5

Yes

No

Accepted 317.8K | Submissions 375.9K | Acceptance Rate 84.6%

Topics

Depth-First Search

Breadth-First Search

Companies

0 - 3 months

Meta 64

Amazon 2

0 - 6 months

LinkedIn 4

6 months ago

Apple 2

eBay 2

Similar Questions

Nested List Weight Sum II 🔒

Medium

Array Nesting

Medium

Employee Importance

Medium

Discussion (31)