Pair of linear equation in two variables

harshita (paidisettyharshita@sriprakashschools.com)

July 19, 2023

Class 10^{th} Maths - Chapter 3

1. On comparing $\frac{a_1}{a_2}$, $\frac{b_1}{b_2}$ $\frac{c_1}{c_2}$ Find out whether the following pair of linear equation are consistent, or inconsistent.

$$5x - 3y = 11; (1)$$

$$-10x + 6y = 22 (2)$$

Solution:

equations can be written as:

$$\begin{pmatrix} 5 & -3 \\ -10 & 6 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 11 \\ 22 \end{pmatrix} \tag{3}$$

The values of x and y are:

$$x = \frac{\begin{vmatrix} b & a2 \end{vmatrix}}{\begin{vmatrix} a1 & a2 \end{vmatrix}} = \frac{\begin{vmatrix} 11 & 22 \\ -3 & 6 \end{vmatrix}}{\begin{vmatrix} 5 & -10 \\ -3 & 6 \end{vmatrix}} = \frac{\begin{vmatrix} 11*-3 - |22*6|}{|5*6 - |-10*-3|} = \frac{66-66}{30-30} = \frac{0}{0}$$
(5)

1

$$y = \frac{\begin{vmatrix} a1 & b \\ 1a1 & a2 \end{vmatrix}}{\begin{vmatrix} a1 & a2 \end{vmatrix}} = \frac{\begin{vmatrix} 5 & -10 \\ 11 & 22 \end{vmatrix}}{\begin{vmatrix} 5 & -10 \\ -3 & 6 \end{vmatrix}} = \frac{\begin{vmatrix} 5 \times 22 \\ -10 \times 11 \end{vmatrix}}{\begin{vmatrix} 6 \times 5 \\ -10 \times -3 \end{vmatrix}} = \frac{-10 + 10}{30 - 30} = \frac{0}{0}$$
(6)

The solution derived is cosistent as the ${\bf x}$ and ${\bf y}$ values are undefined so, no solution exsists.