Name: Harshit Jain Access ID: hmj5262

Recitation: 8

Problem 0 Points:

Acknowledgements

- (a) I did not work in a group.
- (b) I did not consult without anyone my group members.
- (c) I did not consult any non-class materials.

Problem 1

Points:

Analyze Running Time

1. **Runtime:** $\theta(n^2)$

When i = 1, then j = 1. Here, while loop runs for $\frac{n-1}{5}$ times.

When i = 2, then j = 2. Here, while loop runs for $\frac{n-2}{5}$ times.

:

When i = (n-5), then j = (n-5). Here, while loop runs for 1 time.

:

When i = (n-1), then j = (n-1). Here, while loop runs for 1 time.

Total time: $\frac{n-1}{5} + \frac{n-2}{5} + \dots + \frac{5}{5} + 1 + 1 + 1 + 1 + 1 \Rightarrow \theta(n^2)$

2. **Runtime:** $\theta(n^2)$

When i = 1, then while loop runs from 4 to n. Time = (n - 4)

When i = 2, then while loop runs from 8 to n. Time = (n - 8)

:

When $i = \frac{n}{4}$, then while loop will run 1 time. Time = 1

For loop will run for total of n times.

$$\Rightarrow \theta(n^2)$$

3. **Runtime:** $\theta(nlogn)$

The main while loop will run for $\theta(logn)$ time as *i* is reduced to its half after every iteration.

The inner for loop will run for $\frac{n}{2} + \frac{n}{2^2} + \frac{n}{2^3} + \cdots + 1 = \theta(n)$

So, total time $\Rightarrow \theta(nlogn)$

Problem 2

Points:

Polynomials and Horner's rule

$$P(x) = a_0 + a_1 x^1 + a_2 x^2 + \dots + a_n x^n; x = x_0$$

(a) For Addition: The Brute Force Algorithm will perform $\bf n$ additions in total.

So, Time taken for additions = O(n)

For Multiplication: $P(x) = a_0 + [a_1 \cdot x_0] + [a_2 \cdot x_2 \cdot x_2] + \dots + [a_n \cdot x_n \cdot x_n$

So, Time taken = $T_n = 1 + 2 + 3 + \dots + n$

$$= \frac{n(n+1)}{2}$$
$$= \frac{n^2}{2} + \frac{n}{2}$$

$$= O(n^2)$$

Therefore, total time complexity for Brute Force Algorithm = $O(n^2) + O(n) = O(n^2)$

- (b)
- (c)

Problem 3 Points:

Solving recurrences

(a) Height = $\log_2 n$

Branching factor = 2

The size of the sub-problems at depth $k = \frac{n}{2^k}$

Number of sub-problems at depth $k = 2^k$

Total work done =

$$\sum_{k=0}^{\log_2 n} (2^k) \cdot (\frac{n}{2^k})^{\frac{1}{2}} \Rightarrow \sqrt{n} \cdot \sum_{k=0}^{\log_2 n} (2^{\frac{k}{2}}) \Rightarrow \sqrt{n} \cdot \theta(\sqrt{n}) \Rightarrow \theta(n)$$

(Note that, $\sum_{k=0}^{\log_2 n} (2^{\frac{k}{2}})$ is a geometric series $=2^{\frac{0}{2}}+2^{\frac{1}{2}}+\cdots+2^{\frac{\log_2 n}{2}}=\Theta(\sqrt{n})$)

(b) Height = $\log_3 n$

Branching factor = 2

The size of the sub-problems at depth $k = \frac{n}{3^k}$

Number of sub-problems at depth $k = 2^k$

Total work done =

$$\sum_{k=0}^{\log_3 n} (2^k) \cdot (\frac{n}{3^k})^0 \Rightarrow \sum_{k=0}^{\log_3 n} (2^k) \Rightarrow 2^0 + 2^1 + 2^3 + \dots + 2^{\log_3 n} \Rightarrow \Theta(n^{\log_2 3})$$

(c) Height = $\log_4 n$

Branching factor = 5

The size of the sub-problems at depth $k = \frac{n}{4^k}$

Number of sub-problems at depth $k = 5^k$

Total work done =

$$\sum_{k=0}^{\log_4 n} (5^k) \cdot (\frac{n}{4^k})^1 \Rightarrow n \cdot \sum_{k=0}^{\log_4 n} ((\frac{5}{4})^k) \Rightarrow n \cdot ((\frac{5}{4})^0 + (\frac{5}{4})^1 + \dots + (\frac{5}{4})^{\log_4 n}) \Rightarrow n \cdot \theta(\frac{5^{\log_4 n}}{n}) \Rightarrow \theta(n^{\log_5 4})$$

(d) Height = $\log_7 n$

Branching factor = 7

The size of the sub-problems at depth $k = \frac{n}{7^k}$

Number of sub-problems at depth $k = 7^k$

Total work done =

$$\sum_{k=0}^{\log_7 n} (7^k) \cdot (\frac{n}{7^k})^1 \Rightarrow n \cdot \sum_{k=0}^{\log_7 n} 1 \Rightarrow n \cdot \log n \Rightarrow \theta(n \log n)$$

(e) Height = $\log_3 n$

Branching factor = 9

The size of the sub-problems at depth $k = \frac{n}{3^k}$

Number of sub-problems at depth $k = 9^k$

Total work done =

$$\sum_{k=0}^{\log_3 n} (9^k) \cdot (\frac{n}{3^k})^2 \Rightarrow n^2 \cdot \sum_{k=0}^{\log_3 n} 1 \Rightarrow n^2 \cdot \log n \Rightarrow \theta(n^2 \log n)$$