Lecture 7 Signals and Systems (ELL205)

By Dr. Abhishek Dixit

Dept. of Electrical Engineering

IIT Delhi

Heinrich Hertz

(22 February 1857 – 1 January 1894; 36 years) was a German physicist and student of Kirchhoff and Helmholtz.

He provided the conclusive evidence of existence of EM waves.

The unit of frequency — <u>cycle per</u> <u>second</u> — was named the <u>"Hertz"</u> in his honor.

He was also a great scholar of languages and mastered languages like Arabic and Sanskrit.

He had one wife and 2 daughters (Mathilde Carmen Hertz, famous biologist).



Outline of the Lecture

System Properties

- 1. Memoryless
- 2. Causal
- 3. Invertible
- 4. Stable
- 5. Time invariant
- 6. Linear
- 7. Incrementally Linear

Outline of the Lecture

System Properties

- 1. Memoryless
- 2. Causal
- 3. Invertible
- 4. Stable
- 5. Time invariant
- 6. Linear
- 7. Incrementally Linear

Stable

A system is said to be stable if bounded inputs lead to bounded outputs.

Stable

A system is said to be stable if bounded inputs lead to bounded outputs.

Stability is BIBO stability

How many of them are stable systems?

A)	$y[n] = \sum_{k=-N}^{N} x[k]$
В)	$y(t) = \int_{-\infty}^{t} x(\tau) d\tau$
C)	$y(t) = \frac{dx(t)}{dt}$
D)	$y[n] = \sum_{k=-\infty}^{n} x[k]$

How many of them are stable systems?

A)	$y[n] = \sum_{k=-N}^{N} x[k]$
B)	$y(t) = \int_{-\infty}^{t} x(\tau) d\tau$
C)	$y(t) = \frac{dx(t)}{dt}$
D)	$y[n] = \sum_{k=-\infty}^{n} x[k]$

Stable

A system is said to be stable if bounded inputs lead to bounded outputs.

Stability is BIBO stability

Trick:

System is unstable if:

Infinite summation

Infinite integration

Differentiators

Outline of the Lecture

System Properties


- 1. Memoryless
- 2. Causal
- 3. Invertible
- 4. Stable
- 5. Time invariant
- 6. Linear
- 7. Incrementally Linear

Time-invariance

A system is said to be time-invariant if a time delay or time advance in the input produces an identical time shift in the output.

Time-invariance

A system is said to be time-invariant if a time delay or time advance in the input produces an identical time shift in the output.



$$y(t) = \sin(x(t))$$

$$y(t) = \sin(x(t))$$

Step 1: Change input x(t) to x(t-to)

$$\sin(x(t-to))$$

Equ. 1

$$y(t) = \sin(x(t))$$

Step 1: Change input x(t) to x(t-to)

$$\sin(x(t-to))$$

Equ. 1

Step 2: Change output y(t) to y(t - to)

$$y(t - to) = \sin(x(t - to))$$
 Equ. 2

$$y(t) = \sin(x(t))$$

Step 1: Change input x(t) to x(t-to)

$$\sin(x(t-to))$$
 Equ. 1

Step 2: Change output y(t) to y(t - to)

$$y(t - to) = \sin(x(t - to))$$
 Equ. 2

Step 3: TI if Equ. 1 and Equ. 2 matches

A)	$y(t) = \sin^2(x(t))$
В)	y[n] = nx[n]
C)	y(t) = x(2t)
D)	$y[n] = \sum_{k=-\infty}^{n} x[k]$

A)	$y(t) = \sin^2(x(t))$
В)	y[n] = nx[n]
C)	y(t) = x(2t)
D)	$y[n] = \sum_{k=-\infty}^{n} x[k]$

A)	$y(t) = x^2(t)$
В)	$y[n] = \text{Ev}\{x[n]\}$
C)	$y(t) = x^2(t/2)$
D)	$y(t) = t\sin(x(t))$

A)	$y(t) = x^2(t)$
В)	$y[n] = \text{Ev}\{x[n]\}$
C)	$y(t) = x^2(t/2)$
D)	$y(t) = t\sin(x(t))$

Outline of the Lecture

System Properties

- 1. Memoryless
- 2. Causal
- 3. Invertible
- 4. Stable
- 5. Time invariant
- 6. Linear
- 7. Incrementally Linear

A system is said to be linear iff it satisfies superposition (additivity & homogeneity)

A system is said to be linear iff it satisfies superposition (additivity & homogeneity)

Additivity

$$x_1(t) \longrightarrow y_1(t)$$

$$x_2(t) \longrightarrow y_2(t)$$

$$x_1(t) + x_2(t) \longrightarrow y_1(t) + y_2(t)$$

A system is said to be linear iff it satisfies superposition (additivity & homogeneity)

Homogeneity

$$x_1(t) \longrightarrow y_1(t)$$

$$ax_1(t) \longrightarrow ay_1(t)$$

A system is said to be linear iff it satisfies superposition (additivity & homogeneity)

Homogeneity

$$x_1(t) \longrightarrow y_1(t)$$

$$ax_1(t) \longrightarrow ay_1(t)$$

If
$$a = 0$$

$$0 \longrightarrow 0$$
 ZIZO

How many of them are linear systems?

A)	$y(t) = x^2(t)$
В)	$y[n] = \text{Ev}\{x[n]\}$
C)	y[n] = x[n] + 5
D)	$y(t) = \sin(x(t))$

How many of them are linear systems?

A)	$y(t) = x^2(t)$
В)	$y[n] = \text{Ev}\{x[n]\}$
C)	y[n] = x[n] + 5
D)	$y(t) = \sin(x(t))$