

# COL 352 Introduction to Automata and Theory of Computation

Nikhil Balaji

Bharti 420  
Indian Institute of Technology, Delhi  
[nbalaji@cse.iitd.ac.in](mailto:nbalaji@cse.iitd.ac.in)

February 1, 2023

Lecture 9: Non-regularity: Pumping Lemma

# Limitations of Finite Automata

$$L_{0,1} := \{0^n 1^n \mid n \geq 0\}$$

$$PAL := \{ww^R \mid w \in \Sigma^*\}$$

# Generalise?

These arguments seem to be example specific.. can they be generalised?

## Pumping Lemma

From the previous argument, we filter out a property of regular languages .

# Pumping Lemma

## Lemma

Let  $L$  be a regular language. Then

- ▶ there exists a constant  $n$  such that
- ▶ for every word  $w \in L$  such that  $|w| \geq n$

# Pumping Lemma

## Lemma

Let  $L$  be a regular language. Then

- ▶ there exists a constant  $n$  such that
- ▶ for every word  $w \in L$  such that  $|w| \geq n$
- ▶ we can break  $w$  in to three parts  $w = xyz$ , such that

# Pumping Lemma

## Lemma

Let  $L$  be a regular language. Then

- ▶ there exists a constant  $n$  such that
- ▶ for every word  $w \in L$  such that  $|w| \geq n$
- ▶ we can break  $w$  in to three parts  $w = xyz$ , such that
  - 1  $y \neq \epsilon$
  - 2  $|xy| \leq n$  and
  - 3 For all  $k \geq 0$ ,  $xy^kz \in L$ .

# Pumping Lemma

## Lemma

Let  $L$  be a regular language. Then

- ▶ there exists a constant  $n$  such that
- ▶ for every word  $w \in L$  such that  $|w| \geq n$
- ▶ we can break  $w$  in to three parts  $w = xyz$ , such that
  - 1  $y \neq \epsilon$
  - 2  $|xy| \leq n$  and
  - 3 For all  $k \geq 0$ ,  $xy^kz \in L$ .
- ▶ Lets write it using symbols

# Pumping Lemma

## Lemma

Let  $L$  be a regular language. Then

- ▶ there exists a constant  $n$  such that
- ▶ for every word  $w \in L$  such that  $|w| \geq n$
- ▶ we can break  $w$  in to three parts  $w = xyz$ , such that
  - 1  $y \neq \epsilon$
  - 2  $|xy| \leq n$  and
  - 3 For all  $k \geq 0$ ,  $xy^kz \in L$ .
- ▶ Lets write it using symbols

$$\exists n \forall w \in L (|w| \geq n \implies \exists xyz. (xyz = w \wedge y \neq \epsilon \wedge |xy| \leq n \wedge (\forall k \quad xy^kz \in L)))$$



# Understanding the lemma

It is called pumping lemma because we can "pump"  $y$  any number of times including 0.

# Understanding the lemma

It is called pumping lemma because we can "pump"  $y$  any number of times including 0.

- ▶ How many quantifier alternations?

# Understanding the lemma

It is called pumping lemma because we can "pump"  $y$  any number of times including 0.

- ▶ How many quantifier alternations?
- ▶ Exercise: Prove it holds for finite languages.

# Understanding the lemma

It is called pumping lemma because we can "pump"  $y$  any number of times including 0.

- ▶ How many quantifier alternations?
- ▶ Exercise: Prove it holds for finite languages.
- ▶ What happens if  $y = \epsilon$ ?

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.
- ▶ Consider word  $w = w_1 \dots w_m$ , where  $m \gg n$ .

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.
- ▶ Consider word  $w = w_1 \dots w_m$ , where  $m \gg n$ .
- ▶  $q_s = q_0 \xrightarrow{w_1} q_1 \xrightarrow{w_2} \dots \xrightarrow{w_m} q_m = q_f$  is a run of  $A$ .



# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.
- ▶ Consider word  $w = w_1 \dots w_m$ , where  $m \gg n$ .
- ▶  $q_s = q_0 \xrightarrow{w_1} q_1 \xrightarrow{w_2} \dots \xrightarrow{w_m} q_m = q_f$  is a run of  $A$ .
- ▶ By PHP, there exists  $i \neq j \in [m]$  s.t.  $q_i = q_j = q'$

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.
- ▶ Consider word  $w = w_1 \dots w_m$ , where  $m \gg n$ .
- ▶  $q_s = q_0 \xrightarrow{w_1} q_1 \xrightarrow{w_2} \dots \xrightarrow{w_m} q_m = q_f$  is a run of  $A$ .
- ▶ By PHP, there exists  $i \neq j \in [m]$  s.t.  $q_i = q_j = q'$
- ▶ Break  $w = xyz$

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.
- ▶ Consider word  $w = w_1 \dots w_m$ , where  $m \gg n$ .
- ▶  $q_s = q_0 \xrightarrow{w_1} q_1 \xrightarrow{w_2} \dots \xrightarrow{w_m} q_m = q_f$  is a run of  $A$ .
- ▶ By PHP, there exists  $i \neq j \in [m]$  s.t.  $q_i = q_j = q'$
- ▶ Break  $w = xyz$ 
  - ▶  $x = w_1 \dots w_i$
  - ▶  $y = w_{i+1} \dots w_j$
  - ▶  $z = w_{j+1} \dots w_m$
- ▶ By construction,  $y \neq \epsilon$ ,  $|xz| \leq n$

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.
- ▶ Consider word  $w = w_1 \dots w_m$ , where  $m \gg n$ .
- ▶  $q_s = q_0 \xrightarrow{w_1} q_1 \xrightarrow{w_2} \dots \xrightarrow{w_m} q_m = q_f$  is a run of  $A$ .
- ▶ By PHP, there exists  $i \neq j \in [m]$  s.t.  $q_i = q_j = q'$
- ▶ Break  $w = xyz$ 
  - ▶  $x = w_1 \dots w_i$
  - ▶  $y = w_{i+1} \dots w_j$
  - ▶  $z = w_{j+1} \dots w_m$
- ▶ By construction,  $y \neq \epsilon$ ,  $|xz| \leq n$
- ▶ For any  $k > 0$ ,  $xy^kz$  takes  $A$  to  $q_f$ , hence accepting.

# Proof of Pumping Lemma

- ▶ If  $L$  is regular, then there exists DFA  $A$  s.t.  $L = L(A)$ .
- ▶ Suppose  $A$  has  $n$  states.
- ▶ Consider word  $w = w_1 \dots w_m$ , where  $m \gg n$ .
- ▶  $q_s = q_0 \xrightarrow{w_1} q_1 \xrightarrow{w_2} \dots \xrightarrow{w_m} q_m = q_f$  is a run of  $A$ .
- ▶ By PHP, there exists  $i \neq j \in [m]$  s.t.  $q_i = q_j = q'$
- ▶ Break  $w = xyz$ 
  - ▶  $x = w_1 \dots w_i$
  - ▶  $y = w_{i+1} \dots w_j$
  - ▶  $z = w_{j+1} \dots w_m$
- ▶ By construction,  $y \neq \epsilon$ ,  $|xz| \leq n$
- ▶ For any  $k > 0$ ,  $xy^kz$  takes  $A$  to  $q_f$ , hence accepting.
- ▶ Therefore,  $\forall k > 0$ ,  $xy^kz \in L$

# Pumping Lemma


## Lemma

Let  $L$  be a regular language. Then

- ▶ there exists a constant  $n$  such that
- ▶ for every word  $w \in L$  such that  $|w| \geq n$
- ▶ we can break  $w$  in to three parts  $w = xyz$ , such that
  - 1  $y \neq \epsilon$
  - 2  $|xy| \leq n$  and
  - 3 For all  $k \geq 0$ ,  $xy^kz \in L$ .


# Example

Consider DFA  $A$  below.



## Example

Consider DFA  $A$  below.




The size of DFA is  $n = 3$ .



## Example

Consider DFA  $A$  below.




The size of DFA is  $n = 3$ .

Choose  $w = abaaba$ .  $|w| = 6 \geq 3$  and  $w$  is in  $L(A)$ .

## Example

Consider DFA  $A$  below.



The size of DFA is  $n = 3$ .

Choose  $w = abaaba$ .  $|w| = 6 \geq 3$  and  $w$  is in  $L(A)$ .

The run is  $q_0 \xrightarrow{a} q_1 \xrightarrow{b} q_0 \xrightarrow{a} q_1 \xrightarrow{a} q_2 \xrightarrow{b} q_1 \xrightarrow{a} q_2$

$q_1$  is repeated in the first 4 states of the run.

Choose  $x = a$ ,  $y = ba$ ,  $z = aba$

Therefore,  $a(ba)^k aba \in L(A)$

# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?

# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?
- ▶ What does the contrapositive of the lemma say?

# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?
- ▶ What does the contrapositive of the lemma say?
- ▶  $A \implies B$  iff  $\neg B \implies \neg A$ .

# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?
- ▶ What does the contrapositive of the lemma say?
- ▶  $A \implies B$  iff  $\neg B \implies \neg A$ .
- ▶  $L$  is regular  $\implies$  Pumping Conditions.  
 $\neg(\text{Pumping Conditions}) \implies L$  is not regular.

# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?
- ▶ What does the contrapositive of the lemma say?
- ▶  $A \implies B$  iff  $\neg B \implies \neg A$ .
- ▶  $L$  is regular  $\implies$  Pumping Conditions.  
 $\neg(\text{Pumping Conditions}) \implies L$  is not regular.

Recall the conditions of the lemma.

# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?
- ▶ What does the contrapositive of the lemma say?
- ▶  $A \implies B$  iff  $\neg B \implies \neg A$ .
- ▶  $L$  is regular  $\implies$  Pumping Conditions.  
 $\neg(\text{Pumping Conditions}) \implies L$  is not regular.

Recall the conditions of the lemma.

$$\exists n \forall w \in L (|w| \geq n \implies \exists xyz. (xyz = w \wedge y \neq \epsilon \wedge |xy| \leq n \wedge (\forall k \ xy^k z \in L)))$$



# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?
- ▶ What does the contrapositive of the lemma say?
- ▶  $A \implies B$  iff  $\neg B \implies \neg A$ .
- ▶  $L$  is regular  $\implies$  Pumping Conditions.  
 $\neg(\text{Pumping Conditions}) \implies L$  is not regular.

Recall the conditions of the lemma.

$$\exists n \forall w \in L (|w| \geq n \implies \exists xyz. (xyz = w \wedge y \neq \epsilon \wedge |xy| \leq n \wedge (\forall k \ xy^k z \in L)))$$

What is the contrapositive??

# Using the Pumping Lemma

- ▶ Pumping lemma gives a useful property of regular languages. How do we use it?
- ▶ What does the contrapositive of the lemma say?
- ▶  $A \implies B$  iff  $\neg B \implies \neg A$ .
- ▶  $L$  is regular  $\implies$  Pumping Conditions.  
 $\neg(\text{Pumping Conditions}) \implies L$  is not regular.

Recall the conditions of the lemma.

$$\exists n \forall w \in L (|w| \geq n \implies \\ \exists xyz. (xyz = w \wedge y \neq \epsilon \wedge |xy| \leq n \wedge (\forall k \ xy^k z \in L)))$$

What is the contrapositive??

$$\forall n \exists w \in L (|w| \geq n \wedge \\ \forall xyz. (xyz = w \wedge y \neq \epsilon \wedge |xy| \leq n \implies (\exists k \ xy^k z \notin L)))$$

# In words...

## Theorem

If

- ▶ for every  $n$
- ▶ there is a word  $w \in L$  where  $|w| \geq n$ .
- ▶ for every breakup of  $w$  into three words  $w = xyz$  such that
  - 1  $y \neq \epsilon$
  - 2  $|xy| \leq n$

there is a  $k \geq 0$  such that  $xy^kz \notin L$ , then

$L$  is not regular.

# In words...

## Theorem

If

- ▶ for every  $n$
- ▶ there is a word  $w \in L$  where  $|w| \geq n$ .
- ▶ for every breakup of  $w$  into three words  $w = xyz$  such that
  - 1  $y \neq \epsilon$
  - 2  $|xy| \leq n$

there is a  $k \geq 0$  such that  $xy^kz \notin L$ , then

$L$  is not regular.

How do we apply this lemma?

# How to use the lemma?

Consider language  $L$

- ▶ Let  $n$  be an arbitrary number (pumping length).
- ▶ (Cleverly) Find a representative string  $w$  of  $L$  of size  $\geq n$ .
- ▶ Try out all ways to break the string into  $xyz$  triplet satisfying that  $|y| > 0$  and  $|xy| \leq n$ . There will be finitely many cases to consider.
- ▶ For every triplet show that for some  $i$  the string  $xy^iz$  is not in  $L$ , and hence it yields contradiction with pumping lemma.

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$ . The breaks  $x$  and  $y$  are to be from within  $a^n$ .



# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$ . The breaks  $x$  and  $y$  are to be from within  $a^n$ .
- ▶ Let  $x = a^i, y = a^j$  where  $i + j \leq n$  and  $j \neq 0$ .

$$w = a^i a^j a^{n-i-j} b^n$$

# Applying Pumping Lemma: Example 1

Consider

$$L_{a,b} = \{a^n b^n \mid n \geq 0\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$ . The breaks  $x$  and  $y$  are to be from within  $a^n$ .
- ▶ Let  $x = a^i, y = a^j$  where  $i + j \leq n$  and  $j \neq 0$ .

$$w = a^i a^j a^{n-i-j} b^n$$

- ▶ Choose  $k = 0$  for each  $i, j$ . The corresponding word is

$$a^i (a^j)^0 a^{n-i-j} b^n = a^{n-j} b^n \notin L_{a,b}$$

## Example 2

Consider

$$L = \{w \in \{a, b\}^* \mid w \text{ has equal number of a's and b's}\}$$

## Example 2

Consider

$$L = \{w \in \{a, b\}^* \mid w \text{ has equal number of a's and b's}\}$$

- For each  $n$ , we need a word. Let  $w = a^n b^n$ .

## Example 2

Consider

$$L = \{w \in \{a, b\}^* \mid w \text{ has equal number of a's and b's}\}$$

- ▶ For each  $n$ , we need a word. Let  $w = a^n b^n$ .
- ▶ The first  $n$  characters of  $w$  are  $a^n$ . The breaks  $x$  and  $y$  are to be from within  $a^n$ .
- ▶ Let  $x = a^i, y = a^j$  where  $i + j \leq n$  and  $j \neq 0$ .

$$w = a^i a^j a^{n-i-j} b^n$$

- ▶ Choose  $k = 0$  for each  $i, j$ . The corresponding word is

$$a^i (a^j)^0 a^{n-i-j} b^n = a^{n-j} b^n \notin L_{a,b}$$

## Example 2 again

Consider

$$L = \{w \in \{a, b\}^* \mid w \text{ has equal number of a's and b's}\}$$

## Example 2 again

Consider

$$L = \{w \in \{a, b\}^* \mid w \text{ has equal number of a's and b's}\}$$

Exercise: What is  $L \cap a^*b^*$ ?