

COL 352 Introduction to Automata and Theory of Computation

Nikhil Balaji

Bharti 420
Indian Institute of Technology, Delhi
nbalaji@cse.iitd.ac.in

January 12, 2023

Lecture 5: Nondeterminism: Subset Construction

Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language.

Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.


We will work it out for an example.

Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

We will work it out for an example.




Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

We will work it out for an example.




Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

We will work it out for an example.




\emptyset $\{0\}$ $\{1\}$ $\{2\}$ $\{0,1\}$ $\{0,2\}$ $\{1,2\}$ $\{0,1,2\}$

Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

We will work it out for an example.




	\emptyset	$\{0\}$	$\{1\}$	$\{2\}$	$\{0,1\}$	$\{0,2\}$	$\{1,2\}$	$\{0,1,2\}$
a	\emptyset	$\{0,1\}$	$\{2\}$	\emptyset	$\{0,1,2\}$	$\{0,1\}$	$\{2\}$	$\{0,1,2\}$

Power of NFAs

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

We will work it out for an example.



	\emptyset	$\{0\}$	$\{1\}$	$\{2\}$	$\{0,1\}$	$\{0,2\}$	$\{1,2\}$	$\{0,1,2\}$
a	\emptyset	$\{0,1\}$	$\{2\}$	\emptyset	$\{0,1,2\}$	$\{0,1\}$	$\{2\}$	$\{0,1,2\}$
b	\emptyset	$\{0\}$	$\{2\}$	\emptyset	$\{0,2\}$	$\{0\}$	$\{2\}$	$\{0,2\}$

Subset construction

Lemma

Let A be an NFA. Then $L(A)$ is a regular language.

Subset construction

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

Subset construction

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

Proof.

Let $A = (Q, \Sigma, q_0, F, \delta)$. We will construct a DFA $A' = (Q', \Sigma, q'_0, F', \Delta)$ such that $L(A') = L(A)$.

Subset construction

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

Proof.

Let $A = (Q, \Sigma, q_0, F, \delta)$. We will construct a DFA $A' = (Q', \Sigma, q'_0, F', \Delta)$ such that $L(A') = L(A)$.

Subset construction

$$Q' = 2^Q$$

Subset construction

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

Proof.

Let $A = (Q, \Sigma, q_0, F, \delta)$. We will construct a DFA $A' = (Q', \Sigma, q'_0, F', \Delta)$ such that $L(A') = L(A)$.

Subset construction

$$Q' = 2^Q,$$

$$q'_0 = \{q_0\}$$

Subset construction

Lemma

Let A be an NFA. Then $L(A)$ is a regular language. That is, NFA and DFA accept the same set of languages.

Proof.

Let $A = (Q, \Sigma, q_0, F, \delta)$. We will construct a DFA $A' = (Q', \Sigma, q'_0, F', \Delta)$ such that $L(A') = L(A)$.

Subset construction

$$Q' = 2^Q,$$

$$q'_0 = \{q_0\},$$

$$F' = \{S \subseteq Q \mid S \cap F \neq \emptyset\}.$$

$$\text{For each } S \subseteq Q, \Delta(S, a) = \bigcup_{p \in S} \delta(p, a).$$



Correctness of subset construction

We will prove the following first.

Lemma

For any NFA A , there is an equivalent DFA A' such that $L(A') = L(A)$.

Correctness of subset construction

We will prove the following first.

Lemma

For any NFA A , there is an equivalent DFA A' such that $L(A') = L(A)$.

Definition

Let $A = (Q, \Sigma, \delta, q_0, F)$ be an NFA.

Let $\hat{\delta} : Q \times \Sigma^* \subseteq Q$ be defined as follows:

Correctness of subset construction

We will prove the following first.

Lemma

For any NFA A , there is an equivalent DFA A' such that $L(A') = L(A)$.

Definition

Let $A = (Q, \Sigma, \delta, q_0, F)$ be an NFA.

Let $\hat{\delta} : Q \times \Sigma^* \subseteq Q$ be defined as follows:

Let $q \in Q$

Correctness of subset construction

We will prove the following first.

Lemma

For any NFA A , there is an equivalent DFA A' such that $L(A') = L(A)$.

Definition

Let $A = (Q, \Sigma, \delta, q_0, F)$ be an NFA.

Let $\hat{\delta} : Q \times \Sigma^* \subseteq Q$ be defined as follows:

Let $q \in Q$

$$\hat{\delta}(q, \epsilon) := q$$

Correctness of subset construction

We will prove the following first.

Lemma

For any NFA A , there is an equivalent DFA A' such that $L(A') = L(A)$.

Definition

Let $A = (Q, \Sigma, \delta, q_0, F)$ be an NFA.

Let $\hat{\delta} : Q \times \Sigma^* \subseteq Q$ be defined as follows:

Let $q \in Q$

$$\hat{\delta}(q, \epsilon) := q$$

$$\hat{\delta}(q, xa) := \{p \mid \exists q' \in \hat{\delta}(q, x) \text{ s.t. } p \in \delta(q', a)\}$$

Correctness of subset construction

We will prove the following first.

Lemma

For any NFA A , there is an equivalent DFA A' such that $L(A') = L(A)$.

Definition

Let $A = (Q, \Sigma, \delta, q_0, F)$ be an NFA.

Let $\hat{\delta} : Q \times \Sigma^* \subseteq Q$ be defined as follows:

Let $q \in Q$

$$\hat{\delta}(q, \epsilon) := q$$

$$\hat{\delta}(q, xa) := \{p \mid \exists q' \in \hat{\delta}(q, x) \text{ s.t. } p \in \delta(q', a)\}$$

Proof of Correctness

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof.

Base case: Let $w = \epsilon$. $\hat{\delta}(\{q_0\}, \epsilon) = q_0 = \Delta(\{q_0\}, \epsilon) = \hat{\Delta}(\{q_0\}, \epsilon)$

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof.

Base case: Let $w = \epsilon$. $\hat{\delta}(\{q_0\}, \epsilon) = q_0 = \Delta(\{q_0\}, \epsilon) = \hat{\Delta}(\{q_0\}, \epsilon)$

Induction step: Let $w = x \cdot a$ where $w \in \Sigma^*$, $a \in \Sigma$

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof.

Base case: Let $w = \epsilon$. $\hat{\delta}(\{q_0\}, \epsilon) = q_0 = \Delta(\{q_0\}, \epsilon) = \hat{\Delta}(\{q_0\}, \epsilon)$

Induction step: Let $w = x \cdot a$ where $w \in \Sigma^*$, $a \in \Sigma$

- ▶ $\hat{\delta}(q_0, x) = \hat{\Delta}(\{q_0\}, x) = S$ (say) (By Induction Hypothesis)

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof.

Base case: Let $w = \epsilon$. $\hat{\delta}(\{q_0\}, \epsilon) = q_0 = \Delta(\{q_0\}, \epsilon) = \hat{\Delta}(\{q_0\}, \epsilon)$

Induction step: Let $w = x \cdot a$ where $w \in \Sigma^*$, $a \in \Sigma$

- ▶ $\hat{\delta}(q_0, x) = \hat{\Delta}(\{q_0\}, x) = S$ (say) (By Induction Hypothesis)
- ▶ $\hat{\delta}(q_0, xa) = \bigcup_{q \in S} \delta(q, a)$ (By definition of $\hat{\delta}$).

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof.

Base case: Let $w = \epsilon$. $\hat{\delta}(\{q_0\}, \epsilon) = q_0 = \Delta(\{q_0\}, \epsilon) = \hat{\Delta}(\{q_0\}, \epsilon)$

Induction step: Let $w = x \cdot a$ where $w \in \Sigma^*$, $a \in \Sigma$

- ▶ $\hat{\delta}(q_0, x) = \hat{\Delta}(\{q_0\}, x) = S$ (say) (By Induction Hypothesis)
- ▶ $\hat{\delta}(q_0, xa) = \bigcup_{q \in S} \delta(q, a)$ (By definition of $\hat{\delta}$).
- ▶ $\hat{\Delta}(\{q_0\}, xa) = \Delta(S, a) = \bigcup_{q \in S} \delta(q, a)$ (By definition of $\hat{\Delta}$).

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof.

Base case: Let $w = \epsilon$. $\hat{\delta}(\{q_0\}, \epsilon) = q_0 = \Delta(\{q_0\}, \epsilon) = \hat{\Delta}(\{q_0\}, \epsilon)$

Induction step: Let $w = x \cdot a$ where $w \in \Sigma^*$, $a \in \Sigma$

- ▶ $\hat{\delta}(q_0, x) = \hat{\Delta}(\{q_0\}, x) = S$ (say) (By Induction Hypothesis)
- ▶ $\hat{\delta}(q_0, xa) = \bigcup_{q \in S} \delta(q, a)$ (By definition of $\hat{\delta}$).
- ▶ $\hat{\Delta}(\{q_0\}, xa) = \Delta(S, a) = \bigcup_{q \in S} \delta(q, a)$ (By definition of $\hat{\Delta}$).
- ▶ Therefore, $\hat{\delta}(q_0, xa) = \hat{\Delta}(\{q_0\}, xa)$.

Proof of correctness

Lemma

For all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$.

Proof.

Base case: Let $w = \epsilon$. $\hat{\delta}(\{q_0\}, \epsilon) = q_0 = \Delta(\{q_0\}, \epsilon) = \hat{\Delta}(\{q_0\}, \epsilon)$

Induction step: Let $w = x \cdot a$ where $w \in \Sigma^*$, $a \in \Sigma$

- ▶ $\hat{\delta}(q_0, x) = \hat{\Delta}(\{q_0\}, x) = S$ (say) (By Induction Hypothesis)
- ▶ $\hat{\delta}(q_0, xa) = \bigcup_{q \in S} \delta(q, a)$ (By definition of $\hat{\delta}$).
- ▶ $\hat{\Delta}(\{q_0\}, xa) = \Delta(S, a) = \bigcup_{q \in S} \delta(q, a)$ (By definition of $\hat{\Delta}$).
- ▶ Therefore, $\hat{\delta}(q_0, xa) = \hat{\Delta}(\{q_0\}, xa)$.



From Lemma, we have for all $w \in \Sigma^*$, $\hat{\delta}(q_0, w) = \hat{\Delta}(\{q_0\}, w)$. Hence, for all w , $\hat{\delta}(q_0, w) \cap F \neq \emptyset$ iff $\hat{\Delta}(\{q_0\}, w) \in F'$.

Complexity of Subset Construction

- ▶ If NFA has n states, then DFA has 2^n states.

Complexity of Subset Construction

- ▶ If NFA has n states, then DFA has 2^n states.
- ▶ But not all these states need to be reachable!

Complexity of Subset Construction

- ▶ If NFA has n states, then DFA has 2^n states.
- ▶ But not all these states need to be reachable!

Two questions

- ▶ Does this blowup really occur when only considering reachable states?
- ▶ On examples where it does not occur can we have a subset construction that is efficient?

Worst case blowup of Subset Construction

- ▶ Is one NFA enough? A family of examples such that for each n there is an example NFA parametrized by n .

Worst case blowup of Subset Construction

- ▶ Is one NFA enough? A family of examples such that for each n there is an example NFA parametrized by n .
- ▶ For each example in family show that if a smaller DFA than 2^n exists, there is a contradiction.

Worst case blowup of Subset Construction

- ▶ Is one NFA enough? A family of examples such that for each n there is an example NFA parametrized by n .
- ▶ For each example in family show that if a smaller DFA than 2^n exists, there is a contradiction.

For each $n \in \mathbb{N}$, let

$$L_n = \{w \in \{0, 1\}^* \mid n\text{-th symbol from the end is a } 1\}$$

Worst case blowup of Subset Construction

- ▶ Is one NFA enough? A family of examples such that for each n there is an example NFA parametrized by n .
- ▶ For each example in family show that if a smaller DFA than 2^n exists, there is a contradiction.

For each $n \in \mathbb{N}$, let

$$L_n = \{w \in \{0, 1\}^* \mid n\text{-th symbol from the end is a } 1\}$$

- ▶ Exercise: give an NFA for L_n . How many states does it have?
- ▶ Can L_n be accepted by a DFA with less than 2^n states?

Worst case blowup of Subset Construction

- ▶ Is one NFA enough? A family of examples such that for each n there is an example NFA parametrized by n .
- ▶ For each example in family show that if a smaller DFA than 2^n exists, there is a contradiction.

For each $n \in \mathbb{N}$, let

$$L_n = \{w \in \{0, 1\}^* \mid n\text{-th symbol from the end is a } 1\}$$

- ▶ Exercise: give an NFA for L_n which accepts L_n . How many states does it have?
- ▶ Can L_n be accepted by a DFA with less than 2^n states?

$$L = \{x \in \{a\}^* \mid |x| \text{ is divisible by } 3 \text{ or } 5\}$$

Worst case blowup of Subset Construction

- ▶ Is one NFA enough? A family of examples such that for each n there is an example NFA parametrized by n .
- ▶ For each example in family show that if a smaller DFA than 2^n exists, there is a contradiction.

For each $n \in \mathbb{N}$, let

$$L_n = \{w \in \{0, 1\}^* \mid n\text{-th symbol from the end is a } 1\}$$

- ▶ Exercise: give an NFA for A_n which accepts L_n . How many states does it have?
- ▶ Can L_n be accepted by a DFA with less than 2^n states?

; divisible by 3 or 5}



Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

- ▶ Suppose not. For some $n \in \mathbb{N}$ let L_n be accepted by DFA A with less than 2^n states.

Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

- ▶ Suppose not. For some $n \in \mathbb{N}$ let L_n be accepted by DFA A with less than 2^n states.
- ▶ Then, there exist $a_1 \dots a_n$ and $b_1 \dots b_n$ that must both end in the same state q .

Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

- ▶ Suppose not. For some $n \in \mathbb{N}$ let L_n be accepted by DFA A with less than 2^n states.
- ▶ Then, there exist $a_1 \dots a_n$ and $b_1 \dots b_n$ that must both end in the same state q .
- ▶ Suppose a_i and b_i are different and wlog let $a_i = 1, b_i = 0$.

Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

- ▶ Suppose not. For some $n \in \mathbb{N}$ let L_n be accepted by DFA A with less than 2^n states.
- ▶ Then, there exist $a_1 \dots a_n$ and $b_1 \dots b_n$ that must both end in the same state q .
- ▶ Suppose a_i and b_i are different and wlog let $a_i = 1, b_i = 0$.
- ▶ Consider words

$$w = a_1 \dots a_i \dots a_n 0^{i-1} \in L_n$$

$$w' = b_1 \dots b_i \dots b_n 0^{i-1} \notin L_n$$

Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

- ▶ Suppose not. For some $n \in \mathbb{N}$ let L_n be accepted by DFA A with less than 2^n states.
- ▶ Then, there exist $a_1 \dots a_n$ and $b_1 \dots b_n$ that must both end in the same state q .
- ▶ Suppose a_i and b_i are different and wlog let $a_i = 1, b_i = 0$.
- ▶ Consider words

$$w = a_1 \dots a_i \dots a_n 0^{i-1} \in L_n$$

$$w' = b_1 \dots b_i \dots b_n 0^{i-1} \notin L_n$$

- ▶ But A is a DFA, so starting from q on reading 0^{i-1} we reach the same state.

Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

- ▶ Suppose not. For some $n \in \mathbb{N}$ let L_n be accepted by DFA A with less than 2^n states.
- ▶ Then, there exist $a_1 \dots a_n$ and $b_1 \dots b_n$ that must both end in the same state q .
- ▶ Suppose a_i and b_i are different and wlog let $a_i = 1, b_i = 0$.
- ▶ Consider words

$$w = a_1 \dots a_i \dots a_n 0^{i-1} \in L_n$$

$$w' = b_1 \dots b_i \dots b_n 0^{i-1} \notin L_n$$

- ▶ But A is a DFA, so starting from q on reading 0^{i-1} we reach the same state. Hence A will either accept both w and w' or reject both.

Exponential Blowup of Subset Construction

Theorem

For all $n \in \mathbb{N}$, every DFA accepting L_n has at least 2^n states.

Proof.

- ▶ Suppose not. For some $n \in \mathbb{N}$ let L_n be accepted by DFA A with less than 2^n states.
- ▶ Then, there exist $a_1 \dots a_n$ and $b_1 \dots b_n$ that must both end in the same state q .
- ▶ Suppose a_i and b_i are different and wlog let $a_i = 1, b_i = 0$.
- ▶ Consider words

$$w = a_1 \dots a_i \dots a_n 0^{i-1} \in L_n$$

$$w' = b_1 \dots b_i \dots b_n 0^{i-1} \notin L_n$$

- ▶ But A is a DFA, so starting from q on reading 0^{i-1} we reach the same state. Hence A will either accept both w and w' or reject both. Contradiction!