

Lecture 6

Signals and Systems (ELL205)

By Dr. Abhishek Dixit
Dept. of Electrical Engineering
IIT Delhi

Leonhard Euler

(15 April 1707 – 18 September 1783) was a Swiss mathematician.

His collected works fill 60 to 80 quarto volumes, more than anybody in the field.

In 1766, he got completely blind, and then he quoted “now I have a fewer distractions.”

After that he produced almost one mathematical paper every week.

He had two wives. The second of whom he took at the age 69. He had 13 children.




Outline of the Lecture

System Properties

1. Memoryless
2. Causal
3. Invertible
4. Stable
5. Time invariant
6. Linear
7. Incrementally Linear

System



System



Outline of the Lecture

System Properties

1. Memoryless
2. Causal
3. Invertible
4. Stable
5. Time invariant
6. Linear
7. Incrementally Linear

Memoryless

A system is said to be memoryless if $y(t)$ depends only on input at time at time t , i.e.,

$$y(t) = f(x(t))$$

How many of them are memoryless systems?

A)	$y(t) = x(t^2)$
B)	$y(t) = tx(t+1)$
C)	$y[n] = nx[n] + y[n-1]$
D)	$y(t) = \frac{1}{C} \int_{-\infty}^t x(\tau) d\tau$

NONE!!

Outline of the Lecture

System Properties

1. Memoryless
2. Causal
3. Invertible
4. Stable
5. Time invariant
6. Linear
7. Incrementally Linear

Causal

A system is said to be causal if the output depends only on the current input and on the inputs in the past.

Causal

A system is said to be causal if the output depends only on the current input and on the inputs in the past.

Output does not depend on inputs in future (non-anticipatory system)

Causal

A system is said to be causal if the output depends only on the current input and on the inputs in the past.

Output does not depend on inputs in future (non-anticipatory system)

Formal:

$$x_1(t) \equiv x_2(t) \quad \forall \quad t < t_0$$

$$y_1(t) \equiv y_2(t) \quad \forall \quad t < t_0$$

How many of them are causal systems?

A)	$y(t) = \cos(t + 1)x(t)$
B)	$y(t) = \cos(t)x(t + 1)$
C)	$y[n] = x[-n]$
D)	$y[n] = \sum_{k=-M}^M x[n - k]$

How many of them are causal systems?

A)	$y(t) = \cos(t + 1)x(t)$
B)	$y(t) = \cos(t)x(t + 1)$
C)	$y[n] = x[-n]$
D)	$y[n] = \sum_{k=-M}^M x[n - k]$

Outline of the Lecture

System Properties


1. Memoryless
2. Causal
3. Invertible
4. Stable
5. Time invariant
6. Linear
7. Incrementally Linear

Invertible

A system is said to be invertible if distinct inputs lead to distinct outputs.

Invertible

A system is said to be invertible if distinct inputs lead to distinct outputs.



How many of them are invertible systems?


A)	$y(t) = x^2(t)$
B)	$y(t) = 0$
C)	$y[n] = x[n]$
D)	$y[n] = \sum_{k=-\infty}^n x[k]$

How many of them are invertible systems?

A)	$y(t) = x^2(t)$
B)	$y(t) = 0$
C)	$y[n] = x[n]$
D)	$y[n] = \sum_{k=-\infty}^n x[k]$

Invertibility: Discrete Time Systems


1. $y[n] = 0$



System is **not** invertible.

Invertibility: Discrete Time Systems


2. $y[n] = x[n]$



System is invertible.

Invertibility: Discrete Time Systems


3. $y[n] = \sum_{k=-\infty}^n x[k]$ *Hint: $y[n] - y[n-1] = x[n]$*



System is invertible.

Invertibility: Discrete Time Systems

3. $y[n] = x[n] - x[n - 1]$



System is **not** invertible.

Invertibility: Discrete Time Systems

3. $y[n] = x[n] - x[n - 1]$

$x[n - 1]$ is **unknown**.

Hence, system is **not** invertible.

But, if $x[n - 1] = 0$, system becomes invertible.

