

# Vector Analysis - II

PYL101: Electromagnetics & Quantum Mechanics  
Semester I, 2020-2021

Prof. Rohit Narula<sup>1</sup>


<sup>1</sup>Department of Physics  
The Indian Institute of Technology, Delhi



# Copyright Notice

- ▶ My lectures and course materials, including presentations, tests, exams, outlines, and similar materials, are protected by copyright and for the sole use of students registered for PYL101: Electromagnetics & Quantum Mechanics in Sem-I 2020-2021 only. I am the exclusive owner of copyright in those materials I create. I encourage you to take notes and make copies of course materials for your own educational use. However, you may not, nor may you knowingly allow others to reproduce or distribute lecture notes, audio, video, screenshots and other course materials publicly or to internal university repositories such as the BSW without my express written consent. This includes providing materials to commercial course material suppliers such as CourseHero and other similar services. Students who publicly distribute or display or help others publicly distribute or display copies or modified copies of my course materials may be in violation of applicable law.

# Line Integrals



- ▶ A **line integral** is an expression of the form

$$\int_a^b \underbrace{\mathbf{v} \cdot}_{\text{dot product}} d\mathbf{l}$$


where  $\mathbf{v}$  is a vector function, and  $d\mathbf{l}$  is the infinitesimal displacement vector.

- ▶ The integral **must be** carried out **along a prescribed path**  $\mathcal{P}$  from point  $\mathbf{a}$  to point  $\mathbf{b}$ , and we should instead write

$$\int_a^b \int_{\mathcal{P}} \mathbf{v} \cdot d\mathbf{l}$$

explicitly mentioning the path  $\mathcal{P}$ .

# Line Integrals




- If the path  $\mathcal{P}$  is **closed loop** (i.e.,  $\mathbf{a} = \mathbf{b}$ ) we put a **circle** around the integral sign as,

$$\oint_{\mathcal{P}} \mathbf{v} \cdot d\mathbf{l} \neq 0$$

where it's important to note that, in general, closed line integrals don't have to be zero, *in general!*

## A Line Integral Example




- **Problem:** Calculate the line integral for the vector function

$$\mathbf{v} = y^2 \hat{\mathbf{x}} + 2x(y+1) \hat{\mathbf{y}}$$

from the point  $a = (1, 1, 0)$  to  $b = (2, 2, 0)$ , along the paths (1) and (2) in the above figure.


## A Line Integral Example: Along path (1)



- ▶ Along the **horizontal segment of (1)** (only  $x$  varies)  $dy = dz = 0$ , so

$$d\mathbf{l} = dx\hat{\mathbf{x}}, y = 1, \mathbf{v} \cdot d\mathbf{l} = y^2 dx, \int \mathbf{v} \cdot d\mathbf{l} = (1)^2 \int_1^2 dx = \mathbf{1}$$

## A Line Integral Example: Along path (l)



- For the **vertical segment of (l)** (only  $y$  varies)  $dx = dz = 0$ , so


$$d\mathbf{l} = dy\hat{y}, x = 2, \mathbf{v} \cdot d\mathbf{l} = 2x(y+1)dy, \int \mathbf{v} \cdot d\mathbf{l} = 4 \int_1^2 (y+1)dy = 10$$

- Thus, summing the horizontal and vertical parts of (l) we get,

$$\int_{(l)} \mathbf{v} \cdot d\mathbf{l} = 1 + 10 = 11$$



## A Line Integral Example: Along path (2)



- For path (2),  $x = y \Rightarrow dx = dy$ <sup>1</sup> and,  $dz = 0$

$$d\mathbf{l} = dx\hat{\mathbf{x}} + dy\hat{\mathbf{y}}, \mathbf{v} \cdot d\mathbf{l} = x^2 dx + 2x(x+1)dx = (3x^2 + 2x)dx,$$


, and finally,

$$\int_{(2)} \mathbf{v} \cdot d\mathbf{l} = 10$$

---

<sup>1</sup>This step is crucial for a *general* path where variables are interdependent.


## A Line Integral Example: $a \rightarrow (1) \rightarrow (2) \rightarrow a$



- For the *circuitous* path  $a \rightarrow (1) \rightarrow (2) \rightarrow a$ , we simply **sum** the contributions of paths (1) and (2) and get,

$$\oint \mathbf{v} \cdot d\mathbf{l} = 11 - 10 = \mathbf{1}$$

# Surface/Double and Flux Integrals



- ▶ A **surface/double integral** is an expression of the form

$$\int_{\mathcal{S}} f \, da$$

where  $f$  is a scalar function,  $da$  is an infinitesimal patch of area over a prescribed surface  $\mathcal{S}$ .


- ▶ However, the **flux**<sup>2</sup> of a vector function  $\mathbf{v}$  through  $\mathcal{S}$  is defined as,

$$\int_{\mathcal{S}} \mathbf{v} \cdot \underbrace{\quad}_{\text{dot product}} da$$

---

<sup>2</sup>in analogy with liquid flow, i.e., if  $\mathbf{v}$  describes the flow a mass of liquid per unit area per unit time.

# Flux Integrals




- ▶ Since there are two opposite directions for the surface normal, we *choose* the direction that points **radially outward** with the origin as a reference point.
- ▶ If the surface is **closed**<sup>3</sup> we write,

$$\oint_{\mathcal{S}} \mathbf{v} \cdot d\mathbf{a} \neq 0$$

---

<sup>3</sup>A closed surface in 3-d is exemplified by a *balloon*.

## A Flux Example




- **Problem:** Calculate the flux

$$\mathbf{v} = 2xz\hat{\mathbf{x}} + (x+2)\hat{\mathbf{y}} + y(z^2 - 3)\hat{\mathbf{z}}$$

over five sides (*excluding the bottom*).

## A Flux Example: Surface 1




(i)  $x = 2$ ,  $d\mathbf{a} = dy dz \hat{\mathbf{x}}$ ,  $\mathbf{v} \cdot d\mathbf{a} = 2xz dy dz = 4z dy dz$ , so

$$\int \mathbf{v} \cdot d\mathbf{a} = 4 \int_0^2 dy \int_0^2 z dz = 16.$$

- Note, that for surface (i) variables  $x$  and  $y$  are independent of each other<sup>4</sup>, and the integrals for each can be carried out independently.

<sup>4</sup>Why?


## A Flux Example: Surface 2



(ii)  $x = 0$ ,  $d\mathbf{a} = -dy dz \hat{\mathbf{x}}$ ,  $\mathbf{v} \cdot d\mathbf{a} = -2xz dy dz = 0$ , so

$$\int \mathbf{v} \cdot d\mathbf{a} = 0.$$

## A Flux Example: Surface 3




(iii)  $y = 2$ ,  $d\mathbf{a} = dx \, dz \, \hat{\mathbf{y}}$ ,  $\mathbf{v} \cdot d\mathbf{a} = (x + 2) \, dx \, dz$ , so

$$\int \mathbf{v} \cdot d\mathbf{a} = \int_0^2 (x + 2) \, dx \int_0^2 dz = 12.$$




## A Flux Example: Surface 4



(iv)  $y = 0$ ,  $d\mathbf{a} = -dx dz \hat{\mathbf{y}}$ ,  $\mathbf{v} \cdot d\mathbf{a} = -(x + 2) dx dz$ , so

$$\int \mathbf{v} \cdot d\mathbf{a} = - \int_0^2 (x + 2) dx \int_0^2 dz = -12.$$


## A Flux Example: Surface 5



(v)  $z = 2$ ,  $d\mathbf{a} = dx dy \hat{\mathbf{z}}$ ,  $\mathbf{v} \cdot d\mathbf{a} = y(z^2 - 3) dx dy = y dx dy$ , so

$$\int \mathbf{v} \cdot d\mathbf{a} = \int_0^2 dx \int_0^2 y dy = 4.$$

## A Flux Example: Total




The *total* flux is

$$\int_{\text{surface}} \mathbf{v} \cdot d\mathbf{a} = 16 + 0 + 12 - 12 + 4 = 20.$$

- Note that the surface  $\mathcal{S}$  over which the flux is calculated need not be closed.

## Evaluating a Surface Integral over a triangular $\mathcal{S}$



- **Problem:** Evaluate


$$\iint_{\mathcal{S}} xy \, dx \, dy$$

over the triangular region with  $\perp$  sides of unity, as above.

- Our first **strategy** is to:

1. Calculate  $T(x, y = \text{const.}) \times$  the area of each infinitesimally short **horizontal strip** of height  $dy$  [INNER SUM:  $x : 0 \rightarrow (1 - y)$ ], and then,
2. Sum these horizontal strips up one by one [OUTER SUM:  $y : 0 \rightarrow 1$ ].

## Evaluating the Surface Integral over a triangular $\mathcal{S}$



- Thus, keeping **y constant**, let's focus now on the [INNER SUM:  $x: 0 \rightarrow (1 - y)$ ], i.e.,

$$H(y) = \int_{x=0}^{1-y} \underbrace{xy}_{y=\text{const.}} dx = \frac{y(1-y)^2}{2}$$

where the  $(1 - y)$  limit accounts for the fact that the length of these strips varies.


- Finally, we perform the [OUTER SUM:  $y: 0 \rightarrow 1$ ] and

$$\int_{y=0}^1 H(y) dy = \int_{y=0}^1 \frac{y(1-y)^2}{2} dy = \frac{1}{24}$$

- HW: Find  $\int_{\mathcal{S}} xy \, dx dy$  by instead summing **vertical** strips.

# Evaluating the Surface Integral for a Horizontally-Simple Region

optional




- ▶ A **horizontally-simple region** is a region where every horizontal line drawn inside it shares the same left-function  $g_1(y)$  and right-function  $g_2(y)$ .
- ▶ It's evaluated as

$$\iint_{\mathcal{S}} f(x, y) dA = \int_{y=c}^{y=d} \left[ \int_{x=g_1(y)}^{x=g_2(y)} \underbrace{f(x, y)}_{y=\text{const}} dx \right] dy$$

- ▶ You should be able to *guess* the math for surface integrals over **vertically-simple regions**.

# Evaluating the **Flux** for a surface described by $z = g(x, y)$

optional



- When the surface  $\mathcal{S}$  can be described by the equation

$$z = g(x, y)$$

the **flux** of the vector function  $F(x, y, z)$  through  $\mathcal{S}$  is given by


$$\int_{\mathcal{S}} \mathbf{F} \cdot d\mathcal{S} = \iint_A \mathbf{F} \cdot \left( -\frac{\partial g}{\partial x} \hat{\mathbf{x}} - \frac{\partial g}{\partial y} \hat{\mathbf{y}} + 1 \hat{\mathbf{z}} \right) dx dy$$

where the region  $A$  is the **projection** of  $\mathcal{S}$  on the  $(x, y)$ -plane<sup>5</sup>.

<sup>5</sup>Note the subtlety in the above formula which is that in the integrand of the RHS, after calculating the dot product you must replace any instance of  $z$  by  $g(x, y)$ .

# Evaluating the **Flux** for a surface described by $z = g(x, y)$

optional



- **Problem:** Calculate the flux of  $\mathbf{F} = x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}$  over the surface given by the planar region  $\mathcal{S} : x + y + z = 1$  above. The projection of  $\mathcal{S}$  along the  $(x, y)$ -plane is the shaded region  $A$ .
- Therefore<sup>6</sup>,  $z = g(x, y) = 1 - x - y$ , and  $\frac{\partial g}{\partial x} = -1$ ,  $\frac{\partial g}{\partial y} = -1$

$$\begin{aligned}\int_{\mathcal{S}} \mathbf{F} \cdot d\mathcal{S} &= \iint_A (x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}) \cdot (\hat{\mathbf{x}} + \hat{\mathbf{y}} + \hat{\mathbf{z}}) dA \\ &= \iint_A (x + y + z) dx dy \quad (x + y + z = 1 : \mathcal{S}) \\ &= \frac{1}{2}\end{aligned}$$

---

<sup>6</sup>In the integrand above,  $z \neq 0$  despite the projected surface  $A$  lying on the  $(x, y)$ -plane. Instead replace any mention of the variable  $z = g(x, y) = 1 - x - y$ .



## Calculating Fluxes Numerically

- ▶ Let the vector function  $\mathbf{v}(x, y) = xy^2 \hat{\mathbf{z}}$ , and  $\mathcal{S}$  be a square region of sides unity, with its south-west corner located at the origin.
- ▶ Divide the square region into:
  - ▶ 4 subsquares of length  $dx_i = 0.25$  and breath  $dy_i = 0.25$
  - ▶ 16 subsquares of length  $dx_i = 0.0625$  and breath  $dy_i = 0.0625$
  - ▶ 64 subsquares of length  $dx_i = 0.015625$  and breath  $dy_i = 0.015625 \dots$and calculate the value  $\mathbf{v}(x_i, y_i)$  at the *center*  $(x_i, y_i)$  of each of the subsquares  $i$ .
- ▶ The area element  $d\mathbf{a} = dx_i dy_i \hat{\mathbf{z}}$ .
- ▶ The sum<sup>7</sup>  $\sum_i \mathbf{v}(x_i, y_i) dx_i dy_i$  is the required answer!
- ▶ As you **increase** the number of subsquares, the value obtained from the sum above **converges** to the **exact** value:

$$\int_{\mathcal{S}} \mathbf{v} \cdot d\mathbf{a} = \frac{1}{6}$$

---

<sup>7</sup>A practical way to do this is to write a small computer program.

# Volume Integrals

- ▶ Given a **scalar** function  $T$ , volume integrals are expressed as,

$$\int_{\mathcal{V}} T d\tau$$

and  $d\tau$  is an infinitesimal volume element.

- ▶ For a **vector** function  $\mathbf{v}$ , on the other hand, the associated volume integral is,

$$\begin{aligned}\int \mathbf{v} d\tau &= \int (v_x \hat{\mathbf{x}} + v_y \hat{\mathbf{y}} + v_z \hat{\mathbf{z}}) d\tau \\ &= \hat{\mathbf{x}} \int v_x d\tau + \hat{\mathbf{y}} \int v_y d\tau + \hat{\mathbf{z}} \int v_z d\tau\end{aligned}$$


- ▶ In **Cartesian coordinates**,<sup>8</sup>

$$d\tau = dx dy dz$$

---


<sup>8</sup>Spherical and cylindrical coordinates are other candidates.

## A Volume Integral Example



- ▶ **Problem:** Calculate the volume integral of  $T = xyz^2$ .
- ▶ It's easy to see from the prismatic volume that  $z$  does not depend on either  $x$  or  $y$ , and ranges from 0 to 3.
- ▶ This means that the integral over  $z$  can be **factored out**.


## A Volume Integral Example



$$\begin{aligned}\int xyz^2 d\tau &= \int_0^3 z^2 dz \underbrace{\iint xy \, dx dy}_{\text{done earlier!}} \\ &= 9 \int_{y=0}^1 y \left[ \int_{x=0}^{1-y} x \, dx \right] dy \\ &= \frac{9}{2} \int_0^1 y(1-y)^2 dy = \frac{3}{8}\end{aligned}$$

# Volume Integrals Over More General Regions

optional



- ▶ Let  $R$  be a **solid region**<sup>9</sup> bounded below and above by the functions  $g_1(x, y)$  and  $g_2(x, y)$ , respectively such that

$$g_1(x, y) \leq z \leq g_2(x, y)$$

- ▶ The region  $D$  is the **projection** of  $R$  onto the  $xy$ -plane. The triple integral is then given by


$$\int_R f(x, y, z) dV = \iint_D \left[ \int_{z=g_1(x, y)}^{g_2(x, y)} \underbrace{f(x, y, z) dz}_{x, y = \text{const}} \right] dA = \iint_D h(x, y) dA$$

which is a double-integral over the region  $D$  in the  $(x, y)$ -plane.

<sup>9</sup>Such a solid region is known as a "z-simple" solid.

# Volume Integrals Over General Regions

optional



- **Problem:** Evaluate

$$\int_R (x+2y) dV$$

where  $R$  is the **tetrahedron** bounded by the planes  $x=0, y=0, z=0$  and  $x+y+z=2$ .


- We can rewrite the equation of the plane  $x+y+z=2$  as  $z=2-x-y$ . Note that  $0 \leq z \leq 2-x-y$ . Hence, we have<sup>10</sup>

$$\iint_D \left[ \int_0^{2-x-y} \underbrace{(x+2y)}_{x,y=\text{const}} dz \right] dA = \iint_D (x+2y)(2-x-y) dA = \mathbf{2}$$

---

<sup>10</sup>The required answer was **incorrectly** given as  $\frac{2}{3}$  in the video lecture.

# The Fundamental Theorem for Gradients



- ▶ Given a **scalar** function  $T(x, y, z)$ , by changing  $x$ ,  $y$  and  $z$  *infinitesimally*, i.e., by  $dx$ ,  $dy$  and  $dz$ , the variation in  $T$  is,

$$dT = (\nabla T) \cdot d\mathbf{l}$$

- ▶ If we keep advancing from  $a$  to  $b$  along units of  $d\mathbf{l}_i$  we can **accumulate** the total change in the scalar function  $T$ ,


$$\int_a^b (\nabla T) \cdot d\mathbf{l} = T(b) - T(a)$$

also known as the **fundamental theorem for gradients**<sup>11</sup>.

---

<sup>11</sup>There is also a direct proof using the integral in the LHS above.

# The Fundamental Theorem for Gradients



- ▶ Remarkably, the RHS, i.e.,  $T(b) - T(a)$  makes no reference to the actual **path** taken.
- ▶ It implies that,  $\int_a^b (\nabla T) \cdot d\mathbf{l}$  is **independent** of the path taken from  $a$  to  $b$ .<sup>12</sup>
- ▶ It also implies that,

$$\oint (\nabla T) \cdot d\mathbf{l} = 0$$

---

<sup>12</sup>In practice, even though the integral is independent of the path, we **must** pick a specific (if *convenient*) route in order to evaluate it explicitly.




# The Fundamental Theorem for Divergences

- ▶ The fundamental theorem for divergences states that:

$$\int_{\mathcal{V}} (\nabla \cdot \mathbf{v}) d\tau = \oint_{\mathcal{S}} \mathbf{v} \cdot d\mathbf{a}$$


- ▶ It is alternatively known as, **Gauss' theorem**, **Green's theorem**, or simply the **divergence theorem**.
- ▶ **Physical Analogy:** Imagine we're interested in measuring the flux (RHS) of an *incompressible* fluid through a **closed** area  $\mathcal{S}$ .
- ▶ The divergence theorem states that instead of measuring the flux directly, we could've *equivalently*, sum up all the (liquid) **sources** inside the volume  $\mathcal{V}$  enclosed by the surface  $\mathcal{S}$ .

# The Fundamental Theorem for Divergences



- Q: Verify the divergence theorem for  $\mathbf{v} = 2x\hat{\mathbf{x}} + y^2\hat{\mathbf{y}} + z^2\hat{\mathbf{z}}$  for the sphere of radius unity. [Ans: LHS = RHS =  $\frac{8\pi}{3}$ ]

# The Fundamental Theorem for Curls



- ▶ Also known as **Stoke's theorem** states that,

$$\int_{\mathcal{S}} (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = \oint_{\mathcal{P}} \mathbf{v} \cdot d\mathbf{l}$$


which says that the integral of a curl (of a vector function) over a surface  $\mathcal{S}$  is equal to the value of the function integrated over the boundary  $\mathcal{P}$ ) *enclosing* that surface.

- ▶ Sticking with the liquid analogy, since the curl measures the *twist* of  $\mathbf{v}$ , if we are interested in the **total swirl**, we can equivalently just measure how much the flow  $\mathbf{v}$  follows the closed boundary  $\mathcal{P}$  enclosing the surface  $\mathcal{S}$ .
- ▶ As earlier, by convention we select the orientation of  $d\mathbf{a}$  pointing outward, and the sense of the line  $d\mathbf{l}$  to be anti-clockwise.

# The Fundamental Theorem for Curls


- ▶ As a consequence of the theorem,  $\int_{\mathcal{S}} (\nabla \times \mathbf{v}) \cdot d\mathbf{a}$  depends only on the boundary line  $\mathcal{P}$ , **but not on the particular surface  $\mathcal{S}$  used**, as long as it's circumscribed by  $\mathcal{P}$ . Think of an intact **soap bubble** across a fixed loop. It doesn't matter whether the bubble is convex, concave, or combinations thereof, as long as the loop circumscribing it is fixed.
- ▶ As a consequence we may **deform**  $\mathcal{S}$  for *mathematical convenience* as long as it satisfies the boundary  $\mathcal{P}$ .
- ▶  $\oint_{\mathcal{S}} (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$  for **any closed surface**, since the boundary line, like the mouth of a balloon, shrinks down to a point.
- ▶ HW: Do example 1.11 of [IED].

# Spherical Coordinates



- An *alternative* to using the usual Cartesian coordinates described by  $(x, y, z)$  is to use **spherical coordinates** described by  $(r, \theta, \phi)$  where,
  1.  $r$  is the distance from the origin (the magnitude of the position vector  $\mathbf{r}$ )
  2.  $\theta$  (the angle down from the  $z$  axis) is called the **polar angle**
  3.  $\phi$  (the angle around from the  $x$  axis) is the **azimuthal angle**.

# Spherical Coordinates



- From the above figure we observe that,

$$x = r \sin \theta \cos \phi$$

$$y = r \sin \theta \sin \phi$$


$$z = r \cos \theta$$

- A general vector  $\mathbf{A}$  can be represented in spherical coordinates as,

$$\mathbf{A} = A_r \hat{\mathbf{r}} + A_\theta \hat{\boldsymbol{\theta}} + A_\phi \hat{\boldsymbol{\phi}}$$

where  $\hat{\mathbf{r}}$ ,  $\hat{\boldsymbol{\theta}}$ , and  $\hat{\boldsymbol{\phi}}$  form an **orthogonal** basis set.

# Spherical Coordinates



- In terms of Cartesian unit vectors we can write,<sup>13</sup>,

$$\hat{\mathbf{r}} = \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}}$$


$$\hat{\boldsymbol{\theta}} = \cos \theta \cos \phi \hat{\mathbf{x}} + \cos \theta \sin \phi \hat{\mathbf{y}} - \sin \theta \hat{\mathbf{z}}$$

$$\hat{\boldsymbol{\phi}} = -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}}$$

---

<sup>13</sup>Do the proof.

# The infinitesimal elements in spherical coordinates



- Beware that in the spherical coordinate system,

$$dl \neq dr \hat{r} + d\theta \hat{\theta} + d\phi \hat{\phi}$$

$$d\tau \neq dr d\theta d\phi$$

- An infinitesimal displacement in the  $\hat{r}$  direction is simply  $dr$ , and thus,

$$dl_r = dr$$

- An infinitesimal element of length in the  $\hat{\theta}$  direction is,


$$dl_\theta = r d\theta$$

- An infinitesimal element of length in the  $\hat{\phi}$  direction is,

$$dl_\phi = r \sin \theta d\phi$$



## The infinitesimal elements in spherical coordinates




- ▶ Thus the general **infinitesimal displacement**  $d\mathbf{l}$  is,

$$d\mathbf{l} = dr\hat{\mathbf{r}} + r d\theta\hat{\boldsymbol{\theta}} + r \sin\theta d\phi\hat{\boldsymbol{\phi}}$$

- ▶ The **infinitesimal volume element**  $d\tau$ , in spherical coordinates, is the *product* of the three infinitesimal displacements:

$$d\tau = dl_r dl_\theta dl_\phi = r^2 \sin\theta dr d\theta d\phi$$

## The surface element in spherical coordinates



- Suppose you're integrating over the *surface* of a sphere of radius  $r$ , here,

$$d\mathbf{a}_1 = dl_\theta dl_\phi \hat{\mathbf{r}} = r^2 \sin\theta d\theta d\phi \hat{\mathbf{r}}$$

- On the other hand, if the *surface* lies in the  $xy$  plane, as above<sup>14</sup>

$$d\mathbf{a}_2 = dl_r dl_\phi \hat{\boldsymbol{\theta}} = r dr d\phi \hat{\boldsymbol{\theta}}$$

---

<sup>14</sup>What happened to the  $\sin\theta$  term in the RHS of  $dl_\phi = r \sin\theta d\phi$ ?

## More Formulae for Spherical Coordinates

*Gradient:*

$$\nabla T = \frac{\partial T}{\partial r} \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial T}{\partial \theta} \hat{\boldsymbol{\theta}} + \frac{1}{r \sin \theta} \frac{\partial T}{\partial \phi} \hat{\boldsymbol{\phi}}.$$

*Divergence:*

$$\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta v_\theta) + \frac{1}{r \sin \theta} \frac{\partial v_\phi}{\partial \phi}.$$

*Curl:*

$$\begin{aligned} \nabla \times \mathbf{v} = & \frac{1}{r \sin \theta} \left[ \frac{\partial}{\partial \theta} (\sin \theta v_\phi) - \frac{\partial v_\theta}{\partial \phi} \right] \hat{\mathbf{r}} + \frac{1}{r} \left[ \frac{1}{\sin \theta} \frac{\partial v_r}{\partial \phi} - \frac{\partial}{\partial r} (r v_\phi) \right] \hat{\boldsymbol{\theta}} \\ & + \frac{1}{r} \left[ \frac{\partial}{\partial r} (r v_\theta) - \frac{\partial v_r}{\partial \theta} \right] \hat{\boldsymbol{\phi}}. \end{aligned}$$

*Laplacian:*

$$\nabla^2 T = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial T}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial T}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 T}{\partial \phi^2}.$$

► Things get complicated<sup>15</sup>...

---

<sup>15</sup>The above formulae will be provided, if required, during the exam, and needn't be memorized.

# Cylindrical Coordinates

- Do as HW.

## Need for the Dirac Delta Function

- ▶ Given a body of mass  $M$  located at  $\mathbf{r}_0$ , we may express it in terms of its mass density  $\rho(\mathbf{r})$  as,

$$M = \int_{\mathcal{V}} \rho(\mathbf{r}) d\tau$$

- ▶ But what does the mass density of a **point mass** located at  $\mathbf{r}_0$  look like?
- ▶ It can be unequal to zero only at a single point, *i.e.*,

$$\rho(\mathbf{r}) = 0 \quad \forall \mathbf{r} \neq \mathbf{r}_0$$

- ▶ The volume integral, however,

$$\int_{\mathcal{V}} \rho(\mathbf{r}) d\tau = M \quad (\text{finite})$$

is **finite** provided  $\mathbf{r}_0$  lies within the volume  $\mathcal{V}$ .

# Need for the Dirac Delta Function

- ▶ It was a long time coming.
- ▶ Since there was no function that encoded such a property, **Paul A. M. Dirac** **invented one**, writing  $\rho(\mathbf{r})$  as,


$$\rho(\mathbf{r}) = M \times \delta(\mathbf{r} - \mathbf{r}_0)$$

, and *requiring* that,

$$\int_{\mathcal{V}} \delta(\mathbf{r} - \mathbf{r}_0) d\tau = \begin{cases} 1 & \text{if } \mathbf{r}_0 \in \mathcal{V} \\ 0 & \text{otherwise} \end{cases}$$

$$\delta(\mathbf{r} - \mathbf{r}_0) = 0 \quad \forall \mathbf{r} \neq \mathbf{r}_0$$

# The One-Dimensional Dirac Delta Function



- ▶ The one-dimensional **Dirac delta function**<sup>16</sup>,  $\delta(x - a)$ , can be pictured as an infinitely high, but infinitesimally narrow spike, with area 1, located at  $x = a$ , *i.e.*,

$$\delta(x - a) \equiv \begin{cases} 0 & x \neq a \\ \infty & x = a \end{cases}$$

and,

$$\int_{-\infty}^{\infty} \delta(x - a) dx = 1$$

---

<sup>16</sup>If  $x$  has units of length, what's the unit of  $\delta(x)$ ?

# Properties of the One-Dimensional Dirac Delta Function

- ▶ *Technically*,  $\delta(x)$  is not a function at all, since its value is not finite at  $x = 0$ .
- ▶ It's **even**, i.e.,  $\delta(x) = \delta(-x)$ .
- ▶ An important characteristic of the Dirac delta function is its **sifting property**<sup>17</sup>,

$$\int_{-\infty}^{\infty} f(x) \delta(x - x_0) dx = f(x_0) \int_{-\infty}^{\infty} \delta(x - x_0) dx = f(x_0)$$

where, *loosely speaking*, the delta function *picks out* the value of  $f(x)$  at  $x = x_0$ , i.e.,  $f(x_0)$ .

- ▶ Another **curious** property of the Dirac delta function is,

$$\delta(kx) = \frac{1}{|k|} \delta(x)$$

where  $k$  is any (non-zero) constant

---

<sup>17</sup>Prove the sifting property!



# The One-Dimensional Dirac Delta Function

- ▶ If  $f(x)$  is differentiable,

$$\int_{-\infty}^{\infty} f(x) \delta'(x - x_0) dx = -f'(x_0)$$

- ▶ The  $\delta(x)$  may also be seen as the derivative of the **Heaviside step function**<sup>18</sup>,

$$\delta(x - a) = \frac{d}{dx} \Theta(x - a)$$

---

<sup>18</sup>This property might be invoked while discussing square potential barriers in the QM part of this course.

# The Three-Dimensional Dirac Delta Function

- ▶ In three dimensions,

$$\underbrace{\delta^3(\mathbf{r})}_{\text{shorthand}} \equiv \delta(x)\delta(y)\delta(z)$$


and,

$$\int_{\text{all space}} \delta^3(\mathbf{r}) d\tau \equiv 1$$

- ▶ Also,

$$\int_{\text{all space}} f(\mathbf{r}) \delta^3(\mathbf{r} - \mathbf{r}_0) d\tau = f(\mathbf{r}_0)$$

## Divergence of $\hat{\mathbf{r}}/r^2$



- ▶ Consider the vector function<sup>19</sup>,

$$\mathbf{v} = \frac{1}{r^2} \hat{\mathbf{r}}$$

- ▶ From the figure, evidently  $\mathbf{v}$  has a **LARGE positive divergence** at the center,
- ▶ ...and **yet...** (do the math)

▶


$$\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) = \frac{1}{r^2} \frac{\partial}{\partial r} (1) = 0 \quad \text{everywhere!?!}$$

which is certainly not what we were expecting!

---

<sup>19</sup>Which is eerily reminiscent of  $\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0 r^2} \hat{\mathbf{r}}$ , the electric field due to a **single, static point** charge.

## Divergence of $\hat{\mathbf{r}}/r^2$



- ▶ However, when we consider the **divergence theorem**, i.e.,


$$\int_V (\nabla \cdot \mathbf{v}) d\tau = \oint_{\mathcal{S}} \mathbf{v} \cdot d\mathbf{a}$$

the RHS is...

- ▶ ...,

$$\oint_{\mathcal{S}} \mathbf{v} \cdot d\mathbf{a} = \int \left( \frac{1}{r^2} \hat{\mathbf{r}} \right) \cdot (r^2 \sin\theta d\theta d\phi \hat{\mathbf{r}}) = 4\pi$$

## Divergence of $\hat{\mathbf{r}}/r^2$




- ▶ But we'd just (*albeit naïvely*) found that the LHS, i.e.,  $\int_V (\nabla \cdot \mathbf{v}) d\tau = 0$ , which **contradicts** the **divergence theorem**!
- ▶ However,  $\int_V (\nabla \cdot \mathbf{v}) d\tau = 0$  is **incorrect** –the **source** of the problem being the point  $r = 0$ , where  $\mathbf{v} = \infty$ , i.e., it BLOWS UP.
- ▶ Indeed,  $\nabla \cdot \mathbf{v}$  is actually *zero* everywhere **except the origin**.<sup>20</sup>

---

<sup>20</sup>**Beware!** The intuition of observing the spreading of sawdust at any of the **non-central** points seems to suggest a non-zero divergence, at least to my eye.

## Divergence of $\hat{\mathbf{r}}/r^2$



- ▶ We resolve this *paradox* by realizing that the volume integral of  $\nabla \cdot (\hat{\mathbf{r}}/r^2)$  must yield a constant  $4\pi$ . (Why because the RHS of the **divergence theorem** just said so.)
- ▶ Thus, we can write using **the definition of the Dirac delta function**,

$$\boxed{\nabla \cdot (\hat{\mathbf{r}}/r^2) = 4\pi\delta^3(\mathbf{r})}$$

- ▶ *Alternatively*,  $\frac{1}{4\pi}\nabla \cdot (\hat{\mathbf{r}}/r^2)$  is a **concrete representation** of the Dirac delta function  $\delta(\mathbf{r})$ .

## The Scalar Potential $V$

- ▶ When  $\nabla \times \mathbf{E} = 0$  everywhere<sup>21</sup>, **Stokes' theorem** tells us that,

$$\oint \mathbf{E} \cdot d\mathbf{l} = 0$$

- ▶ On the other hand, the **fundamental theorem of gradients** allows us to write,

$$\oint -\nabla V \cdot d\mathbf{l} = 0$$

- ▶ Thus,  $\mathbf{E}$  can be written as<sup>22</sup> the **gradient of a scalar potential  $V$** ,

$$\mathbf{E} = -\nabla(V + \text{const.})$$

- ▶ The potential is **not unique**, *i.e.*, any **constant** can be added to  $V$  **without affecting** its gradient, *i.e.*, the (negative) electric field  $-\mathbf{E}$ .

---

<sup>21</sup> $\nabla \times \mathbf{E} = 0$  is guaranteed only in the *electrostatic* regime.

<sup>22</sup>The negative sign is purely a matter of convention.

## Curl-less (or irrotational) fields $\mathbf{v}_{irr}$

- ▶ The condition  $\nabla \times \mathbf{E} = 0$  everywhere is **equivalent** to,
  - ▶  $\int_a^b \mathbf{E} \cdot d\mathbf{l}$  is independent of path, for any given end points, a consequence of the fundamental theorem of gradients.
  - ▶  $\oint \mathbf{E} \cdot d\mathbf{l} = 0$ .
  - ▶  $\mathbf{E}$  is the gradient of some scalar function:  $\mathbf{E} = -\nabla V$
- ▶ Irrotational fields  $\mathbf{v}_{irr}$ , i.e.,  $\nabla \times \mathbf{v}_{irr} = 0$ , are also known as **conservative fields**.<sup>23</sup>
- ▶ We'll soon **prove** in the chapter on **Electrostatics** that the **work done** in moving a test charge against a background of **static** charges is **independent** of the path taken.

---

<sup>23</sup>In **mechanics**, a **conservative force**  $\mathbf{F} = -\nabla U$  is a force with the property that the total work done  $W = \int_a^b \mathbf{F} \cdot d\mathbf{l}$  in moving a particle between point  $a$  to  $b$  is *independent* of the path taken.



# The Vector Potential $\mathbf{A}$

- ▶ **Maxwell's equations** *guarantee* that

$$\nabla \cdot \mathbf{B} = 0$$

- ▶ This allows  $\mathbf{B}$  to be written as the curl of a **vector potential**  $\mathbf{A}$ ,

$$\mathbf{B} = \nabla \times (\mathbf{A} + \nabla T)$$

since mathematically (4) **the divergence of a curl is always zero**, i.e.,  $\nabla \cdot (\nabla \times \mathbf{v}) = 0$

- ▶ The vector potential  $\mathbf{A}$  is **not unique**, i.e., *the gradient of any scalar function*  $+\nabla T$  can be added to  $\mathbf{A}$  without affecting its curl, i.e., the magnetic field  $\mathbf{B}$ , since mathematically (2) **the curl of a gradient is always zero**, i.e.,  $\nabla \times (\nabla T) = 0$ .

## Divergence-less (or solenoidal) fields

- ▶ The condition  $\nabla \cdot \mathbf{B} = 0$  everywhere is *equivalent* to,
  - ▶  $\oint \mathbf{B} \cdot d\mathbf{a} = 0$  for any closed surface, a direct consequence of the divergence theorem.
  - ▶  $\mathbf{B}$  is the curl of some vector function:  $\mathbf{B} = \nabla \times \mathbf{A}$
  - ▶  $\int_{\mathcal{S}} \mathbf{B} \cdot d\mathbf{a}$  is independent of the details of the precise **open** surface  $\mathcal{S}(\mathcal{P})$ , once its periphery  $\mathcal{P}$  is set.<sup>24</sup>

---

<sup>24</sup>This means we can deform the surface as we please as long as the boundary is kept fixed.

# The Helmholtz Theorem

- ▶ "A well-behaved (*i.e.*, goes to zero at infinity) vector field is **uniquely** specified by its **divergence and curl** (and, in the case of a *finite region*, additionally by its normal component over the entire boundary.)"
- ▶ *i.e.*, suppose we know over all space.<sup>25</sup>

$$\nabla \cdot \mathbf{F}(\mathbf{r}) = D(\mathbf{r}) \quad \text{and,} \quad \nabla \times \mathbf{F}(\mathbf{r}) = \mathbf{C}(\mathbf{r})$$

- ▶ Then the **unique** vector field  $\mathbf{F}$  is given by

$$\mathbf{F}(\mathbf{r}) = -\nabla U(\mathbf{r}) + \nabla \times \mathbf{W}(\mathbf{r}) \quad \text{Helmholtz Decomposition}$$

where,

$$U(\mathbf{r}) = \frac{1}{4\pi} \int_{\text{all space}} d^3 r' \overbrace{\frac{\nabla' \cdot \mathbf{F}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}}^{D(\mathbf{r}')} \quad \text{and,} \quad \mathbf{W}(\mathbf{r}) = \frac{1}{4\pi} \int_{\text{all space}} d^3 r' \overbrace{\frac{\nabla' \times \mathbf{F}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}}^{\mathbf{C}(\mathbf{r}'')}$$

where the  $\nabla'$  denotes that the derivatives are to be taken *w.r.t.* source points  $\mathbf{r}'$ .

---

<sup>25</sup>Note that since (4):  $\nabla \cdot (\nabla \times \mathbf{v}) = 0$ , we must have  $\nabla \cdot \mathbf{C} = 0$  for consistency.

## Why's Helmholtz Theorem Useful?

- ▶ Since a vector field is completely specified once its divergence and curl are known (a purely mathematical result), and we know that the study of electromagnetism involves the vector fields  $\mathbf{E}$  and  $\mathbf{B}$ , we can already *guess*...
- ▶ ...the laws of electromagnetism:

$$\nabla \cdot \mathbf{E} = \text{something}_1$$

$$\nabla \times \mathbf{E} = \text{something}_2$$

$$\nabla \cdot \mathbf{B} = \text{something}_3$$

$$\nabla \times \mathbf{B} = \text{something}_4$$

which look exactly like **Maxwell's equations**<sup>26</sup>, which are *thus mathematically sufficient* to reconstruct  $\mathbf{E}$ , and  $\mathbf{B}$ .

- ▶ Even further, we can *almost solve* for the fields  $\mathbf{E}$  and  $\mathbf{B}$  via the **Helmholtz decomposition**, even without explicitly knowing what the  $RHS = \text{something}_i$  are!

<sup>26</sup>If you're wondering where  $\mathbf{D}$  and  $\mathbf{H}$  went, here we're discussing the so-called **microscopic** representation of Maxwell's equations.  $\mathbf{D}$  and  $\mathbf{H}$  appear in the **macroscopic** formulation where the material medium is built into the equations. Both formulations are *equally* general.

# Why's Helmholtz Theorem Useful?

- ▶ Considering the **static**<sup>27</sup> version of Maxwell's equations

$$\nabla \cdot \mathbf{E} = \frac{\rho(\mathbf{r})}{\epsilon_0}$$

$$\nabla \times \mathbf{E} = 0$$

$$\nabla \cdot \mathbf{B} = 0$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{j}(\mathbf{r})$$

we're **gratified** by noting that a given set of **stationary** charges  $\rho(\mathbf{r})$ , and **steady** currents  $\mathbf{j}(\mathbf{r})$ , may only generate **one** possible steady  $\mathbf{E}$ , and **one** possible steady  $\mathbf{B}$  field.

- ▶ Similarly, if all the sources  $\rho$ , and currents  $\mathbf{j}$  are **zero everywhere**, then the **only** physical solution is  $\mathbf{E} = \mathbf{B} = 0$ .
- ▶ This implies that **static electric and magnetic fields cannot generate themselves**, *i.e.*, there must be stationary charges and steady currents generating them!

---

<sup>27</sup>*i.e.*, all charges are stationary, and currents steady.

# Limitations of our Treatment of the Helmholtz Theorem

- ▶ Our discussion for the Helmholtz theorem only works for **time-independent** sources, and currents.
- ▶ For the **time-dependent** case, while the divergence and curl still uniquely identify the vector field, the **Helmholtz decomposition** looks a bit different, though I will not be writing it down explicitly.
- ▶ A second issue is that we haven't grappled with the application of the **boundary conditions** (*i.e., the normal component of the field must be known at all points of the periphery*) that need to be imposed if we're considering a **finite region of space**.