

Physics Behind the Simulation: A CS251 Report by Group 11.

Anmol Arora

Roll No. 130050027 anmolarora59@gmail.com

Pranjal Khare

Roll No. 130050028 pranjalkhare7@gmail.com

Aman Goel

Roll No. 130050041 amangoelvsec@gmail.com

Sept 29, 2014

Overview

- 1 Introduction
- 2 Body
 - Part 1: Pendulum
 - Part 2: Flying Ball
 - Part 3: Collision
- 3 Conclusion
- 4 References



The Purpose of this Report

We identified the "behind the scenes" equations which are used for simulation by Box2D.

The Purpose of this Report

We identified the "behind the scenes" equations which are used for simulation by Box2D. In this report we have identified three physical phenomenon which are involved in the rube goldberg simulation.

The Purpose of this Report

We identified the "behind the scenes" equations which are used for simulation by Box2D. In this report we have identified three physical phenomenon which are involved in the rube goldberg simulation. We have tried to describe the equations and point out the specific occurrences of the corresponding physical phenomenon in our simulation

The Purpose of this Report

We identified the "behind the scenes" equations which are used for simulation by Box2D. In this report we have identified three physical phenomenon which are involved in the rube goldberg simulation. We have tried to describe the equations and point out the specific occurrences of the corresponding physical phenomenon in our simulation

This report is made through beamer which is a very useful tool. It is much more powerful than other presentation softwares such as MS Power Point.

The Purpose of this Report

We identified the "behind the scenes" equations which are used for simulation by Box2D. In this report we have identified three physical phenomenon which are involved in the rube goldberg simulation. We have tried to describe the equations and point out the specific occurrences of the corresponding physical phenomenon in our simulation

This report is made through beamer which is a very useful tool. It is much more powerful than other presentation softwares such as MS Power Point. There are many predefined templates, which can be applied very easily, and make the presentation attractive and interactive.

The Purpose of this Report

We identified the "behind the scenes" equations which are used for simulation by Box2D. In this report we have identified three physical phenomenon which are involved in the rube goldberg simulation. We have tried to describe the equations and point out the specific occurrences of the corresponding physical phenomenon in our simulation

This report is made through beamer which is a very useful tool. It is much more powerful than other presentation softwares such as MS Power Point. There are many predefined templates, which can be applied very easily, and make the presentation attractive and interactive.

Part 1: Pendulum

The Formula for the time period of a pendulum is: [3]

Part 1: Pendulum

The Formula for the time period of a pendulum is: [3]

$$T = 2\pi\sqrt{\frac{l}{g}} \quad (1)$$

Part 1: Pendulum

The Formula for the time period of a pendulum is: [3]

$$T = 2\pi\sqrt{\frac{l}{g}} \quad (1)$$

- Here l is the length of thread from bob to pivot point and g is the acceleration due to gravity ($\approx 9.81m/sec^2$)

Part 1: Pendulum

The Formula for the time period of a pendulum is: [3]

$$T = 2\pi\sqrt{\frac{l}{g}} \quad (1)$$

- Here l is the length of thread from bob to pivot point and g is the acceleration due to gravity ($\approx 9.81m/sec^2$)
- g is the acceleration due to gravity.

Part 1: Pendulum

The Formula for the time period of a pendulum is: [3]

$$T = 2\pi\sqrt{\frac{l}{g}} \quad (1)$$


- Here l is the length of thread from bob to pivot point and g is the acceleration due to gravity ($\approx 9.81m/sec^2$)
- g is the acceleration due to gravity.
- v_x is the initial velocity of ball in x-direction (No acceleration in x-direction).

Part 1: Pendulum

The Formula for the time period of a pendulum is: [3]

$$T = 2\pi\sqrt{\frac{l}{g}} \quad (1)$$

- Here l is the length of thread from bob to pivot point and g is the acceleration due to gravity ($\approx 9.81m/sec^2$)
- g is the acceleration due to gravity.
- v_x is the initial velocity of ball in x-direction (No acceleration in x-direction).



Part 2: Flying Ball

The equation of motion of a ball in free fall is: [2]

$$s_y = u_y t + \frac{1}{2} g t^2 \quad (2)$$

$$s_x = v_x t \quad (3)$$

- s_y is the displacement in the y direction and s_x is the displacement in x direction.

Part 2: Flying Ball

The equation of motion of a ball in free fall is: [2]

$$s_y = u_y t + \frac{1}{2} g t^2 \quad (2)$$

$$s_x = v_x t \quad (3)$$

- s_y is the displacement in the y direction and s_x is the displacement in x direction.
- g is the acceleration due to gravity.

Part 2: Flying Ball

The equation of motion of a ball in free fall is: [2]

$$s_y = u_y t + \frac{1}{2} g t^2 \quad (2)$$

$$s_x = v_x t \quad (3)$$

- s_y is the displacement in the y direction and s_x is the displacement in x direction.
- g is the acceleration due to gravity.
- v_x is the initial velocity of ball in x-direction (No acceleration in x-direction).


Part 2: Flying Ball

The equation of motion of a ball in free fall is: [2]

$$s_y = u_y t + \frac{1}{2} g t^2 \quad (2)$$

$$s_x = v_x t \quad (3)$$

- s_y is the displacement in the y direction and s_x is the displacement in x direction.
- g is the acceleration due to gravity.
- v_x is the initial velocity of ball in x-direction (No acceleration in x-direction).



Part 3: Collision

Collision is dependent on the surfaces involved to a large extent.
The principle equations involved are:[1]

Part 3: Collision

Collision is dependent on the surfaces involved to a large extent.
The principle equations involved are:[1]

$$m_1 v_{i1} + m_2 v_{i2} = m_1 v_{f1} + m_2 v_{f2} \quad (4)$$

Part 3: Collision

Collision is dependent on the surfaces involved to a large extent.
The principle equations involved are:[1]

$$m_1 v_{i1} + m_2 v_{i2} = m_1 v_{f1} + m_2 v_{f2} \quad (4)$$

$$e = \frac{v_{f2} - v_{f1}}{v_{i2} - v_{i1}} \quad (5)$$

Part 3: Collision

Collision is dependent on the surfaces involved to a large extent.
The principle equations involved are:[1]

$$m_1 v_{i1} + m_2 v_{i2} = m_1 v_{f1} + m_2 v_{f2} \quad (4)$$

$$e = \frac{v_{f2} - v_{f1}}{v_{i2} - v_{i1}} \quad (5)$$

- e is a constant for two surfaces and is called coefficient of restitution .It is dimensionless.

Part 3: Collision

Collision is dependent on the surfaces involved to a large extent.
The principle equations involved are:[1]

$$m_1 v_{i1} + m_2 v_{i2} = m_1 v_{f1} + m_2 v_{f2} \quad (4)$$

$$e = \frac{v_{f2} - v_{f1}}{v_{i2} - v_{i1}} \quad (5)$$

- e is a constant for two surfaces and is called coefficient of restitution .It is dimensionless.
- v_{1i} , v_{1f} , v_{2i} and v_{2f} are the velocities of body 1 and body 2 before and after the collision.


Part 3: Collision

Collision is dependent on the surfaces involved to a large extent.
The principle equations involved are:[1]

$$m_1 v_{i1} + m_2 v_{i2} = m_1 v_{f1} + m_2 v_{f2} \quad (4)$$

$$e = \frac{v_{f2} - v_{f1}}{v_{i2} - v_{i1}} \quad (5)$$

- e is a constant for two surfaces and is called coefficient of restitution .It is dimensionless.
- v_{1i} , v_{1f} , v_{2i} and v_{2f} are the velocities of body 1 and body 2 before and after the collision.



Conclusion

We have covered the following:

Conclusion

We have covered the following:

- Identifying three different Physics phenomenon applicable to our Box2D simulation.

Conclusion

We have covered the following:

- Identifying three different Physics phenomenon applicable to our Box2D simulation.
 - Colissions

Conclusion

We have covered the following:

- Identifying three different Physics phenomenon applicable to our Box2D simulation.
 - Colissions
 - Free Fall in gravity

Conclusion

We have covered the following:

- Identifying three different Physics phenomenon applicable to our Box2D simulation.
 - Colissions
 - Free Fall in gravity
 - Simple Harmonic Motion - Motion of pendulum.

Conclusion

We have covered the following:

- Identifying three different Physics phenomenon applicable to our Box2D simulation.
 - Colissions
 - Free Fall in gravity
 - Simple Harmonic Motion - Motion of pendulum.
- Identifying the equations Box2D engine might be using to simulate these phenomenon.

Conclusion

We have covered the following:

- Identifying three different Physics phenomenon applicable to our Box2D simulation.
 - Colissions
 - Free Fall in gravity
 - Simple Harmonic Motion - Motion of pendulum.
- Identifying the equations Box2D engine might be using to simulate these phenomenon.
- Describing the identified equations.

Conclusion

We have covered the following:

- Identifying three different Physics phenomenon applicable to our Box2D simulation.
 - Colissions
 - Free Fall in gravity
 - Simple Harmonic Motion - Motion of pendulum.
- Identifying the equations Box2D engine might be using to simulate these phenomenon.
- Describing the identified equations.

References



Wikipedia.
Collision.

<http://en.wikipedia.org/wiki/Collision>.



Wikipedia.
Equations for a falling body.

http://en.wikipedia.org/wiki/Equations_for_a_falling_body.



Wikipedia.
Pendulum (mathematics).

[http://en.wikipedia.org/wiki/Pendulum_\(mathematics\)](http://en.wikipedia.org/wiki/Pendulum_(mathematics)).