

A 19 years old woman approaches the doctor with her boyfriend for second trimester medical termination of pregnancy. Enumerate ethical issues and communication strategies.



AAB0100022201-P1

5. Short Answer Questions (Any 4 out of 5) :

[4 × 7 = 28]

- a) Define adverse drug reaction. Mention teratogenic drugs with their effects
- b) Classify hypolipidemic agents. Write mechanism of action, adverse effects and uses of atorvastatin.
- c) Classify alpha blockers. Write six uses and four adverse effects of prazosin.
- d) Describe pharmacotherapy of bronchial asthma.
- e) Classify anticoagulants. Compare and contrast between low molecular weight and unfractionated heparin.

6. Structured Long Answer Question (Compulsory) :

[1 × 12 = 12]

Classify para-sympatholytic agents. Write their uses and adverse effects.

~~~~~




**5. Short Answer Questions (Any 4 out of 5) :****[4 × 7 = 28]**

- a) Define adverse drug reaction. Mention teratogenic drugs with their effects
- b) Classify hypolipidemic agents. Write mechanism of action, adverse effects and uses of atorvastatin.
- c) Classify alpha blockers. Write six uses and four adverse effects of prazosin.
- d) Describe pharmacotherapy of bronchial asthma.
- e) Classify anticoagulants. Compare and contrast between low molecular weight and unfractionated heparin.

**6. Structured Long Answer Question (Compulsory) :****[1 × 12 = 12]**

Classify para-sympholytic agents. Write their uses and adverse effects.





14. Contraction of circular muscle of iris resulting in miosis is an example of action on
- ☒ A) muscarinic receptors
  - B) nicotinic receptors
  - C) alpha - adrenergic receptors
  - D) beta - adrenergic receptors
15. The clinical use of dobutamine is due to its action mainly on
- A) D1 receptor
  - B) D2 receptor
  - C) alpha1 - adrenergic receptor
  - ☒ D) beta1 - adrenergic receptor
16. An eye drop which causes miosis and increase in trabecular outflow of aqueous humor is
- A) dorzolamide
  - B) apraclonidine
  - C) levobunolol
  - ☒ D) pilocarpine
17. The duration of action is longest in the case of the following skeletal muscle relaxant
- A) d-Tubocurarine
  - ☒ B) Pancuronium
  - C) Vecuronium
  - D) Succinylcholine
18. The daily dosage of methylcobalamin to treat peripheral neuropathy is
- ☒ A) 0.5-1.5mg
  - B) 1.5-2.5 mg
  - C) 2.5-3.5mg
  - ☒ D) 3.5-4.5mg
19. Response to erythropoietin is poor in
- A) anaemia of CKD
  - ☒ B) nutritional anaemia
  - C) cancer chemotherapy induced anaemia
  - D) anaemia in HIV patients
20. The important adverse effect associated with niacin therapy for dyslipidemia is
- A) hypoglycaemia
  - B) acute kidney disease
  - ☒ C) serious liver damage
  - D) pancreatitis