mhs-poisson-ltau: implementation

Technical notes on `run.mode = "mhs-poisson-ltau"`. HSEMHS, 24 Oct 2026.

1. Running it

```
[mpiexec -n R] julia --project=. -t T scripts/run.jl config/modes/mhs-poisson-ltau.toml
```

Driver `run_mhs_poisson_ltau` (`src/RunModes.jl`); solver `PoissonTau.solve_tau` (`src/PoissonTau.jl`); MPI layer `src/Slab.jl`; PCG and threaded vector updates from `PoissonZ`. Input and `[tau_grid]` keys as `column-hse-ltau`; in addition:

Key	Default	Meaning
`poisson.logtau_cut`	-4	solved levels $\log\tau \geq$ cut (nearest level k_c)
`poisson.max_steps`	100	Levenberg–Marquardt steps
`poisson.rtol`, `n_consec`	10^{-4}, 3	stop when $(\Phi_{\rm prev}-\Phi)/\Phi_{\rm prev} <$ rtol on `n_consec` consecutive accepted steps (objective only)
`poisson.cg_tol`, `cg_maxit`	10^{-3}, 80	PCG per trial
`poisson.dc_cap_m`	3×10^5	largest height change of a column per step (whole-step scaling)
`poisson.lm_mu0`	0	initial damping

2. Flow

`column_hse_tau` (aligned, gauged) gives $u=\ln P$ and z_0 on all levels; `solve_tau` solves for u on levels $k>k_c$ and $c=z_{k_c}$ per column, with $u_{k\leq k_c}$ and $z_k-z_{k_c}$ $(k<k_c)$ fixed; the heights are gauged (mean $z(\tau{=}1)=0$); output in the `column-hse-ltau` layout.

3. State, heights and residuals (`evaluate!`)

Nodes $m=1\ldots n_1$ are levels $k=k_c\ldots n_\tau$. The unknown vector per column has n_1 slots: slot $1=c$, slots $2\ldots n_1=u$ at $k>k_c$ (one array (n_1,N_y,n_x), so FFTs and halos treat it uniformly). At each node the EOS gives $\ln\rho$, $\partial\ln\rho/\partial\ln P$ and $\ln\chi$; $\partial\ln\chi/\partial\ln P$ by a central difference ($\varepsilon=10^{-4}$, the table returns no opacity derivative). With $a=\tau/\chi$, $q=\rho/P$:

$$z_{k+1}=z_k-\tfrac12\Delta\ln\tau_k\,(a_k+a_{k+1}),\qquad z_{k_c}=c.$$

Lorentz force at all levels (`lorentz_tau!`): for each component f of $\mathbf{B}$, $\partial_z f=(f_{k-1}-f_{k+1})/(z_{k-1}-z_{k+1})$ (one-sided at the ends), $\partial_x f|_z=D_x f-D_x z\,\partial_z f$, likewise y; $\mathbf{J}=\nabla\times\mathbf{B}/\mu_0$; $\mathbf{L}=\mathbf{J}\times\mathbf{B}$. Residuals with $W=gq-L_z e^{-u}$, $w=Wa$:

$$R_x=-D_x u-W\,D_x z+L_x e^{-u},\qquad R_{z,f}=u_{f+1}-u_f-\tfrac12\Delta\ln\tau_f\,(w_f+w_{f+1}).$$

Weights from the start state (fixed): $\bar H_m$ global median of $1/(gq)$, $\overline{\Delta z}_f$ global median of the level thickness, $w_c=\overline{\Delta z}_{\rm node}\bar H^2$, $w_f=\bar H_f\bar H_{f+1}/\overline{\Delta z}_f$; $\Phi=\sum w_c(R_x^2+R_y^2)+\sum w_f R_z^2$.

4. Jacobian

Exact, including $\mathbf{L}$'s dependence on the heights (`Jop!`, adjoint `JTop!`):

- **Triangular operator.** $\delta z_m=\delta c+\sum_{f<m}\frac12\Delta\ln\tau_f\,(a\chi_u)_f\delta u_f+(a\chi_u)_{f+1}\delta u_{f+1}$ with $\chi_u=\partial\ln\chi/\partial\ln P$: a cumulative sum down each column. Its adjoint is a reverse cumulative sum.
- Vertical: $\delta R_{z,f}=\delta u_{f+1}-\delta u_f-\frac12\Delta\ln\tau_f(w'_f\delta u_f+w'_{f+1}\delta u_{f+1})$, $w'=\partial w/\partial u=W'a-Wa\chi_u$, $W'=gq(\rho_u-1)+L_z e^{-u}$.
- Horizontal: $\delta R_x=-D_x\delta u-W\,D_x\delta z+\ell_x\delta u$, $\ell_x=-D_x z\,W'-L_x e^{-u}$.
- **Lorentz through the heights.** With $\zeta_m=(\delta z_{m-1}-\delta z_{m+1})/(z_{m-1}-z_{m+1})$ (the relative change of the vertical spacing; δz above the cut $=\delta c$): $\delta(\partial_z f)=-\partial_z f\,\zeta$, $\delta(\partial_x f|_z)=-\partial_z f\,D_x\delta z+D_x z\,\partial_z f\,\zeta$. Hence $\delta\mathbf{J}=\mathbf{J}_\zeta\zeta+\mathbf{J}_X D_x\delta z+\mathbf{J}_Y D_y\delta z$ with $\mathbf{J}_\zeta=(z_yB_{z,z}+B_{y,z},\,-B_{x,z}-z_xB_{z,z},\,z_xB_{y,z}-z_yB_{x,z})/\mu_0$, $\mathbf{J}_X=(0,B_{z,z},-B_{y,z})/\mu_0$, $\mathbf{J}_Y=(-B_{z,z},0,B_{x,z})/\mu_0$, and $\delta\mathbf{L}=\delta\mathbf{J}\times\mathbf{B}$. It enters δR_x through $e^{-u}(\delta L_x+z_x\delta L_z)$ and δR_z through $e^{-u}a\,\delta L_z$: nine node-local coefficient arrays.
- Checks (`MHS_CHECK` in the script version): adjoint $\langle Jv,r\rangle=\langle v,J^\top r\rangle$ to 4×10^{-16}; Jv against central differences of $\mathbf{R}$ to 10^{-9} (with $\mathbf{B}$), 10^{-10} (without). With $\mathbf{L}$ held fixed the directional error was 100 % in magnetic regions and the solve stalled after one step.

5. Linear solve and preconditioner

Normal equations $(J^\top W J + E + \mu D)\,\delta = -J^\top W \mathbf{R}$ by PCG with global inner products; $E = 10^{-2}\kappa^2 D$, $D = (\sum w_c \bar{W}^2,\ w_{c,2}, \dots)$ (the damping shape; slot 1 for c, which no vertical equation sees).

Preconditioner (`setprec!`, `Minv!`): the operator with layer-median coefficients ($\overline{a\chi_u}$, $\overline{w'}$, $\bar{W}$) is, per horizontal mode, $M(s^2) = A + s^2 B$ with s^2 as in `mhs-poisson-z`; $A = G_z^\top W_f G_z + E + \mu D$ (vertical rows), $B = G_h^\top W_c G_h$, $G_h = -(I + \bar{W}\,S)$ the horizontal symbol divided by ik including the triangular matrix S. Both are dense $n_1 \times n_1$ ($n_1 = 55$). Once per step: Cholesky $A = LL^\top$, eigen-decomposition $L^{-1}BL^{-\top} = V\Lambda V^\top$, $Q = V^\top L^{-1}$; then $M(s^2)^{-1} = Q^\top \operatorname{diag}(1/(1 + s^2\lambda))\,Q$ for every mode: two complex GEMMs over all local modes of the transposed FFT layout (`SlabFFT`, as in `mhs-poisson-z`), BLAS threaded during the solve.

6. Steps

Levenberg–Marquardt: for μ (start `lm_mu0`) solve, scale the *whole* step by $\min(1,\ 2/\max|\delta u|,\ \texttt{dc_cap}/\max|\delta c|)$, try up to 4 halvings on the true Φ (EOS status checked); if none decreases Φ, $\mu \leftarrow \max(4\mu, 1)$ and re-solve (up to 8 times). After acceptance (source rule): $\mu \leftarrow \mu/3$ if the accepted step length factor is exactly $\alpha = 1$; unchanged if $\alpha \geq s_c/2$ (s_c the whole-step scale); $\times 2$ otherwise; $\mu < 10^{-3} \to 0$. When the whole-step scaling is active ($s_c < 1$), a full scaled step ($\alpha = s_c$) therefore does *not* lower μ; whether that was intended is a code question (follow-up). Element-wise clamping of δc (the first version) turned the direction into an ascent direction where the step asked for > 1 Mm in a few columns: Φ rose linearly with the step length.

Termination. (a) $(\Phi_{\text{prev}} - \Phi)/\Phi_{\text{prev}} <$ `rtol` on `n_consec` consecutive accepted steps (log line `converged: ...`); (b) no damping level gives a decrease (`no decrease up to μ = ...; stopping`); (c) `max_steps` reached (no message). Stopping uses the objective only. Two earlier rules are gone: a single small step (it stopped one run after 17 steps while strong-field columns still moved 80 km per step), and a height shortcut (one small step plus $\max|\delta c| < 100$ m), removed on 24 Oct 2026 in line with the Φ-only policy. The logged $\max|\delta c|$ is the largest change of a column's *cut-level* height in the step, not the change of its $\tau = 1$ height (which also depends on the changed layer thicknesses). The driver returns `converged`, exit 0, in all three cases: read the log (code follow-up: propagate the termination). A met rule does not certify small residuals.

7. Parallelisation

x slabs; halo exchange in x for every D_x (heights, u, $\mathbf{B}$, and in the Jacobian); global sums; exact global medians for the weights and the preconditioner (bisection on order-preserving integer keys); the distributed FFT (rfft in y, `Alltoallv` transpose, FFT in x) and GEMMs on the local k_y slab. Threads in all node loops; BLAS is raised to `min(32, nthreads)` for the solve and reset to 1 afterwards; FFTW plans are single-threaded, applied from Julia threads.

The field values on the τ levels are used unchanged at the new heights; no discrete $\nabla \cdot \mathbf{B} = 0$ is imposed (the staggered construction of `full-mhs-z` needs a fixed geometry). Above the cut the levels stay at their column-HSE pressures and relative heights and move rigidly with c; the global height offset is fixed only by the final gauge.

8. Verification and performance

- 96^2 magnetic crop (manual checks, no automated test), ranks $\times$ threads 1×8, 1×32, 4×8, consecutive-step rule: 37 steps each; $z(\tau{=}1)$ differences median 1 m, 99th percentile 12–24 m, max 150 m (weakly determined strong-field columns). Agreement to rounding only; identical trajectories across layouts are not promised. After the removal of the height shortcut (commit 813a3af) the 1×32 run again took 37 steps.
- Against the original script `scripts/poisson/poisson_tau_direct.jl` (single-step rule) on the same crop: $z(\tau{=}1)$ differences median 3.6 m, 99th percentile 95 m, max 1 km, because the two stopped one step apart while some columns were still moving.
- Full cube ($504 \times 504 \times 95$, 54 solved $\ln P$ levels + 1 height per column, 1.4×10^7 unknowns), with the superseded single-step rule: one node, 96 threads, 17 steps, 14 min, 11 GB (commit b96bb4e); 8 ranks on 4 nodes, 8.3 min, 3.1 GB per rank, same summary statistics (commit 13666b7, log `out/modes/mhs-poisson-ltau.log` in the cluster worktree `MHS_modes`). The current rule has not yet been run on the full cube. PCG reaches its 80-iteration cap in every step (relative residual 10^{-2}–2×10^{-1}): the next speed-up is a better preconditioner.

9. Result on the production cube

$z(\tau{=}1)$ relative to the cube mean, median [5 %, 95 %] in km, by $|\mathbf{B}|$ at $\tau = 1$ (fixed-point = the superseded scheme):

$\lvert\mathbf{B}\rvert$ at $\tau = 1$	column HSE in τ	fixed-point scheme	**mhs-poisson-ltau**
0–200 G	−38 [−85, 187]	−55 [−257, 380]	−10 [−35, 73]
200–500 G	−68 [−88, 58]	+120 [−180, 549]	−26 [−52, 24]
500–1000 G	−86 [−104, −36]	+183 [−198, 587]	−51 [−89, −17]
1000–1500 G	−121 [−144, −92]	+314 [−117, 563]	−96 [−125, −63]
1500–3000 G	−156 [−175, −140]	+402 [58, 568]	−130 [−167, −104]

Force imbalance after the solve, RMS in units of ρg: 0.04–0.09 horizontal and 0.04–0.06 vertical for $\log\tau < 0$ (start 0.3–0.8).

10. Design history

1. Fixed point (`scripts/poisson/poisson_tau.jl`): map to a height grid, Poisson solve there, re-integrate the τ columns, repeat. Each pass is a full Poisson solve (1 h on the full cube), and it diverged where $\mathbf{L}$ carries the gas: the heights were integrated with a pressure that was then replaced. A corrected variant (heights integrated with the Poisson pressure) settled but lifted the strong-field columns instead of depressing them (median +400 km above the cube mean for 1500–3000 G).
2. Direct solve with $\mathbf{L}$ fixed in the Jacobian: exact without $\mathbf{B}$, stalled in magnetic regions.
3. Exact Jacobian + LM with whole-step scaling + the consecutive-step stop rule: the present mode.