© Yuri Balasanov, 2014

All Rights Reserved

No part of this lecture notes document or any of its contents may be reproduced, copied, modified or adapted without the prior written consent of the author, unless otherwise indicated for stand-alone materials.

The content of these lectures, any comments, opinions and materials are put together by the author especially for the course Statistical Analysis, they are sole responsibilities of the author, but not of the author's employers or clients.

The author cannot be held responsible for any material damage as a result of use of the materials presented in this document or in this course. For any inquiries contact the author, Yuri Balasanov, at ybalasan@uchicago.edu or yuri.balasanov@research-soft.com .

4日 4日 4日 4日 4日 日 り90

```
Probability Space.
   Every random experiment is associated with
   the object [II, F, P] called Probability Space.
   I - all elementary outcomes.
  F - all possible events related to elementary outcomes
   P - probability measure that assigns probability
        number to each event.
   Example. Jossing a fair coin once.
    D= 5"Heads", "Tails"}
   F= [Ø, "Heads", "Tails", I }
    P={Ø}=0; P{"Heads"}=P{"Tails"}===; P{I}=1.
    Event Ø includes all "impossible" events
    (the coin overcomes the force of gravity and
     goes into space).
    Event I means that either "Heads" or "Tails" occurs.
Balasanov (University of Chicago, MScA)
                         Math Challenge
                                           © Y. Balasanov, 2014
```

Coin tossed 2 times. What is the probability space? Let the outcomes of each tors be: "H"=0, "T"=1. The Probability space { , T, F, P} is:

SL={00,01,10,11}. (either both Heads or both Tails)

F={ Ø,00,01,10,11,00+01,00+10,00+11,01+10,01+11,10+11, 

00+01+10,00+01+11,01+10+11,00+01+10+11}

Balasanov (University of Chicago, MScA)

Math Challenge

```
A coin is tossed 2 times. Define random variables
                                 associated with this experiment
  \Omega = \{HH, HT, TH, TT\}.
 1. R.V. X: D→R is the number of Tails in the first toss.
  X = \begin{cases} \infty_0 = 0 : HH, HT, \\ \infty_1 = 1 : TH, TT. \end{cases}
2. R.V. Y: IL -> R is the total number of Tails
  Y= { 3, = 0 : HH
3, = 1 : HT, TH
3, = 2 : TT
   marginal distributions:
 ア{X=xo}=ア{X=xa}=立; ア{Y=yo}=ア{Y=yo}=ン: ア{Y=yo}=士: ア{Y=yo}=士:
 Joint distribution: (Events)
                       ア{X=x,,Y=z,}=ア{X=x,,Y=z,}=女,
ア{X=x,,Y=z,}=ア{X=x,,Y=z2}=女,
                              PIX=x, Y=y, Y=PIX=x, Y=yo = 0.
Balasanov (University of Chicago, MScA)
                                Math Challenge
                                                       © Y. Balasanov, 2014
```

