

Mathematics 4MB3/6MB3 Mathematical Biology
2018 ASSIGNMENT 1

Group Name: [πrates](#)

Group Members: [Adeyemi Fakorede](#), [Megan Hartwell](#), [Ahmad Mahmood](#), [Bradley Montgomery](#), [Courtney Mulholland](#)

1 Analysis of the SI model

The SI model can be written

$$\frac{dI}{dt} = \beta I(N - I), \quad (1)$$

where I denotes prevalence and $N = S + I$ is the total population size.

- (a) Prove that the endemic equilibrium (EE) is a globally asymptotically stable (GAS) equilibrium by finding an appropriate Lyapunov function. Note that “global” here refers to all biologically relevant initial conditions except the (unstable) disease free equilibrium (DFE).

Hint: Lyapunov functions often look paraboloidal.

Note: Notions of stability and Lyapunov functions were discussed in Math 3F03 Lecture 27 in 2013 (<http://www.math.mcmaster.ca/earn/3F03>).

NOTE: When referring to equation (1) make sure to use `equation~\eqref{E:SI}` in your source file so that you obtain a hyperlinked reference that is automatically numbered correctly.

Proof. An appropriate Lyapunov function to prove that the endemic equilibrium is a globally asymptotically stable equilibrium is given by

$$L = S^2 = (N - I)^2 > 0 \quad \forall X \in \mathcal{O} \setminus X_* \quad (2)$$

$$\dot{L} = 2S\dot{S} = 2(N - I)(-\dot{I}) = -2\beta I^2(N - I)^2 \leq 0 \quad \forall X \in \mathcal{O} \setminus X_* \quad (3)$$

which satisfies both of the conditions in Theorem 1 (Lyapunov’s Direct Method) stated in the section “Notes on Lyapunov functions”.

□

- (b) In class we proved only stability of the EE, not asymptotic stability. Prove GAS “directly” in two distinct ways:
- (i) find the exact solution of the model and take the limit as $t \rightarrow \infty$, and conclude that every solution that starts in the interval $(0, N)$ converges to the EE (this approach works only in situations where you can find the exact solution);

Proof. ...beautifully clear and concise text to be inserted here... Since an exact solution is requested, there will no doubt be an equation of the form

$$I(t) = \cdots \text{blah blah blah} \cdots, \quad (4)$$

which will be extremely enlightening. \square

- (ii) given $\epsilon > 0$, prove that for any $I(0) \in (0, N) \exists t < \infty$ such that $I(t) \in [N - \epsilon, N)$ and use this to establish GAS. (Do not use your exact solution in this part; the point is to use an approach that also works for models that cannot be solved exactly.)

Proof. ...beautifully clear and concise text to be inserted here... \square

2 Analysis of the basic SIR model

The basic SIR model is specified by the following system of differential equations.

$$\frac{dS}{dt} = -\mathcal{R}_0 SI \quad (5a)$$

$$\frac{dI}{dt} = \mathcal{R}_0 SI - I \quad (5b)$$

$$\frac{dR}{dt} = I \quad (5c)$$

The state variables S , I and R are the proportions of the population that are susceptible, infectious and removed, respectively. The parameter \mathcal{R}_0 is the basic reproduction number. The time unit has been chosen to be the mean infectious period for convenience.

- (a) A quantity of some practical importance is the **peak prevalence** of disease in the population, *i.e.*, the maximum proportion of the population that is simultaneously infected. Find an exact expression for the peak prevalence, given initial conditions (S_0, I_0) . Why might a public health official want to know this quantity?

Solution. ...beautifully clear and concise text to be inserted here... \square

- (b) It would be helpful to have an analytical expression for the solution of the model. Most valuable would be a formula for $I(t)$, which is most closely related to time series data. You probably will not find a formula for $I(t)$ (extra credit if you do!!) but it is definitely possible to find an exact expression that relates R (proportion removed) and t (time).
- (i) Find such an expression. *Hint:* Combine the equations for dS/dt and dR/dt into one equation that can be solved for S as a function of R . Then recall that $S + I + R = 1$ and use the dR/dt equation again. *Note:* You will end up with an expression for t as a function of R , not R as a function t .

Solution.

$$\frac{\frac{dS}{dt}}{\frac{dR}{dt}} = \frac{dS}{dR} = \frac{-\mathcal{R}_0 SI}{I} = -\mathcal{R}_0 S \implies S = S_0 e^{-\mathcal{R}_0 R} \quad (6)$$

Recall $S + I + R = 1$ so we have that

$$S_0 e^{-\mathcal{R}_0 R} + I + R = 1 \implies R = 1 - I - S_0 e^{-\mathcal{R}_0 R} \quad (7)$$

From equation \eqref{E:S}...

$$\frac{dR}{dt} = 1 - R - S_0 e^{-\mathcal{R}_0 R} \quad (8)$$

We can solve this as a separable equation to get the expression for $t(R)$

$$t = \int \frac{1}{1 - R - S_0 e^{-\mathcal{R}_0 R}} dR \quad (9)$$

□

- (ii) Use your expression for $t(R)$ to find an expression for the time at which peak prevalence will occur. Why might this be useful?

Solution. The peak prevalence corresponds to where the determinant of I is equal to zero:

$$\frac{dI}{dt} = \mathcal{R}_0 SI - I = 0 \implies I = 0 \quad \text{or} \quad \mathcal{R}_0 = \frac{1}{S} \quad (10)$$

Since $I = 0$ is trivial, take $\mathcal{R}_0 = 1/S$ and plug this into the expression for $t(R)$ from 2(b)(i) giving

$$t = \int \frac{1}{1 - R - S_0 e^{-R/S}} dR \quad (11)$$

This expression may be useful because the expression relates time to peak prevalence. This means that if one can solve this expression, they will know at what amount of time units peak prevalence will occur.

□

- (iii) How could your expressions be used to compare with the time series for pneumonia and influenza in Philadelphia in 1918? (Don't actually do it; just clearly explain your thinking including any assumptions you are making.) Would you advise your assistant who just graduated with a degree in math and biology to do this (to help you prepare your report for the public health agency)? Why or why not?

Solution. The P & I 1918 time series shows deaths as a function of time, whereas the model achieved in this solution is time as a function of those recovered; recovered including both those that died and those that are over the illness and now immune. If the $t(R)$ model from this question was solved explicitly (assuming we could find a solution) and an inverse $t^{-1}(R)$ was solved for explicitly, (one again assuming this exists and can be solved) we would have a model similar to the P & I 1918 model which models recovery as a function of time. One further assumption needed is that all of those belonging to the recovered category in the $t^{-1}(R)$ are dead, since alive and immune is not included in the time series data for P & I 1918. I would not advise my assistant to help prepare a report for the public health agency with this model since the assumptions are too extreme and unreasonable. A better model that has an explicit equation and takes into account those recovered that are alive will do a much better job modelling the P & I 1918 data than the one achieved in this question.



□

- (iv) Is it possible to find an exact analytical expression for t as a function S ?

Solution. Yes it is possible to find an exact analytical expression for $t(S)$. The same steps from part (i) can be taken, except R is solved for as a function of S (separable equation) when combining dS/dt and dR/dt , and the dS/dt equation is solved for I to achieve a relation between dS/dt and I to substitute into the $S + I + R = 1$ equation. Carrying on as in part (i) R and I can now be replaced in $S + I + R = 1$ and the equation can be solved as a separable equation, giving t as a function of S or $t(S)$.

□

- (c) Prove that all solutions of the basic SIR model approach $I = 0$ asymptotically, and explain why this makes biological sense. *Hint:* Is the function $L(S, I) = I$ a Lyapunov function? Read the [Notes on Lyapunov functions](#) below.

Answers. ... beautifully clear and concise text to be inserted here... Since no numerical analysis or graphics were requested, there is no need to use  to answer these questions. Note, incidentally, that the `\Rlogo` command that produces  simply includes the image file `images/Rlogo.pdf` and scales it appropriately for text within a paragraph. If we really loved that logo and wanted to display as we would a figure, we easily could... □



(d) Find and classify the stability of all equilibria of the basic SIR model.

Solution. ... beautifully clear and concise text to be inserted here. ...



Notes on Lyapunov functions

Consider Lyapunov's Stability Theorem as stated in [Math 3F03 Lecture 28 in 2013](#):

Theorem 1 (Lyapunov's Direct Method). *Consider an equilibrium X_* of $X' = F(X)$ and an open set \mathcal{O} containing X_* . If \exists a differentiable function $L : \mathcal{O} \rightarrow \mathbb{R}$ such that*

$$(a) \ L(X_*) = 0 \quad \text{and} \quad L(X) > 0 \quad \forall X \in \mathcal{O} \setminus \{X_*\} \quad (L \text{ positive definite on } \mathcal{O})$$

$$(b) \ \dot{L}(X) \leq 0 \quad \forall X \in \mathcal{O} \setminus \{X_*\} \quad (\dot{L} \text{ negative semi-definite on } \mathcal{O})$$

then X_ is stable and L is called a **Lyapunov function**. If, in addition,*

$$(c) \ \dot{L}(X) < 0 \quad \forall X \in \mathcal{O} \setminus \{X_*\} \quad (\dot{L} \text{ negative definite on } \mathcal{O})$$

then X_ is asymptotically stable and L is called a **strict Lyapunov function**.*

Theorem 1 can be generalized for analysis of stability of sets more complicated than isolated equilibria, such as periodic orbits or line segments. If you think through the proof of the theorem above (*e.g.*, [?, §9.2, theorem stated on p. 193 and proved on p. 196]), you should be able to convince yourself that the proof still works if the equilibrium X_* is replaced by any *closed forward-invariant set* (often simply called a *closed invariant set*). This observation allows us to state the following more general theorem.

Theorem 2 (Lyapunov's Direct Method for Closed Invariant Sets). *Consider a closed invariant set \mathcal{C} of $X' = F(X)$ and an open set \mathcal{O} containing \mathcal{C} . If \exists a differentiable function $L : \mathcal{O} \rightarrow \mathbb{R}$ such that*

$$(a) \ L(X) = 0 \quad \forall X \in \mathcal{C} \quad \text{and} \quad L(X) > 0 \quad \forall X \in \mathcal{O} \setminus \mathcal{C} \quad (L \text{ positive definite on } \mathcal{O})$$

$$(b) \ \dot{L}(X) \leq 0 \quad \forall X \in \mathcal{O} \setminus \mathcal{C} \quad (\dot{L} \text{ negative semi-definite on } \mathcal{O})$$

*then \mathcal{C} is stable and L is called a **Lyapunov function**. If, in addition,*

$$(c) \ \dot{L}(X) < 0 \quad \forall X \in \mathcal{O} \setminus \mathcal{C} \quad (\dot{L} \text{ negative definite on } \mathcal{O})$$

*then \mathcal{C} is asymptotically stable and L is called a **strict Lyapunov function**.*

Note in the above theorems that open sets are defined relative to the subset of interest; in our case this subset is $\Delta = \{(S, I) : S \geq 0, I \geq 0, S + I \leq 1\}$, not all of \mathbb{R}^2 . An open set of Δ is a set of the form $U \cap \Delta$ where U is an open set of \mathbb{R}^2 . (These sets are said to be open in the **relative topology** on Δ .) In particular, note that Δ is *open* as a subset of itself, in spite of the fact that it is *not open* as a subset of \mathbb{R}^2 , whereas Δ is closed in both the relative topology on Δ and the usual topology on \mathbb{R}^2 .

— END OF ASSIGNMENT —

Compile time for this document: January 20, 2018 @ 14:59