

Prefecture-Level Older-Adult Solo Household Rate and Large-Volume Infusion Therapy Utilization in Japan: An Ecological Study

Haruki Saito*¹, Tetsuya Ohira^{1,2}

1. Department of Epidemiology, Fukushima Medical University School of Medicine,
Fukushima, Japan

2. Radiation Medical Science Center for the Fukushima Health Management Survey,
Fukushima Medical University, Fukushima, Japan

*Corresponding author. Email: m211039@fmu.ac.jp

Abstract

Background

Older adults living alone are often considered vulnerable to heat because of reduced informal monitoring, delayed help-seeking, and barriers to cooling. Prefecture-level indicators such as elderly solo household rate have been proposed as routinely available markers of this vulnerability, but such indicators may also reflect a prefecture's underlying population age structure. We examined the association between elderly solo household rate and large-volume infusion therapy utilization, and tested whether this association remained after accounting for population age structure, healthcare supply, and summer heat exposure.

Methods

We conducted an ecological study of 47 Japanese prefectures using fiscal year 2023 data. Large-volume infusion therapy utilization (procedure code G004, ≥ 500 mL) from the National Database (NDB) Open Data was used as the outcome, expressed per 100,000 population. Elderly solo household rate, defined as the percentage of all households consisting of one person aged ≥ 65 years, was derived from the 2020 National Census and used as the primary exposure. We fit cumulative regression models adding, in turn, prefectural ageing rate, general hospital beds per 100,000 population, and official daily-maximum wet-bulb globe temperature (WBGT) days ($\geq 28^\circ\text{C}$, June–September 2023). Nonlinearity, metropolitan influence, and an alternative exposure using the older population as the denominator were also examined.

Results

Elderly solo household rate was associated with infusion therapy utilization in the unadjusted analysis ($\beta = 656.2$; 95% CI, 326.9–985.6; $p < 0.001$). Adding prefectural ageing rate attenuated the estimate by 53.3% ($\beta = 306.1$; 95% CI, –71.3–683.5; $p = 0.112$), and the association was no longer statistically significant. Further adjustment for healthcare supply and WBGT reduced the estimate toward zero ($\beta = -9.4$; 95% CI, –493.9–475.1; $p = 0.970$).

Conclusions

The prefectural elderly solo household rate was associated with large-volume infusion therapy utilization in the unadjusted analysis. However, the estimate was substantially attenuated and was no longer statistically significant after adjustment for population age structure. Further adjustment for healthcare supply and summer heat exposure reduced the estimate toward zero. These findings do not support an independent ecological association between living-alone indicators and infusion therapy utilization and highlight the need to account for population structure when evaluating social vulnerability in heat-health research.

Keywords: heat-health surveillance; elderly solo household rate; population ageing; healthcare utilization; ecological study; Japan

Introduction

Heat-related illness is an increasingly urgent public health challenge in Japan. In 2023, heatstroke caused 1,580 deaths, and emergency transports for suspected heatstroke reached 91,467 cases (Fire and Disaster Management Agency 2023;

MHLW 2024). Climate change, population ageing, and urbanization are expected to increase this burden (IPCC 2021; Günsche et al. 2026).

Heat-health responses have therefore emphasized meteorological surveillance. Temperature, humidity, and WBGT are established proximal determinants of heatstroke (Parsons 2006; Blazejczyk et al. 2012), and Japan has implemented WBGT forecasting and heat alert systems (Ministry of the Environment Japan n.d.). These systems are essential, but they primarily describe environmental hazard and do not by themselves identify which populations may be least able to receive warnings, act on them, or obtain timely assistance.

Older adults living alone are often considered vulnerable to heat because of reduced informal household monitoring (for example, a family member or neighbor noticing signs of heat illness or prompting fluid intake and use of cooling), economic barriers to cooling use, delayed help-seeking, and reduced access to heat-health information, in addition to physiological vulnerability from reduced sweating capacity, diminished thirst perception, impaired thermoregulation, and chronic disease (Kenney and Munce 2003; Bouchama and Knochel 2002; Vandentorren et al. 2006; Canouï-Poitrine et al. 2006; MIC 2015; Farbotko and Waitt 2011; Semenza et al. 1996; Naughton et al. 2002; Oberai et al. 2025; Zhang et al. 2025). Prefecture-level elderly solo household rate, the percentage of all households consisting of one older adult living alone, has been proposed as a routinely available marker of this social vulnerability. However, because such a measure is calculated over all households, it may also reflect a prefecture's overall population age structure rather than a distinct social characteristic, and this potential confounding has not been formally examined in prior ecological work.

Japanese studies have examined individual risk factors and ambulance transport data (Ng et al. 2014; Miyatake et al. 2012), but nationwide ecological evidence linking

elderly solo household rate to large-volume infusion therapy utilization, and on whether such an association is independent of population age structure, healthcare supply, and heat exposure, is limited. We therefore examined whether elderly solo household rate was associated with large-volume infusion therapy utilization, and the extent to which this association remained after accounting for prefectural ageing rate, healthcare supply, and official heat-exposure measures.

Materials and methods

1. Study Design

We conducted a nationwide ecological study using Japanese prefectures (N = 47) as the unit of analysis; the prefectures ranged in total 2023 population from approximately 0.5 million to 14.1 million, together representing Japan's total population of approximately 124.4 million. Prefecture-level aggregate data were used because individual household and geospatial information are not available from NDB Open Data. The study period was fiscal year 2023 (April 2023 through March 2024); heat exposure was measured using official Ministry of the Environment wet-bulb globe temperature (WBGT) monitoring-site observations during the summer season (June–September 2023), an unusually hot summer at the time of data collection (Japan Meteorological Agency 2023).

2. Outcome: Large-Volume Infusion Therapy Utilization

Infusion therapy utilization rates were derived from the 10th NDB Open Data. We extracted claims for infusion procedures involving ≥ 500 mL volume (procedure code G004), used for volume repletion in a range of clinical contexts including, but not

limited to, heat-related dehydration; the code is not diagnosis-specific. We describe the outcome as large-volume infusion therapy utilization, a non-specific healthcare-utilization measure that may include a heat-related dehydration component. Rates were calculated per 100,000 population using the 2023 population estimate, temporally aligned with the FY2023 NDB claims period (a sensitivity analysis using the 2020 Census population denominator is reported in Online Resource 1). We independently reviewed the complete public file structure of the 10th NDB Open Data and confirmed that prefecture-by-age cross-tabulated counts for G004 are not publicly available; sex/age and prefecture counts exist only as separate marginal tables. We therefore could not construct an age-specific or age-standardized outcome and instead adjusted for prefectural ageing rate directly (Section 5). As a construct validity check, we examined the cross-prefectural correlation between infusion therapy utilization and the summer 2023 heatstroke ambulance transport rate from the Fire and Disaster Management Agency (FDMA) daily transport report (Fire and Disaster Management Agency 2023); this provides limited support for the presence of a heat-related component in the outcome rather than validating it as a direct heat-illness measure.

3. Exposure: Elderly Solo Household Rate

Elderly solo household rate was defined as the percentage of all households consisting of one person aged ≥ 65 years, derived from the 2020 National Census. We interpret this measure as a prefecture-level indicator of the burden of older adults living alone relative to all households, rather than a direct individual-level measure of social isolation. As a complementary sensitivity analysis, we also constructed an alternative exposure using the prefecture's population aged ≥ 65 years as the denominator (percentage of the 65+ population living alone; Section 9; Online Resource 1).

4. Climatic Covariates

Heat exposure was assessed using official wet-bulb globe temperature (WBGT) observations from the Ministry of the Environment's Heat Illness Prevention Information monitoring network for June–September 2023. For each monitoring site, we counted the number of days on which daily maximum WBGT met or exceeded each threshold, and then averaged these site-specific day-counts within each prefecture (mean 17.9 sites per prefecture; range 5–163). The primary heat metric used in the revised analysis was the prefecture-mean number of days with daily maximum WBGT $\geq 28^{\circ}\text{C}$; days with WBGT $\geq 31^{\circ}\text{C}$, days with WBGT $\geq 33^{\circ}\text{C}$, and cumulative WBGT excess above 28°C were examined as sensitivity heat metrics (Section 6; Online Resource 1). For descriptive purposes (Table 1), we also report the number of days with daily maximum temperature $\geq 35^{\circ}\text{C}$, averaged across Japan Meteorological Agency stations within each prefecture; this variable was not used as an adjustment variable in the primary models.

5. Prefectural Ageing Rate and Healthcare Supply

Prefectural ageing rate, defined as the percentage of the population aged ≥ 65 years and calculated from the Statistics Bureau of Japan's population estimate as of 1 October 2023, was included as the primary demographic adjustment variable, temporally aligned with the outcome denominator. General hospital beds per 100,000 population (2023 Survey of Medical Institutions, Ministry of Health, Labour and Welfare) were included as the primary healthcare-supply adjustment variable, with general clinics per 100,000 population (same source and year) examined as a sensitivity analysis substituted for beds. Air conditioning prevalence (proportion of households owning at least one air conditioning unit, %), obtained from the 2014 National Survey of Family Income and Expenditure, is retained for descriptive

purposes (Table 1) but was not included in the primary adjusted models, because these data describe ownership rather than use.

6. Statistical Analysis

All models used the originally specified elderly solo household rate as the exposure. We fit four cumulative models: the exposure alone (Model O-A); Model O-A plus prefectural ageing rate (Model O-B); Model O-B plus general hospital beds per 100,000 (Model O-C); and Model O-C plus days with WBGT $\geq 28^{\circ}\text{C}$ (Model O-D). We did not use multicollinearity as a reason to avoid adjustment; variance inflation factors for all covariates in all models were below 3.1 (Table 2). For every model we report the unstandardized coefficient with heteroskedasticity-consistent (HC3) robust 95% confidence intervals and p-value, the fully standardized coefficient, adjusted R^2 , the partial R^2 for the exposure (based on the reduction in ordinary least-squares residual sum of squares, independent of the standard-error estimator), and the corrected Akaike information criterion (AICc).

We conducted the following sensitivity analyses on the fully adjusted model: substituting general clinics for hospital beds; excluding Tokyo, Osaka, and Kanagawa jointly; leave-one-prefecture-out re-estimation (47 iterations); and separately substituting the primary heat metric with WBGT $\geq 31^{\circ}\text{C}$ days, WBGT $\geq 33^{\circ}\text{C}$ days, and cumulative WBGT excess, applying Benjamini-Hochberg false-discovery-rate correction across this exploratory heat-metric set. As a further sensitivity analysis, we repeated all models using the 2020 Census population denominator for the outcome (Online Resource 1). We assessed nonlinearity by comparing the linear model with a centered quadratic exposure term and a natural cubic spline (3 degrees of freedom), using partial F-tests and AICc; a median-split stratified analysis is reported in Online Resource 1 and did not support a simple dose-response interpretation. We also

examined an alternative exposure using the older population as the denominator (Section 9).

Regression coefficients (β), 95% confidence intervals (CIs), R^2 , adjusted R^2 , and p-values are reported, together with regression diagnostics (residuals, leverage, Cook's distance, DFBETAs) for the fully adjusted model. Statistical significance was defined as two-sided $p < 0.05$, interpreted alongside effect estimates and confidence intervals rather than significance alone. Analyses were performed using Python (version 3.14), statsmodels (version 0.14), scikit-learn, and matplotlib.

This study used publicly available aggregate data; individual informed consent was not required, and institutional ethics review was not applicable in accordance with Japanese ethical guidelines for epidemiological research.

Results

1. Descriptive Statistics

Across the 47 Japanese prefectures, mean elderly solo household rate was $12.6 \pm 1.9\%$ (range 9.4–17.8%; Table 1). Mean large-volume infusion therapy utilization was $8,022 \pm 2,389$ per 100,000 population (range 4,231–13,234). Mean prefectural ageing rate was $31.6 \pm 3.3\%$ (range 22.8–39.0%). Mean general hospital beds were 794 ± 149 per 100,000 (range 509–1,147). Mean number of days with daily maximum WBGT $\geq 28^\circ\text{C}$ (official Ministry of the Environment data) was 72.9 ± 14.1 (range 27.3–112.3); for descriptive comparison, mean number of days with maximum temperature $\geq 35^\circ\text{C}$ was 17.9 ± 12.1 (range 0.0–44.0). Mean air conditioning prevalence was $89.7 \pm 13.6\%$ (range 26.6–98.2%).

2. Association Between Elderly Solo Household Rate and Infusion Therapy Utilization: Unadjusted and Adjusted Models

Elderly solo household rate was associated with infusion therapy utilization in the unadjusted model (Model O-A: $\beta = 656.2$; 95% CI, 326.9–985.6; standardized $\beta = 0.521$; $R^2 = 0.272$; adjusted $R^2 = 0.256$; $p < 0.001$; Table 2, Fig. 1). Adding prefectural ageing rate (Model O-B) changed the coefficient to 306.1 (95% CI, –71.3–683.5; standardized $\beta = 0.243$; $p = 0.112$), a descriptive attenuation of 53.3%; the 95% confidence interval crossed zero and statistical significance was lost, although the interval remains compatible with both a modest positive association and little or no association. Adding general hospital beds per 100,000 (Model O-C) reduced the coefficient further ($\beta = 94.1$; 95% CI, –302.4–490.6; $p = 0.642$); the estimate was similarly attenuated when clinics were substituted for beds (Online Resource 1). Adding WBGT $\geq 28^\circ\text{C}$ days (Model O-D) reduced the coefficient to near zero and reversed its sign ($\beta = -9.4$; 95% CI, –493.9–475.1; $p = 0.970$). Maximum VIF across all models was 3.01 (Table 2).

3. Influence Diagnostics and Sensitivity Analyses

Cook's distance for the fully adjusted model identified Hokkaido as the most influential prefecture (Cook's D = 0.667, exceeding the $4/N = 0.085$ threshold by approximately 7.8-fold), followed by Kochi (0.190) and Hiroshima (0.144); Hokkaido's WBGT summary averages 163 monitoring sites spanning substantial internal climatic heterogeneity. Leave-one-prefecture-out re-estimation of the fully adjusted model (47 iterations) produced coefficients ranging from –107.9 to 92.8, with the 95% CI crossing zero in every iteration, indicating that the attenuated result was not driven by any single prefecture, including Hokkaido. Excluding Tokyo, Osaka, and Kanagawa

jointly did not materially change the result ($\beta = -42.0$; 95% CI, -645.6 – 561.6 ; $p = 0.891$).

4. Nonlinearity

A quadratic term for elderly solo household rate did not improve model fit relative to the linear fully adjusted model (partial F test $p = 0.142$; AICc 845.6 versus 845.9), nor did a natural cubic spline (3 degrees of freedom; $p = 0.141$; AICc 845.9). A median-split stratified analysis, in which the association was present in the lower stratum but absent in the higher stratum, is reported in Online Resource 1; it did not support a simple dose-response interpretation and is not used as a primary result.

5. Alternative Exposure (Denominator) Analysis

As a complementary sensitivity analysis, we examined an alternative exposure defined as the percentage of the prefecture's population aged ≥ 65 years who live alone (denominator: 65+ population, rather than all households). This alternative measure showed no association with infusion utilization, either unadjusted (standardized $\beta = 0.0003$; 95% CI, -0.327 – 0.328 ; $p = 0.998$) or in the analogous fully adjusted model (standardized $\beta = -0.034$; 95% CI, -0.298 – 0.231 ; $p = 0.803$), and this pattern was unchanged across the sensitivity analyses described above (Online Resource 1).

6. Construct Validity: FDMA Heatstroke Transport Validation

As a construct validity check, infusion therapy utilization was examined in relation to the summer 2023 heatstroke ambulance transport rate from FDMA (Fire and Disaster Management Agency 2023). A positive association was observed ($\beta = 70.19$; 95% CI, 36.85 – 103.53 ; $r = 0.534$; $p < 0.001$; Online Resource 1), which provides limited

support for the presence of a heat-related component in the infusion therapy outcome, rather than validating it as a direct measure of heat-related dehydration.

7. Comparison-Outcome Analysis

A comparison with general outpatient utilization (R5 Patient Survey) did not support outcome specificity for infusion therapy and is reported in the supplementary material (Online Resource 1), using standardized coefficients only; the previously reported raw-coefficient comparison is not valid because the two outcomes are measured on different scales and has been removed.

Discussion

This nationwide ecological analysis examined whether elderly solo household rate, a routinely available prefecture-level indicator, was associated with large-volume infusion therapy utilization, and whether any such association was independent of population age structure, healthcare supply, and summer heat exposure. An association was present in the unadjusted analysis but was substantially attenuated (53.3%) and lost statistical significance after adjustment for prefectural ageing rate alone, and was further attenuated toward zero after additional adjustment for healthcare supply and heat exposure. An alternative exposure using the older population as the denominator showed no association with the outcome at any stage. These findings do not support an independent ecological association between elderly solo household rate and infusion therapy utilization.

Physiological vulnerability to heat among older adults, including reduced thermoregulation and blunted thirst perception, remains biologically plausible and is well documented (Kenney and Munce 2003; Bouchama and Knochel 2002; Bach et al.

2024; Blatteis 2012). However, the present ecological results indicate that a prefecture-level living-alone indicator does not, by itself, capture an independent surveillance-relevant signal once population age structure is taken into account.

Regional infusion use may also reflect healthcare supply and local treatment practice in addition to underlying morbidity: the estimate was further attenuated after adjusting for general hospital beds, and this pattern was unchanged when clinics were substituted for beds. We did not test whether supply-induced demand specifically explains infusion use; we show only that accounting for healthcare supply further attenuates the association between elderly solo household rate and infusion therapy utilization.

Our findings do not contradict international evidence that social isolation is associated with heat-related mortality at the individual and small-area level, including during the 1995 Chicago and 2003 European heat waves (Vandentorren et al. 2006; Canoui-Poitaine et al. 2006; Semenza et al. 1996; Conti et al. 2005). Ecological associations and individual-level associations address different questions, and our attenuated ecological finding at the prefecture level should not be read as evidence against individual-level effects of social isolation on heat vulnerability, which this design cannot test.

The absence of a robust independent association for elderly solo household rate should not be interpreted as evidence that heat exposure is unimportant. Alternative WBGT metrics did not materially alter the interpretation; detailed exploratory results are reported in Online Resource 1. Prefecture-level climatic averages may also miss neighborhood and indoor exposures, and regional heat adaptation may attenuate simple ecological climate-health associations (Hondula et al. 2017; Harlan et al. 2013; Oka et al. 2023).

These findings suggest that prefecture-level social-vulnerability indicators such as elderly solo household rate should not be interpreted as independent heat-vulnerability markers without accounting for population age structure, healthcare supply, and improved heat-exposure measures. This is consistent with, rather than a departure from, the original rationale for examining social indicators in heat-health surveillance: it indicates that such indicators require careful demographic validation before being proposed for that purpose.

Strengths include nationwide coverage of all 47 prefectures, use of NDB Open Data, direct re-examination of the originally specified exposure and outcome, replacement of the original simplified WBGT approximation with official daily-maximum WBGT observations, and a comprehensive set of sensitivity, influence, and nonlinearity analyses.

Several limitations should be considered. First, this is an ecological design; associations at the prefecture level may not reflect individual-level relationships, and we cannot determine whether individuals living alone within a prefecture had higher infusion therapy utilization than those living with family. Second, elderly solo household rate is a household-composition measure and is not equivalent to social isolation. Third, infusion therapy utilization (G004) is not specific to dehydration or heat illness and may be used for volume repletion in other contexts, including perioperative fluid management, gastrointestinal illness, infections, and general supportive care; monthly aggregation of claims is not available in prefecture-level NDB Open Data, so a summer peak in utilization could not be verified directly.

Fourth, prefecture-by-age cross-tabulated G004 counts are not publicly available in the 10th NDB Open Data, precluding an age-specific or age-standardized outcome; we addressed this by adjusting for prefectural ageing rate directly. Fifth, the analysis covers a single fiscal year during an unusually hot summer (2023; Japan

Meteorological Agency 2023), limiting generalizability to other years. Sixth, with N = 47, adjusted confidence intervals remain wide, and this analysis cannot precisely distinguish a small residual association from no association. Seventh, unmeasured confounding cannot be excluded, including reduced thirst perception and lower fluid intake among older adults, chronic disease prevalence, medication use, infections, gastrointestinal illness, and other individual clinical factors associated with infusion therapy that are not captured at the prefecture level.

Eighth, climatic measures are prefecture-level averages and do not capture indoor or neighborhood exposures; the WBGT summary for Hokkaido in particular averages 163 monitoring sites across substantial internal climatic heterogeneity and was the most influential single observation in regression diagnostics, although the qualitative conclusion was unchanged when Hokkaido was excluded (leave-one-prefecture-out analysis). Ninth, source years differ across variables (2020 Census for elderly solo household rate; 2023 estimates for population, ageing rate, and outcome denominator; 2023 Survey of Medical Institutions for healthcare supply; 2014 survey for air conditioning prevalence), and we aligned the outcome denominator to 2023 to match the NDB claims period while retaining a 2020 Census-based denominator analysis as a sensitivity check (Online Resource 1).

Finally, air conditioning prevalence (2014 survey) reflects ownership, not actual use, and cannot establish whether cooling was applied; it is reported descriptively (Table 1) and was not included in the primary adjusted models. The present findings should be interpreted as evidence for cautious, demographically informed use of social-vulnerability indicators in heat-health surveillance, rather than as evidence against considering social factors altogether.

Given the ecological, cross-sectional, and single-year design, these findings should be considered hypothesis-generating. These considerations indicate that firmer

conclusions about an independent heat-vulnerability effect of household living arrangement would require age-specific outcome data, multi-year designs, and finer-grained exposure measurement, in addition to the demographic caution highlighted above.

Conclusions

In this nationwide ecological study, elderly solo household rate was associated with large-volume infusion therapy utilization in unadjusted analysis. However, this association was substantially attenuated and no longer statistically significant after adjustment for prefectural ageing rate, and was further attenuated toward zero after additional adjustment for healthcare supply and summer heat exposure.

These findings do not support elderly solo household rate as an independent ecological correlate of infusion therapy utilization once population age structure, healthcare supply, and heat exposure are taken into account. Given the ecological, cross-sectional, and single-year design, these findings should be regarded as hypothesis-generating. They highlight the need for careful demographic validation, including explicit adjustment for age structure and attention to denominator choice, before routinely available social-vulnerability indicators are proposed for heat-health surveillance.

Future individual-level and longitudinal studies, and age-specific administrative data where available, are needed to clarify whether living alone carries an independent heat-vulnerability signal beyond population age structure, and to establish the role of healthcare supply and improved heat-exposure measures in explaining prefecture-level utilization patterns.

381

382 Acknowledgments

383 The authors thank the Ministry of Health, Labour and Welfare of Japan for providing
384 access to the NDB Open Data. All data used are publicly available aggregate
385 statistics.

386

387 Conflicts of Interest

388 The authors declare no conflicts of interest.

389

390 Data Availability

391 The NDB Open Data used in this analysis are publicly available from the Ministry of
392 Health, Labour and Welfare of Japan
393 (<https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000177182.html>). The analysis
394 code, derived data, and provenance records are openly available on GitHub
395 (<https://github.com/haruki00430/NDB-solo-household-infusion-therapy-japan>) and
396 archived on Zenodo (<https://doi.org/10.5281/zenodo.20740375>), which always
397 resolves to the most recent version.

398

Author Contributions

Haruki Saito: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Visualization, Writing – original draft, Writing – review and editing.

Tetsuya Ohira: Conceptualization, Supervision, Writing – review and editing.

Funding

None declared.

Declaration of Generative AI and AI-Assisted Technologies in the Manuscript Preparation Process

During the preparation of the original submission, the authors used AI-assisted tools to support manuscript drafting and statistical analysis scripting. Cursor 3.0 (Anysphere) and Google Antigravity (Google) were used for AI-assisted writing and Python code development. Large language models used through these platforms included Claude Sonnet 4.6 and Claude Opus 4.8 (Anthropic), GPT-5.5 (OpenAI), and Gemini 3 Pro (Google). During preparation of the revision, the authors used AI-assisted tools (Claude, Anthropic, Claude 5/Sonnet family, accessed via Claude Code) to support text drafting and the development of data-processing and statistical-analysis code. The authors executed the code, verified all source data, numerical results, tables, and figures, and independently determined the statistical

methods and interpretation. No generative image tools were used. The authors reviewed and edited all outputs and take full responsibility for the final manuscript. AI was not listed as an author.

References

- Bach AJE, Cunningham SJK, Morris NR, Xu Z, Rutherford S, Binnewies S, et al (2024) Experimental research in environmentally induced hyperthermic older persons: a systematic quantitative literature review mapping the available evidence. *Temperature (Austin)* 11:4-26. <https://doi.org/10.1080/23328940.2023.2242062>
- Blatteis CM (2012) Age-dependent changes in temperature regulation - a mini review. *Gerontology* 58:289-295. <https://doi.org/10.1159/000333148>
- Blazejczyk K, Epstein Y, Jendritzky G, Staiger H, Tinz B (2012) Comparison of UTCI to selected thermal indices. *Int J Biometeorol* 56:515-535. <https://doi.org/10.1007/s00484-011-0453-2>
- Bouchama A, Knochel JP (2002) Heat stroke. *N Engl J Med* 346:1978-1988. <https://doi.org/10.1056/NEJMra011089>
- Canouï-Poitrine F, Cadot E, Spira A, Groupe Régional Canicule (2006) Excess deaths during the August 2003 heat wave in Paris, France. *Rev Epidemiol Sante Publique* 54:127-135. [https://doi.org/10.1016/S0398-7620\(06\)76706-2](https://doi.org/10.1016/S0398-7620(06)76706-2)
- Conti S, Meli P, Minelli G, et al (2005) Epidemiologic study of mortality during the Summer 2003 heat wave in Italy. *Environ Res* 98:390-399. <https://doi.org/10.1016/j.envres.2004.10.009>

442 Farbotko C, Waitt G (2011) Residential air-conditioning and climate change: voices of
443 the vulnerable. *Health Promot J Austr* 22:13-15. <https://doi.org/10.1071/HE11413>

444 Fire and Disaster Management Agency (2023) Daily number of patients with
445 heatstroke transported by ambulance in Japan. Tokyo: FDMA.
446 <https://www.fdma.go.jp/disaster/heatstroke/post4.html>

447 Günsche S, Borg MA, Anikeeva O, Varghese BM, Li Y, Bhandari D, et al (2026)
448 Mortality, morbidity and healthcare costs of short-term high temperatures and
449 heatwaves exposure in older populations: a global systematic review and meta-
450 analysis. *Environ Int* 208:110129. <https://doi.org/10.1016/j.envint.2026.110129>

451 Harlan SL, DeClet-Barreto JH, Stefanov WL, Petitti DB (2013) Neighborhood effects on
452 heat deaths: social and environmental predictors of vulnerability in Maricopa County,
453 Arizona. *Environ Health Perspect* 121:197-204. <https://doi.org/10.1289/ehp.1104625>

454 Hondula DM, Balling RC, Andrade R, Krayenhoff ES, Middel A, Urban A, Georgescu M,
455 Sailor DJ (2017) Biometeorology for cities. *Int J Biometeorol* 61(Suppl 1):59-69.
456 <https://doi.org/10.1007/s00484-017-1412-3>

457 Intergovernmental Panel on Climate Change (IPCC) (2021) Climate Change 2021: The
458 Physical Science Basis. Contribution of Working Group I to the Sixth Assessment
459 Report. Cambridge University Press, Cambridge.
460 <https://www.ipcc.ch/report/ar6/wg1/>

461 Japan Meteorological Agency (2023) Climate characteristics of summer 2023 (June–
462 August): the warmest summer since observations began in 1898 (mean temperature
463 anomaly +1.76°C, exceeding the previous record of +1.08°C set in 2010). Tokyo: JMA.
464 <https://www.jma.go.jp/jma/press/2309/01b/tenko230608.html>

465 Kenney WL, Munce TA (2003) Invited review: aging and human temperature
 466 regulation. *J Appl Physiol* 95:2598-2603.
 467 <https://doi.org/10.1152/japplphysiol.00202.2003>

468 Ministry of Health, Labour and Welfare (MHLW) (2024) Vital Statistics: Annual status
 469 of deaths due to heatstroke in Japan. Tokyo: MHLW.
 470 <https://www.mhlw.go.jp/toukei/saikin/hw/jinkou/tokusyu/necchusho24/index.html>

471 Ministry of Internal Affairs and Communications (MIC) (2015) National Survey of
 472 Family Income and Expenditure 2014. Tokyo: MIC.

473 Ministry of the Environment, Government of Japan (n.d.) Heat Illness Prevention
 474 Information: Heat Stress Index (WBGT) and Heat Stroke Alerts.
 475 <https://www.wbgt.env.go.jp/en/>

476 Miyatake N, Sakano N, Murakami S (2012) The relation between ambulance
 477 transports stratified by heat stroke and air temperature in all 47 prefectures of Japan
 478 in August, 2009: Ecological study. *Environ Health Prev Med* 17:77-80.
 479 <https://doi.org/10.1007/s12199-011-0221-2>

480 Naughton MP, Henderson A, Mirabelli MC, et al (2002) Heat-related mortality during a
 481 1999 heat wave in Chicago. *Am J Prev Med* 22:221-227.
 482 [https://doi.org/10.1016/S0749-3797\(02\)00421-X](https://doi.org/10.1016/S0749-3797(02)00421-X)

483 Ng CF, Ueda K, Takeuchi A, et al (2014) Sociogeographic variation in the effects of
 484 heat and cold on daily mortality in Japan. *J Epidemiol* 24:15-24.
 485 <https://doi.org/10.2188/jea.JE20130051>

486 Oberai M, Xu Z, Bach A, Forbes C, Jackman E, O'Connor F, et al (2025) A digital heat
 487 early warning system for older adults. *NPJ Digit Med* 8:114.
 488 <https://doi.org/10.1038/s41746-025-01505-5>

489 Oka K, Honda Y, Hui Phung VL, Hijioka Y (2023) Potential effect of heat adaptation on
490 association between number of heatstroke patients transported by ambulance and
491 wet bulb globe temperature in Japan. Environ Res 216:114666.
492 <https://doi.org/10.1016/j.envres.2022.114666>

493 Parsons K (2006) Heat stress standard ISO 7243 and its global application. Ind Health
494 44:368-379. <https://doi.org/10.2486/indhealth.44.368>

495 Semenza JC, Rubin CH, Falter KH, et al (1996) Heat-related deaths during the July
496 1995 heat wave in Chicago. N Engl J Med 335:84-90.
497 <https://doi.org/10.1056/NEJM199607113350203>

498 Vandentorren S, Bretin P, Zeghnoun A, et al (2006) August 2003 heat wave in France:
499 risk factors for death of elderly people living at home. Eur J Public Health 16:583-591.
500 <https://doi.org/10.1093/eurpub/ckl063>

501 Zhang C, Mohamad E, Azlan AA, Wu A, Ma Y, Qi Y (2025) Social media and eHealth
502 literacy among older adults: systematic literature review. J Med Internet Res
503 27:e66058. <https://doi.org/10.2196/66058>

504 **Tables and Figures**

505 **Table 1. Descriptive Statistics of Prefecture-Level Variables (N = 47)**

Variable	Unit	N	Mean	SD	Min	Median	Max
Outcome variable							
Large-volume infusion therapy utilization (G004, ≥500 mL)	per 100,000	47	8,022.2	2,389.3	4,231.3	7,608.7	13,234.1
Exposure variable							
Elderly solo household rate	%	47	12.6	1.9	9.4	12.3	17.8

Variable	Unit	N	Mean	SD	Min	Median	Max
Alternative exposure (sensitivity) variable							
% of population aged ≥65 living alone	%	47	17.7	3.1	12.1	17.1	26.1
Adjustment variables							
Prefectural ageing rate	%	47	31.6	3.3	22.8	31.7	39.0
General hospital beds	per 100,000	47	794.2	148.5	508.7	800.3	1,146.5
General clinics (sensitivity)	per 100,000	47	84.5	11.9	61.8	85.9	113.0
Climatic variables							
Days with WBGT ≥28°C (official, primary)	days	47	72.9	14.1	27.3	74.7	112.3
Days with WBGT ≥31°C (sensitivity)	days	47	28.7	10.6	6.0	26.5	61.6
Days with WBGT ≥33°C (sensitivity)	days	47	2.9	2.7	0.0	1.7	10.8
Days with maximum temperature ≥35°C (descriptive only)	days	47	17.9	12.1	0.0	17.0	44.0
Covariate (descriptive only)							
Air conditioning prevalence	%	47	89.7	13.6	26.6	94.8	98.2

WBGT, wet-bulb globe temperature; SD, standard deviation. WBGT days are based on official Ministry of the Environment daily-maximum observations (June–September 2023). Days with maximum temperature ≥35°C are shown for descriptive comparison and were not used as an adjustment variable. Air conditioning prevalence (2014 National Survey of Family Income and Expenditure) reflects ownership, not use, and was not used as an adjustment variable. Ageing rate and healthcare supply are the primary demographic and healthcare-supply adjustment variables in this analysis.

Table 2. Hierarchical Regression Models: Elderly Solo Household Rate and Large-Volume Infusion Therapy Utilization (N = 47)


Model	Cumulative adjustment	N	Unstd. β (HC3 95% CI)	Std. β (HC3 95% CI)	p	Exposure partial R^2	Adjusted R^2	AICc
O-A	Unadjusted	47	656.2 (326.9, 985.6)	0.521 (0.260, 0.783)	<0.001	0.272	0.256	853.2
O-B	+ ageing rate	47	306.1 (-71.3, 683.5)	0.243 (-0.057, 0.543)	0.112	0.056	0.365	847.1
O-C	+ ageing rate + hospital beds	47	94.1 (-302.4, 490.6)	0.075 (-0.240, 0.390)	0.642	0.004	0.386	846.9
O-D	+ ageing rate + hospital beds + WBGT $\geq$ 28d	47	-9.4 (-493.9, 475.1)	-0.007 (-0.392, 0.377)	0.970	<0.001	0.422	845.6

Model O-A: exposure alone. Model O-B: Model O-A + prefectural ageing rate. Model O-C: Model O-B + general hospital beds per 100,000. Model O-D: Model O-C + days with WBGT $\geq$ 28°C (each model cumulative). CI, confidence interval (HC3 robust); VIF, variance inflation factor; AICc, corrected Akaike information criterion. Using the 2020 Census population denominator (Online Resource 1, Table S3) yields a closely comparable Model O-A estimate; inferential statistics differ because this analysis uses HC3 robust standard errors.

Figure Legends

Fig. 1 Coefficient plot for elderly solo household rate across Models O-A through O-D, showing unstandardized coefficients with HC3 95% confidence intervals and a zero reference line.

528 **Fig. 1**



529

530

531 **Online Resource 1**

532 Online Resource 1 contains supplementary tables and figures, including the 2020
533 Census-denominator model hierarchy, the alternative-exposure (denominator)
534 analysis, sensitivity and influence diagnostics, nonlinearity analyses (including the
535 median-split analysis), exploratory WBGT metrics with FDR-adjusted p-values, and
536 the comparison-outcome analysis with general outpatient utilization.