

CPS 5310

Mathematical and Computer Modeling

Spring 2017

Homework #Final

Md Mahmudul Hasan

SID: 80542629

May 12, 2017

Problem II

Part 1

Given that,

$$\frac{\partial^2 u}{\partial t^2} = -c^2 \frac{\partial^4 u}{\partial x^4} \quad (1)$$

$$\frac{\partial u}{\partial t} = 0, \quad u(x, 0) = x(L - x) \quad (2)$$

$$u(0, t) = u(L, t) = 0 \quad (3)$$

$$u_{xx}(0, t) = u_{xx}(L, t) = 0 \quad (4)$$

Substitution of $u(x, t) = X(x)T(t)$ into (6) gives us,

$$\begin{aligned} XT'' &= -c^2 X^{IV} T \\ \Rightarrow \frac{T''}{-c^2 T} &= \frac{X^{IV}}{X} = \lambda^4 \end{aligned} \quad (5)$$

Where λ is a constant.

Case I: when $\lambda = 0$

$$\frac{X^{IV}}{X} = 0 \Rightarrow X(x) = ax^3 + bx^2 + cx + d$$

Now, we have,

$$X(0)T(t) = 0 \quad \text{and} \quad X(L)T(t) = 0$$

$$\text{So, } X(0) = T(L) = 0, \quad \text{as} \quad T(t) = 0$$

And,

$$X''(0)T(t) = 0 \quad \text{and} \quad X''(L)T(t) = 0$$

$$\text{So, } X''(0) = X''(L) = 0$$

From equation (5), we can write,

$$F(0) = d = 0$$

Now,

$$\begin{aligned}
X'(x) &= 3ax^2 + 2bx + c \\
X''(x) &= 6ax + 2b \\
X''(0) &= 2b = 0 \Rightarrow b = 0 \\
X(L) &= aL^3 + cL = 0 \\
X''(L) &= 6aL = 0 \\
&\Rightarrow a = 0
\end{aligned}$$

So, $c = 0$. And we obtain, $X(x) = 0$, which we do not have any interest.

Case II: when $\lambda \neq 0$

$$\begin{aligned}
\frac{X^{IV}}{X} &= \beta^4 \\
X(x) &= A\cos(\lambda x) + B\sin(\lambda x) + Ee^{\lambda x} + Fe^{-\lambda x} \\
&= A\cos(\lambda x) + B\sin(\lambda x) + \frac{C+D}{2}e^{\lambda x} + \frac{C-D}{2}e^{-\lambda x} \\
&= A\cos(\lambda x) + B\sin(\lambda x) + C\frac{e^{\lambda x} + e^{-\lambda x}}{2} + D\frac{e^{\lambda x} - e^{-\lambda x}}{2} \\
&= A\cos(\lambda x) + B\sin(\lambda x) + C\cosh(\lambda x) + D\sinh(\lambda x)
\end{aligned}$$

Now,

$$\begin{aligned}
X'(x) &= -A\lambda\sin(\lambda x) + B\lambda\cos(\lambda x) + C\lambda\sinh(\lambda x) + D\lambda\cosh(\lambda x) \\
X''(x) &= -A\lambda^2\cos(\lambda x) - B\lambda^2\sin(\lambda x) + C\lambda^2\cosh(\lambda x) + D\lambda^2\sinh(\lambda x)
\end{aligned}$$

Then, $X(0) = A + C = 0$ and $X''(0) = -A\lambda^2 + C\lambda^2 = 0$ gives us $A = C = 0$

Now,

$$X(L) = B\sin(\lambda L) + D\sinh(\lambda L) = 0$$

$$X''(L) = -B\lambda^2\sin(\lambda L) + D\lambda^2\sinh(\lambda L) = 0$$

We got, $D = 0$

Now,

$$\begin{aligned} X(L) &= B \sin(\lambda L) = 0 \\ &= \sin(\lambda L) = 0 \quad [\text{as } B \neq 0] \\ \Rightarrow \lambda &= \frac{n\pi}{L}, \quad n = 1, 2, 3, \dots \end{aligned}$$

So,

$$X_n(x) = B \sin\left(\frac{n\pi x}{L}\right), \quad n = 1, 2, 3, \dots$$

Now,

$$\begin{aligned} T'' + c^2 \lambda^2 T &= 0 \\ \Rightarrow T_n(t) &= a_n \cos(c\lambda^2 t) + b_n \sin(c\lambda^2 t), \quad \text{where, } \lambda = \frac{n\pi}{L} \end{aligned}$$

Therefore,

$$\begin{aligned} u_n(x, t) &= X_n(x) T_n(t) \\ &= \sin\left(\frac{n\pi x}{L}\right) \left(B a_n \cos(c\lambda^2 t) + B b_n \sin(c\lambda^2 t) \right), \quad \text{where, } \lambda = \frac{n\pi}{L} \end{aligned}$$

We know,

$$\begin{aligned} u(x, t) &= \sum_{n=1}^{\infty} u_n(x, t) \\ &= \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{L}\right) \left(B a_n \cos\left(c\left(\frac{n\pi}{L}\right)^2 t\right) + B b_n \sin\left(c\left(\frac{n\pi}{L}\right)^2 t\right) \right) \end{aligned}$$

$$\begin{aligned} \left. \frac{\partial u}{\partial t} \right|_{t=0} &= \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{L}\right) \left(-B a_n c \left(\frac{n\pi}{L}\right)^2 \sin\left(c\left(\frac{n\pi}{L}\right)^2 t\right) + B b_n c \left(\frac{n\pi}{L}\right)^2 \cos\left(c\left(\frac{n\pi}{L}\right)^2 t\right) \right) \Big|_{t=0} \\ \Rightarrow 0 &= \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{L}\right) B b_n c \left(\frac{n\pi}{L}\right)^2 \\ \therefore B b_n &= 0. \end{aligned}$$

Therefore,

$$u(x, t) = \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{L}\right) B a_n \cos\left(c\left(\frac{n\pi}{L}\right)^2 t\right).$$

Now,

$$u(x, 0) = \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{L}\right) Ba_n = f(x) = x(L - x),$$

where,

$$\begin{aligned} Ba_n &= \frac{2}{L} \int_0^L x(L - x) \sin\left(\frac{n\pi x}{L}\right) dx \\ &= \frac{2}{L} \int_0^L Lx \sin\left(\frac{n\pi x}{L}\right) dx - \frac{2}{L} \int_0^L x^2 \sin\left(\frac{n\pi x}{L}\right) dx \\ &= I_1 - I_2. \end{aligned}$$

So,

$$\begin{aligned} I_1 &= \frac{2}{L} L \left[\frac{-x \cos\left(\frac{n\pi x}{L}\right)}{\frac{n\pi}{L}} + \frac{\sin\left(\frac{n\pi x}{L}\right)}{\frac{n^2 \pi^2}{L^2}} \right]_0^L \\ &= \frac{2}{L} \left[-\frac{L^3}{n\pi} \cos(n\pi) + \frac{L^3}{n^2 \pi^2} \sin(n\pi) \right] \\ &= \frac{2}{L} \left[-\frac{L^3}{n\pi} \cos(n\pi) \right] \end{aligned}$$

And

$$\begin{aligned} I_2 &= \frac{2}{L} \left[\frac{-x^2 \cos\left(\frac{n\pi x}{L}\right)}{\frac{n\pi}{L}} + \frac{2x \sin\left(\frac{n\pi x}{L}\right)}{\frac{n^2 \pi^2}{L^2}} + \frac{2 \cos\left(\frac{n\pi x}{L}\right)}{\frac{n^3 \pi^3}{L^3}} \right]_0^L \\ &= \frac{2}{L} \left[-\frac{L^3}{n\pi} \cos(n\pi) + \frac{2L^3}{n^2 \pi^2} \sin(n\pi) + \frac{2L^3}{n^3 \pi^3} \cos(n\pi) - \frac{2L^3}{n^3 \pi^3} \right] \\ &= \frac{2}{L} \left[-\frac{L^3}{n\pi} \cos(n\pi) + \frac{2L^3}{n^3 \pi^3} \cos(n\pi) - \frac{2L^3}{n^3 \pi^3} \right] \quad [\text{since, } \sin(n\pi) = 0] \end{aligned}$$

So,

$$\begin{aligned} Ba_n &= \frac{2}{L} \left[\frac{2L^3}{n^3 \pi^3} - \frac{2L^3}{n^3 \pi^3} \cos(n\pi) \right] \\ &= \frac{4L^2}{n^3 \pi^3} - \frac{4L^2}{n^3 \pi^3} \cos(n\pi) \end{aligned}$$

Therefore,

$$u(x, t) = \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{L}\right) \left[\frac{4L^2}{n^3\pi^3} - \frac{4L^2}{n^3\pi^3} \cos(n\pi) \right] \cos\left(c\left(\frac{n\pi}{L}\right)^2 t\right).$$

Part 2

Given that,

$$\frac{\partial^2 u}{\partial t^2} = -c^2 \frac{\partial^4 u}{\partial x^4} \quad (6)$$

$$\frac{\partial u}{\partial t} = 0, \quad u(x, 0) = x(L - x) \quad (7)$$

$$u(0, t) = u(L, t) = 0 \quad (8)$$

$$u_{xx}(0, t) = u_{xx}(L, t) = 0 \quad (9)$$

Let,

$$\frac{\partial u(t)}{\partial t} = v(t) \quad \text{and} \quad \Rightarrow \frac{\partial^2 u(t)}{\partial t^2} = \frac{\partial v(t)}{\partial t}$$

From equation (6) we get,

$$\frac{\partial v(t)}{\partial t} = -c^2 \frac{\partial^4 u(t)}{\partial x^4}$$

From equation (7) we get,

$$\left. \frac{\partial u}{\partial t} \right|_{t=0} = 0$$

$$\Rightarrow v(0) = 0 = v_0$$

$$u(x, 0) = u(0) = x(L - x) = u_0$$

Therefore,

$$\frac{\partial u(t)}{\partial t} = v(t),$$

$$\frac{\partial v(t)}{\partial t} = -c^2 \frac{\partial^4 u(t)}{\partial x^4},$$

$$u(0) = x(L - x), \quad v(0) = 0.$$

$$F(u(t)) = -c^2 \frac{\partial^4 u(t)}{\partial x^4},$$

$$u_0 = x(L - x),$$

$$v_0 = 0.$$

Part 2

Let,

$$\mathbb{X} = \begin{pmatrix} u \\ v \end{pmatrix}$$

So,

$$\begin{aligned} \frac{d\mathbb{X}}{dt} &= \begin{pmatrix} \frac{\partial u}{\partial t} \\ \frac{\partial v}{\partial t} \end{pmatrix} = \begin{pmatrix} v \\ -c^2 \frac{\partial^4 u}{\partial x^4} \end{pmatrix} \\ \frac{\mathbb{X}(n) - \mathbb{X}(n-1)}{\tau} &= \begin{pmatrix} \frac{u(n) - u(n-1)}{\tau} \\ \frac{v(n) - v(n-1)}{\tau} \end{pmatrix} \\ &= \begin{pmatrix} v(n-1) \\ -c^2 \frac{\partial^4 u(n-1)}{\partial x^4} \end{pmatrix} \end{aligned}$$

Where, $\tau = \frac{1}{3}$

Now,

$$u(n) - u(n-1) = \tau v(n-1)$$

$$v(n) - v(n-1) = -\tau c^2 \frac{\partial^4 u(n-1)}{\partial x^4}$$

Hence,

$$u(n) = u(n-1) + \tau v(n-1)$$

$$v(n) = v(n-1) - \tau c^2 \frac{\partial^4 u(n-1)}{\partial x^4}$$

In matrix form we can write,

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} u(n) \\ v(n) \end{pmatrix} = \begin{pmatrix} u(n-1) + \tau v(n-1) \\ v(n-1) - \tau c^2 \frac{\partial^4 u(n-1)}{\partial x^4} \end{pmatrix}.$$

So,

$$A_\tau = 1, \quad u_\tau = u(n), \quad v_\tau = v(n)$$

$$f_\tau = u(n-1) + \tau v(n-1)$$

$$g_\tau = v(n-1) - \tau c^2 \frac{\partial^4 u(n-1)}{\partial x^4}.$$

Problem III

Part 1

Here,

V = feasible region

$$\begin{aligned}
 &= \text{space of real valued functions } w \text{ on } \overline{\Omega} = \partial\Omega \text{ such that } w, \text{ and } \nabla w \text{ are integrable} \\
 &= \{w : \overline{\Omega} \rightarrow \mathbb{R} | w, \text{ and } \nabla w \text{ are } L^2 \text{ on } \omega \text{ and } w = \nabla \cdot \nabla w = 0 \text{ on } \partial\Omega\} \\
 &= H_0^2(\Omega) \\
 &= \overline{C_0^2(\Omega)}
 \end{aligned}$$

Part 2

Given that,

$$J(v) = \frac{1}{2} \int_{\Omega} \left(\frac{d^2 v}{dx^2}\right)^2 dx - \int_{\Omega} f v dx.$$

We have to find $u \in V$ such that,

$$J(u) = \min_{v \in V} J(v)$$

This minimization problem has a unique solution $u \in V$ under certain condition. We have to find $u \in V$ such that,

$$\langle J'(u), v \rangle = 0 \quad \forall v \in V$$

When $\lambda > 0$,

$$\begin{aligned}
 \frac{J(u + \lambda v)}{\lambda} &= \frac{1}{\lambda} \left[\frac{1}{2} \int_{\Omega} \left(\frac{d^2(u + \lambda v)}{dx^2}\right)^2 dx - \int_{\Omega} f(u + \lambda v) dx - \frac{1}{2} \int_{\Omega} \left(\frac{d^2 u}{dx^2}\right)^2 dx + \int_{\Omega} f u dx \right] \\
 &= \frac{1}{\lambda} \left[\frac{1}{2} \int_{\Omega} \left(\frac{d^2 u}{dx^2} + \lambda \frac{d^2 v}{dx^2}\right)^2 dx - \int_{\Omega} f u dx - \lambda \int_{\Omega} f v dx - \frac{1}{2} \int_{\Omega} \left(\frac{d^2 u}{dx^2}\right)^2 dx + \int_{\Omega} f u dx \right] \\
 &= \frac{1}{\lambda} \left[\frac{1}{2} \int_{\Omega} \left(\frac{d^2 u}{dx^2}\right)^2 dx + 2\lambda \int_{\Omega} \Delta u \cdot \Delta v dx + \lambda^2 \int_{\Omega} \left(\frac{d^2 v}{dx^2}\right)^2 dx \right] - \lambda \int_{\Omega} f v dx - \frac{1}{2} \int_{\Omega} \left(\frac{d^2 u}{dx^2}\right)^2 dx \\
 &= \int_{\Omega} \Delta u \cdot \Delta v dx + \frac{\lambda}{2} \int_{\Omega} \left(\frac{d^2 v}{dx^2}\right)^2 dx - \lambda \int_{\Omega} f v dx
 \end{aligned}$$

So,

$$\lim_{\lambda \rightarrow 0^+} \frac{J(u + \lambda v)}{\lambda} = \int_{\Omega} \Delta u \cdot \Delta v dx - \lambda \int_{\Omega} f v dx = 0$$

Now, we have,

$$\int_{\Omega} \Delta u \cdot \Delta v dx = \int_{\Omega} f v dx, \quad \forall v \in V$$

So, the boundary value problem becomes,

$$\Delta^2 u = f \text{ on } \Omega \tag{10}$$

$$u = 0 \text{ on } \partial\Omega \quad \text{and} \quad \Delta u = 0 \text{ on } \partial\Omega$$

Multiplying equation(10) by v and then using integration by parts, we get,

$$\begin{aligned} \int_{\Omega} \Delta u \Delta v dx - \int_{\partial\Omega} \Delta u \nabla v dx + \int_{\partial\Omega} \Delta u v dx &= \int_{\Omega} f v dx \\ \int_{\Omega} \Delta u \cdot \Delta v dx &= \int_{\Omega} f v dx \quad [\text{Since, } u = 0 \text{ and } \Delta u = 0 \text{ on } \partial\Omega] \end{aligned}$$

Part 3

Here,

$$\begin{aligned} u(x) &= \frac{f_0}{24}(x^4 - 2Lx^3 + L^3x) \\ u'(x) &= \frac{f_0}{24}(4x^3 - 6Lx^2 + L^3) \\ u''(x) &= \frac{f_0}{24}(12x^2 - 12Lx) \\ u'''(x) &= \frac{f_0}{24}(24x - 12L) \\ u^{IV}(x) &= f_0 \end{aligned}$$

Now,

$$\begin{aligned}u(0) &= \frac{f_0}{24}(0 - 0 + 0) \\ &= 0\end{aligned}$$

$$\begin{aligned}u(L) &= \frac{f_0}{24}(L^4 - 2L^4 + L^4) \\ &= 0\end{aligned}$$

$$\begin{aligned}u''(0) &= \frac{f_0}{24}(0 - 0) \\ &= 0\end{aligned}$$

$$\begin{aligned}u''(L) &= \frac{f_0}{24}(12L^2 - 12L^2) \\ &= 0\end{aligned}$$

So, $u(x)$ satisfies the boundary value problem

$$u^{IV}(x) = f(x)$$

with the same boundary conditions

$$u(0) = u(L) = 0$$

$$\text{and } u''(0) = u''(L) = 0$$