

FUNCTIONS

CHAPTER 4

Lines & Quadratics

4

SYLLABUS 1.16 2.1–2.4 2.6 2.7

On 17 January 1995 an earthquake struck near Kobe, in western Japan. The Akashi Kaikyō Bridge was under construction: its two towers already stood in the strait, but the roadway between them did not yet exist. The earthquake moved the towers apart.

The engineers did not start again. They measured the new gap and adjusted the design to fit it. The bridge that opened in 1998 has a main span of 1991 metres instead of the planned 1990.

They could do this because the design was written as equations, not as fixed drawings. The main cables of a suspension bridge carry a level roadway of even weight, and the curve they hang in is a *parabola*, the same curve as a thrown ball. Change one value in the equations and every other measurement follows from it.

That is the power of a **function**. It takes an input and returns exactly one output, so a whole structure becomes a single rule you can graph, study and recalculate. This chapter builds the functions the rest of the course rests on: lines, which change at a steady rate, and quadratics, which give the cable its curve.

By the end of this chapter you will be able to give a function's domain, range and inverse, and to find the equation of a line from whatever you are told. You will solve systems of linear equations, move a quadratic between its factorised, vertex and expanded forms, and use the discriminant to count the roots of an equation.

4.1 Functions

A function is a rule you can rely on. Give it an input and it returns one definite output, and it returns the same output every time. That is what makes it possible to reason about.

DEF A **function** f assigns exactly one output $f(x)$ to each input x . The set of inputs it accepts is the **domain**; the set of outputs that actually occur is the **range**.

The phrase “exactly one” is the whole idea. A rule that could return two different outputs for the same input is not a function. On a graph this becomes the **vertical line test**: a curve represents a function only if no vertical line crosses it more than once.

Reading the domain

A function need not accept every number. Three operations, and only three, put limits on what you may substitute. Learn the three and you can read the domain of almost any function on the syllabus.

Where x appears	Condition	Reason
In a denominator	$\neq 0$	you cannot divide by zero
Inside an even root	≥ 0	a negative number has no real square root
Inside a logarithm	> 0	the logarithm of zero, or of a negative number, does not exist

Note the difference between the last two. A square root allows zero, because $\sqrt{0} = 0$. A logarithm does not, because no power of the base gives zero. That single distinction, $\geq$ against $>$, is the most common slip in this topic.

Function	Condition	Domain
$\sqrt{x+3}$	$x+3 \geq 0$	$x \geq -3$
$\sqrt{5-x}$	$5-x \geq 0$	$x \leq 5$
$\frac{1}{x-2}$	$x-2 \neq 0$	$x \neq 2$
$\ln(x-4)$	$x-4 > 0$	$x > 4$
$\frac{1}{\sqrt{x-1}}$	$x-1 > 0$, since it is both under a root and in a denominator	$x > 1$

When a function contains more than one of these, every condition must hold at once. Write them all down, then take the values that satisfy all of them.

EXAMPLE 4.1

State the domain of $f(x) = \frac{\sqrt{x+2}}{x-1}$.

There are two restrictions here. The square root needs $x+2 \geq 0$, so $x \geq -2$. The denominator needs $x-1 \neq 0$, so $x \neq 1$.

Both must hold, so the domain is

$$x \geq -2, \quad x \neq 1.$$

Notice that $x = 1$ satisfies the first condition but fails the second, so it must still be removed.

EXAMPLE 4.2

State the domain of $g(x) = \ln(9 - x^2)$.

A logarithm needs a positive argument, so $9 - x^2 > 0$, that is $x^2 < 9$. This is a quadratic inequality, solved in §4.7: the parabola $x^2 - 9$ lies below the axis between its roots. So

$$-3 < x < 3.$$

Both ends are excluded, because at $x = \pm 3$ the argument is zero and $\ln 0$ does not exist.

CAUTION Solve the condition, do not guess it. For $\sqrt{5-x}$ the domain is $x \leq 5$, not $x \geq 5$. Write the inequality down and rearrange it every time, however obvious it looks.

Reading the range

The range is the set of outputs that actually occur. There is no single rule, but four facts cover most functions you will meet, and each says that some expression can never go below zero.

Almost every range argument rests on one fact: some part of the function can never be negative. A square is never negative, so $x^2 \geq 0$; a modulus is never negative, so $|x| \geq 0$; the square root symbol means the non-negative root, so $\sqrt{x} \geq 0$; and a positive base raised to any power stays positive, so $a^x > 0$.

To use them, build the function up from the part you know. Ask what that part can be, then follow the operations one at a time.

Function	Reasoning	Range
$x^2 + 1$	$x^2 \geq 0$, then add 1	$y \geq 1$
$ x - 3$	$ x \geq 0$, then subtract 3	$y \geq -3$
$\sqrt{x} + 2$	$\sqrt{x} \geq 0$, then add 2	$y \geq 2$
$3 - x + 1 $	$ x + 1 \geq 0$, so subtracting it takes 3 downward	$y \leq 3$
$-(x - 2)^2 + 5$	$(x - 2)^2 \geq 0$, and the minus sign turns the inequality round	$y \leq 5$

CAUTION A minus sign in front reverses the direction. Since $(x-2)^2 \geq 0$, it follows that $-(x-2)^2 \leq 0$, so $-(x-2)^2 + 5 \leq 5$. Multiplying an inequality by a negative number always flips the sign.

For a quadratic, complete the square first. The vertex form shows the smallest or largest output immediately.

EXAMPLE 4.3

State the range of $f(x) = x^2 - 6x + 11$.

Complete the square:

$$x^2 - 6x + 11 = (x - 3)^2 - 9 + 11 = (x - 3)^2 + 2.$$

Now use the first fact. Since $(x - 3)^2 \geq 0$, adding 2 gives $f(x) \geq 2$. The range is $y \geq 2$. The smallest value, 2, occurs at $x = 3$, which is the vertex.

EXAMPLE 4.4

State the domain and range of $g(x) = \sqrt{x - 2}$.

Domain. The square root needs $x - 2 \geq 0$, so $x \geq 2$.

Range. As x runs from 2 upward, $x - 2$ runs from 0 upward, and so does its square root. So $g(x) \geq 0$.

The domain comes from the inputs the rule permits. The range comes from the outputs those inputs give. Always ask both questions.

EXAMPLE 4.5

State the domain and range of $f(x) = \sqrt{2 - x}$.

The square root needs $2 - x \geq 0$, that is $x \leq 2$. So the domain is $x \leq 2$.

As x decreases from 2, the quantity $2 - x$ increases from 0 upward, and so does its square root. The range is $f(x) \geq 0$.

Compare this with $\sqrt{x - 2}$ above. The two look similar but their domains point in opposite directions, so read the inequality from the expression each time rather than from memory.

TIP On Paper 2, check a range on your GDC. Graph the function and look at the lowest and highest

points the curve reaches. The algebra is still expected in the working, but the graph catches a sign error in seconds.

Reading a graph

Most of what you need from a function can be read straight from its graph. Five features matter, and each has its own name.

Feature	What it is	How to find it
x-intercepts (also called zeros, or roots)	the points where the curve meets the x -axis	set $y = 0$ and solve
y-intercept	the point where the curve meets the y -axis	set $x = 0$
Turning points	the local maximum and minimum points, where the curve changes direction	GDC, or the vertex form for a quadratic
Asymptote	a line that the curve gets closer and closer to, without ever reaching it	look for values the function cannot take
Intersection	a point that lies on both of two graphs	solve the two equations together

Your GDC finds all five directly. The graph menu has zero, min/max and intersect tools. On Paper 2 these are the intended method. Graph the function, read the feature, and write down what you asked the calculator to find. Many equations have no algebraic solution at all. For those the graph is not a faster method. It is the only method.

TIP Know the difference between the command terms: **sketch** means show the shape with key features labelled, not to scale; **draw** means an accurate, to-scale plot. A sketch of a parabola needs the vertex, the intercepts, and the correct opening direction, and nothing else.

Inverse functions

An **inverse function** reverses the rule. If f sends 3 to 7, then f^{-1} sends 7 back to 3. An inverse exists only when no two inputs give the same output. If two did, the way back would have two answers, and a function may return only one.

DEF The **inverse** f^{-1} of a function f undoes it: $f^{-1}(f(x)) = x$. It exists when f is **one-to-one**, meaning no two inputs share an output (the *horizontal* line test).

The inverse swaps input and output. Two things follow. Its graph is the mirror image of f in the line $y = x$, and its domain and range are those of f exchanged. To find a formula, write $y = f(x)$, swap x and y , then solve for y .

EXAMPLE 4.6

Find the inverse of $f(x) = 2x + 3$.

Write $y = 2x + 3$, then swap x and y :

$$x = 2y + 3.$$

Solving for y ,

$$y = \frac{x-3}{2}, \quad \text{so} \quad f^{-1}(x) = \frac{x-3}{2}.$$

Check: f doubles and adds three; f^{-1} subtracts three and halves. Each step of f is undone, in reverse order.




Figure 4.1 A function and its inverse are mirror images in the line $y = x$.

The inverse also gives another way to read an equation. Asking “for which x does $f(x) = 10$?” is the same as asking “what does f^{-1} send 10 to?”

KEY Solving $f(x) = 10$ is equivalent to evaluating $f^{-1}(10)$.

EXAMPLE 4.7

For $f(x) = 2x + 3$, solve $f(x) = 10$ in two ways.

Directly: $2x + 3 = 10$ gives $2x = 7$, so $x = \frac{7}{2}$.

Using the inverse found above, $f^{-1}(x) = \frac{x-3}{2}$, so

$$f^{-1}(10) = \frac{10-3}{2} = \frac{7}{2}.$$

The two routes must agree, because they ask the same question. When a question gives you f^{-1} already, the second route is the shorter one.

CAUTION The notation f^{-1} means the inverse, not the reciprocal. $f^{-1}(x)$ is not $\frac{1}{f(x)}$.

YOUR TURN 4.1 Functions

1. State the domain of each function.

(a) $\sqrt{x+3}$

(b) $\sqrt{7-x}$

(c) $\frac{1}{x-2}$

(d) $\frac{1}{x+5}$

(e) $\ln(x-4)$

(f) $\ln(2x+1)$

(g) $\frac{1}{\sqrt{x-1}}$

(h) $\frac{1}{x^2-9}$

2. State the domain of each function. Each has more than one restriction.

(a) $\frac{\sqrt{x+2}}{x-1}$

(b) $\frac{\sqrt{x}}{x-9}$

3. State the range of each function on the domain given.

(a) $x^2 + 1, \quad x \in \mathbb{R}$

(b) $|x| - 3, \quad x \in \mathbb{R}$

(c) $\sqrt{x} + 2, \quad x \geq 0$

(d) $3 - |x+1|, \quad x \in \mathbb{R}$

4. Find the range of each quadratic by completing the square. Each is defined for all real x . If you have not met this method yet, it is set out in §4.5.

(a) $x^2 - 6x + 11$

(b) $x^2 - 4x$

(c) $-(x-2)^2 + 5$

(d) $-x^2 + 6x - 4$

5. For $f(x) = x^2 - 3x$, find each value.

(a) $f(2)$

(b) $f(-1)$

(c) $f(0)$

(d) the values of x with $f(x) = 0$

6. Find the inverse of each function.

(a) $f(x) = 3x - 6$

(b) $f(x) = \frac{x+1}{2}$

(c) $f(x) = 2x + 7$

(d) $f(x) = \frac{1}{x+2}$

(e) $f(x) = \frac{2x+1}{x-3}$

(f) $f(x) = \frac{x-4}{x+2}$

7. Consider $f(x) = \frac{3x+2}{x-1}$.

- (a) State the domain of f .
(b) Find $f^{-1}(x)$.
(c) State the domain of f^{-1} .
(d) Verify that $f^{-1}(f(2)) = 2$.

8. Decide whether each rule defines y as a function of x , and justify your answer.

- (a) $y^2 = x$ (b) $y = x^2$

9. The function $f(x) = x^2$ has no inverse over all real numbers. State a restricted domain on which an inverse does exist, and give that inverse.

4.2 Linear Functions

The simplest function that actually changes is the line: its output rises at a constant rate. That rate is the **gradient**, the rise divided by the run.

KEY · IB The gradient of the line through (x_1, y_1) and (x_2, y_2) is

$$m = \frac{y_2 - y_1}{x_2 - x_1}.$$

A gradient and one point on the line are enough to fix the line completely. From that fact come three ways to write its equation. All three describe the same line.

KEY · IB A straight line can be written

$$y = mx + c, \quad y - y_1 = m(x - x_1), \quad ax + by + d = 0.$$

Here m is the gradient and c the y -intercept.

Use the first form when you know the gradient and the y -intercept. Use the second when you know the gradient and any point. The third is the tidy general form, with whole-number coefficients.

Two lines are **parallel** when they rise at the same rate. They are **perpendicular** when you get one gradient from the other by turning the fraction upside down and changing the sign.

KEY For two lines with gradients m_1 and m_2 :

$$m_1 = m_2 \text{ (parallel),} \quad m_1 m_2 = -1 \text{ (perpendicular).}$$




Figure 4.2 Perpendicular gradients: turn the fraction upside down and change the sign.

EXAMPLE 4.8

Find the equation of the line through $(1, 2)$ and $(4, 8)$.

First the gradient:

$$m = \frac{8 - 2}{4 - 1} = \frac{6}{3} = 2.$$

Then use the point $(1, 2)$ in point–gradient form:

$$y - 2 = 2(x - 1) \implies y = 2x.$$

EXAMPLE 4.9

Find the equation of the line through $(0, 4)$ perpendicular to $y = 3x - 1$.

The given line has gradient 3, so the perpendicular gradient is $-\frac{1}{3}$. Through $(0, 4)$,

$$y = -\frac{1}{3}x + 4.$$

For a perpendicular gradient, turn the fraction upside down and change the sign.

YOUR TURN 4.2 Linear Functions

10. Find the gradient of the line through each pair of points.

(a) $(1, 3)$ and $(5, 11)$

(b) $(2, 1)$ and $(4, 7)$

(c) $(-1, 4)$ and $(3, -4)$

(d) $(0, 5)$ and $(2, 5)$

11. Find the equation of each line, giving your answer as $y = mx + c$.

- (a) gradient 2, through $(0, -3)$ (b) through $(2, 1)$ and $(4, 7)$
 (c) gradient $-\frac{1}{2}$, through $(4, 0)$ (d) through $(0, 5)$, perpendicular to $y = -2x$

12. Write each equation as $y = mx + c$ and state the gradient.

- (a) $2x + 3y = 6$ (b) $4x - 2y = 8$

13. Answer each question about parallel and perpendicular lines.

- (a) Write down the gradient of a line perpendicular to $y = 4x + 1$.
 (b) Write down the gradient of a line parallel to $3x + y = 7$.
 (c) Are $y = 2x + 1$ and $y = 2x - 3$ parallel? Give a reason.
 (d) Are $y = 3x - 1$ and $y = -\frac{1}{3}x + 2$ perpendicular? Give a reason.

4.3 Systems of Linear Equations HL

Two lines in a plane usually cross at one point. Finding that point means finding the values of x and y that satisfy *both* equations at once, a **system** of equations. The standard tool is **elimination**: combine the equations to cancel one variable, solve for the other, then substitute back.

EXAMPLE 4.10

Solve $2x + y = 7$ and $x - y = 2$.

Adding the two equations cancels y :

$$(2x + y) + (x - y) = 7 + 2 \implies 3x = 9 \implies x = 3.$$

Substituting into $x - y = 2$ gives $3 - y = 2$, so $y = 1$. The lines cross at $(3, 1)$.

Not every system has a single answer, and the picture shows why. Two lines can do three things. They cross once, which gives a **unique solution**. They stay parallel and never meet, so there is **no solution**. Or they are the same line drawn twice, which gives **infinitely many solutions**.

CAUTION If two equations have the same gradient but different intercepts, such as $2x + y = 3$ and $4x + 2y = 10$, the lines are parallel and the system has no solution. Elimination then produces a false statement such as $0 = 4$.

The same idea extends to three equations in three unknowns, where each equation is now a plane in space. You eliminate one variable at a time to reduce a 3×3 system to a 2×2 one,

or read the solution directly from a GDC.

EXAMPLE 4.11

Solve the system

$$x + y + z = 6, \quad 2x - y + z = 3, \quad x + 2y - z = 2.$$

Eliminate z in pairs. Subtracting the first equation from the second:

$$(2x - y + z) - (x + y + z) = 3 - 6 \implies x - 2y = -3.$$

Adding the first and third equations:

$$(x + 2y - z) + (x + y + z) = 2 + 6 \implies 2x + 3y = 8.$$

Now solve this 2×2 system. From $x - 2y = -3$ we get $x = 2y - 3$; substituting,

$$2(2y - 3) + 3y = 8 \implies 7y = 14 \implies y = 2,$$

so $x = 1$. Finally the first equation gives $1 + 2 + z = 6$, hence $z = 3$. The solution is $(x, y, z) = (1, 2, 3)$.

Eliminating one variable at a time, working down the system, is called **row reduction**. It is what your GDC's simultaneous-equation solver does inside. Use the solver to check your work. You still need the algebra in an examination, and you need it when the solver returns only an error message.

The augmented matrix

Look again at the elimination above and notice what you were really changing: never the letters x, y, z , only the numbers in front of them. The letters only marked the columns, so you can stop writing them. Reduce the system to its coefficients and constants, written in a grid called the **augmented matrix**:

$$\begin{cases} x + y + z = 6 \\ 2x - y + z = 7 \\ x + 2y - z = 3 \end{cases} \longrightarrow \left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 2 & -1 & 1 & 7 \\ 1 & 2 & -1 & 3 \end{array} \right)$$

The vertical bar shows where the equals signs were. Nothing new is happening: the matrix records the same elimination in a shorter form, with fewer symbols to copy incorrectly.

Elimination becomes three **row operations**. None of them changes the solution set. Each one is reversible, and each does to a whole equation only what you were always allowed to do:

KEY The three elementary row operations:

$$R_i \leftrightarrow R_j \text{ (swap two rows),} \quad R_i \rightarrow kR_i \text{ (multiply a row by } k \neq 0\text{),} \quad R_i \rightarrow R_i + kR_j.$$

Swapping reorders the equations, scaling multiplies one through by a constant, and the third adds a multiple of one equation to another: the same three moves as elimination, written compactly.

The goal is a step shape, with zeros below the first entry of each row. The last row then contains one unknown, the row above it two, and so on. This shape is called **row echelon form**. Once you reach it, work upwards from the last row, finding one unknown at a time. That is **back-substitution**.

EXAMPLE 4.12

Solve, by row reduction,

$$x + y + z = 6, \quad 2x - y + z = 7, \quad x + 2y - z = 3.$$

Work on the augmented matrix, clearing the first column below the pivot. $R_2 \rightarrow R_2 - 2R_1$ and $R_3 \rightarrow R_3 - R_1$:

$$\left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 2 & -1 & 1 & 7 \\ 1 & 2 & -1 & 3 \end{array} \right) \rightarrow \left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & -3 & -1 & -5 \\ 0 & 1 & -2 & -3 \end{array} \right)$$

A leading entry of 1 gives simpler arithmetic than -3 , so swap $R_2 \leftrightarrow R_3$. Then clear below it with $R_3 \rightarrow R_3 + 3R_2$:

$$\rightarrow \left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 1 & -2 & -3 \\ 0 & -3 & -1 & -5 \end{array} \right) \rightarrow \left(\begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 0 & 1 & -2 & -3 \\ 0 & 0 & -7 & -14 \end{array} \right)$$

The step shape is complete. Now read it from the bottom up. The last row says $-7z = -14$, so $z = 2$. The middle row says $y - 2z = -3$, so $y = 4 - 3 = 1$. The top row gives $x = 6 - y - z = 3$. The solution is $(x, y, z) = (3, 1, 2)$.

Every step here is the elimination you already know. The matrix keeps the working short and the columns lined up, and misaligned columns are the usual cause of mistakes.

Once a system is in row echelon form, the last row determines the outcome. There are only three possible outcomes, and every 3×3 system gives exactly one of them.

Last row reduces to	What it means	The three planes
$(0\ 0\ a\ \ b),\ a \neq 0$	one unique solution	meet at a single point
$(0\ 0\ 0\ \ c),\ c \neq 0$	no solution: the row says $0 = c$, which is false	have no common point; two are parallel, or the three meet in a triangular prism
$(0\ 0\ 0\ \ 0)$	infinitely many solutions: the row says $0 = 0$, which is true but tells us nothing	share a whole line, or a whole plane if two rows vanish

Count the rows that vanish. No vanishing row gives one solution. One vanishing row gives a line of solutions, described by one free variable. Two vanishing rows mean all three equations describe the same plane, and the solutions form that whole plane, described by two free variables.

CAUTION Read the constant on the right before deciding. The rows $(0\ 0\ 0\ |\ 0)$ and $(0\ 0\ 0\ |\ 5)$ look almost identical, and they give opposite answers: infinitely many solutions, and none at all.

When a system has no unique solution

Each equation in three unknowns is a plane in space. Three planes usually meet at exactly one point. That point is the unique solution found in the example above.

Two other arrangements are possible. If two planes are parallel, or if the three meet in three parallel lines like the faces of a triangular prism, no point lies on all three. The system is **inconsistent**, and there is no solution. If instead the three planes share a whole line, like pages sharing a spine, every point of that line is a solution, so there are infinitely many.

You can separate the two cases by algebra alone, without a picture. If elimination produces a false statement such as $0 = 4$, the planes have no common point, and there is no solution. If it produces $0 = 0$, one equation repeats information already contained in the other two. There are infinitely many solutions, and one variable may take any value.

EXAMPLE 4.13

Consider the system

$$x + y + 2z = 2, \quad 2x - y + z = 1, \quad x + 4y + 5z = k.$$

Find the value of k for which the system has infinitely many solutions, and give the general solution. Show that for every other value of k there is no solution.

The question asks for infinitely many solutions, so expect a row that reduces to zeros. Eliminate

x using the first equation. Second minus twice the first:

$$(2x - y + z) - 2(x + y + 2z) = 1 - 4 \Rightarrow -3y - 3z = -3 \Rightarrow y + z = 1.$$

Third minus the first:

$$(x + 4y + 5z) - (x + y + 2z) = k - 2 \Rightarrow 3y + 3z = k - 2 \Rightarrow y + z = \frac{k-2}{3}.$$

The same left side, $y+z$, has appeared twice. The two remaining equations are consistent only if

$$1 = \frac{k-2}{3} \Rightarrow k = 5.$$

If $k \neq 5$, subtracting them gives $0 = \frac{k-5}{3} \neq 0$, a false statement: the system is inconsistent, and no solution exists for any such k .

If $k = 5$, the third equation gives no new information, and the system reduces to two equations. Let the free variable be $z = t$. Then $y = 1 - t$, and the first equation gives

$$x = 2 - y - 2z = 2 - (1 - t) - 2t = 1 - t.$$

The general solution is

$$(x, y, z) = (1 - t, 1 - t, t), \quad t \in \mathbb{R},$$

a whole line of solutions: the line where the three planes meet. Check with $t = 0$: the point $(1, 1, 0)$ satisfies all three original equations. So does $(0, 0, 1)$, from $t = 1$. One free parameter gives one line, so the geometry and the algebra give the same result.

CAUTION “No unique solution” covers two different cases, so always separate them. A row that reduces to $0 = 0$ gives infinitely many solutions. A row that reduces to $0 = c$, with $c \neq 0$, gives none. Examiners ask for both, and for the general solution in terms of a parameter.

YOUR TURN 4.3 Systems of Linear Equations HL

14. Solve each system by elimination.

(a) $x + y = 5, \quad x - y = 1$

(b) $3x - y = 5, \quad x + y = 3$

(c) $x + y = 4, \quad 2x + y = 7$

(d) $2x + 3y = 12, \quad x - y = 1$

15. Solve each system by substitution.

(a) $y = 2x, \quad x + y = 6$

(b) $y = x - 1, \quad 2x + y = 8$

16. State how many solutions each system has, and give a reason.

(a) $2x + y = 4, \quad 4x + 2y = 9$

(b) $y = x + 1, \quad y = x + 3$

(c) $x + y = 3, \quad 2x + 2y = 6$

(d) $x + y = 3, \quad x - y = 1$

17. Solve each system of three equations.

- (a) $x + y + z = 6$, $x - y + z = 2$, $x + y - z = 0$ (b) $x + y + z = 4$, $2x - y + z = 8$, $x + 2y - z = -3$
- (c) $x + y + z = 2$, $2x - y + 3z = -1$, $x - 2y + z = -1$ (d) $2x + y - z = -4$, $x + 3y + 2z = 3$, $3x - y + z = 9$

18. For parts (a) to (c), write the augmented matrix, reduce it to row echelon form, and state the number of solutions.

- (a) $x + y + z = 6$, $x + 2y - z = 2$, $2x + 3y + z = 11$
- (b) $x + y + z = 3$, $2x + 2y + 2z = 7$, $x - y + z = 1$
- (c) $x + y + z = 6$, $x - y + z = 2$, $2x + 2z = 8$
- (d) For $x + y + z = 3$, $2x + y + 3z = 7$, $3x + 2y + 4z = k$, find the value of k for which the system has infinitely many solutions, and state what happens for every other value of k .

4.4 Quadratic Functions

A **quadratic** is a function whose highest power is x^2 . Its graph is the **parabola**, the curve of Galileo's cannonball: a single smooth arch, symmetric about a vertical line through its turning point.

The same quadratic can be written in three forms, and each form shows a different feature immediately.

KEY A quadratic can be written in three forms:

$$y = ax^2 + bx + c, \quad y = a(x - p)(x - q), \quad y = a(x - h)^2 + k.$$

Standard form shows the y -intercept c ; **factored** form shows the roots p and q ; **vertex** form shows the turning point (h, k) .

The sign of a sets the direction. If $a > 0$ the parabola opens upward and the turning point is a minimum. If $a < 0$ it opens downward and the turning point is a maximum. The curve is symmetric, and its mirror line passes through the vertex.

KEY · IB The graph of $f(x) = ax^2 + bx + c$ has axis of symmetry

$$x = -\frac{b}{2a}.$$

To move from standard form to vertex form, **complete the square**. This means rewriting $x^2 + \frac{b}{a}x$ as a perfect square, then subtracting whatever that added. It is the most useful step in the whole topic.

EXAMPLE 4.14

Write $y = x^2 - 6x + 5$ in vertex form, and state its vertex and axis of symmetry.

Take the $x^2 - 6x$ part and complete the square. Half of -6 is -3 , and $(-3)^2 = 9$:

$$x^2 - 6x = (x - 3)^2 - 9.$$

So

$$y = (x - 3)^2 - 9 + 5 = (x - 3)^2 - 4.$$

The vertex is $(3, -4)$ and the axis of symmetry is $x = 3$. Because $a = 1 > 0$, this vertex is the minimum point.

EXAMPLE 4.15

Sketch $y = x^2 - 2x - 3$, showing the vertex, axis of symmetry, and all intercepts.

Factor for the roots: $x^2 - 2x - 3 = (x - 3)(x + 1)$, so the x -intercepts are $x = -1$ and $x = 3$.

The y -intercept is $c = -3$. Completing the square, $y = (x - 1)^2 - 4$, so the vertex is $(1, -4)$ and the axis of symmetry is $x = 1$, which is exactly halfway between the two roots.




Figure 4.3 $y = x^2 - 2x - 3$, with its vertex, axis of symmetry and intercepts.

YOUR TURN 4.4 Quadratic Functions

19. Write each quadratic in vertex form and state its vertex.

(a) $y = x^2 + 4x + 1$

(b) $y = x^2 - 4x$

(c) $y = x^2 - 6x + 11$

(d) $y = x^2 + 2x + 5$

20. State the vertex and the axis of symmetry of each quadratic.

(a) $y = (x - 2)^2 + 5$

(b) $y = x^2 - 8x + 3$

(c) $y = -(x + 1)^2 + 2$

(d) $y = 2(x - 3)^2 - 1$

21. Consider $y = (x - 1)(x + 5)$.

(a) Write down its roots.

(b) Find its y -intercept.

(c) Find its axis of symmetry.

(d) Find its vertex.

22. State whether each quadratic has a maximum or a minimum, and give a reason.

(a) $y = -x^2 + 2x$

(b) $y = 3x^2 - x + 1$

4.5 Quadratic Equations

Setting a quadratic equal to zero is the same as asking where its parabola crosses the x -axis. There are three ways to find those crossings, and all three give the same answers. Choosing the right method saves time in an examination.

Method	Use it when	Be careful with
Factorisation	the quadratic factorises with whole numbers	many quadratics do not factorise, and searching for factors takes time
Completing the square	you also want the vertex, or an exact surd answer	the arithmetic is harder when $a \neq 1$
The formula	always. It works for every quadratic	slower than factorising when factors are obvious

Solving by factorisation

If the quadratic factorises, you can read the roots immediately. The reason is the zero product rule: a product is zero only when one of its factors is zero.

EXAMPLE 4.16

Solve $x^2 - 5x + 6 = 0$.

Find two numbers that multiply to 6 and add to -5 . They are -2 and -3 :

$$(x - 2)(x - 3) = 0.$$

A product is zero only when a factor is zero, so $x - 2 = 0$ or $x - 3 = 0$, giving $x = 2$ or $x = 3$.

CAUTION The zero product rule needs a zero on the right. From $(x - 2)(x - 3) = 2$ you may not conclude $x - 2 = 2$. Expand, collect everything on one side, and factorise again.

Solving by completing the square

Completing the square works whether or not the quadratic factorises, and it gives exact answers in surd form. Rewrite the quadratic as a perfect square plus a constant, then take square roots of both sides.

EXAMPLE 4.17

Solve $x^2 + 8x + 7 = 0$ by completing the square.

Half of 8 is 4, and $4^2 = 16$, so $x^2 + 8x = (x + 4)^2 - 16$. The equation becomes

$$(x + 4)^2 - 16 + 7 = 0 \implies (x + 4)^2 = 9.$$

Take the square root of both sides, remembering both signs:

$$x + 4 = \pm 3 \implies x = -1 \text{ or } x = -7.$$

EXAMPLE 4.18

Solve $x^2 - 4x - 1 = 0$, giving exact answers.

Half of -4 is -2 , and $(-2)^2 = 4$:

$$(x - 2)^2 - 4 - 1 = 0 \implies (x - 2)^2 = 5 \implies x - 2 = \pm\sqrt{5}.$$

So $x = 2 \pm \sqrt{5}$. This quadratic does not factorise with whole numbers, which is why the answer contains a surd.

TIP When $a \neq 1$, divide the whole equation by a first. For $2x^2 - 8x + 5 = 0$, divide by 2 to get $x^2 - 4x + 2.5 = 0$, then complete the square as usual.

Solving by formula

The formula works for every quadratic, and it is in the formula booklet.

KEY · IB The solutions of $ax^2 + bx + c = 0$, with $a \neq 0$, are

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Substitute the three coefficients and the formula returns both roots at once.

The formula is not a rule to memorise without understanding. It is what you get by completing the square once, using letters instead of numbers. Start from $ax^2 + bx + c = 0$, divide by a , complete the square on $x^2 + \frac{b}{a}x$, then solve. The $\pm$ appears because the parabola is symmetric, with one root on each side of the axis.

EXAMPLE 4.19

Solve $2x^2 - 3x - 2 = 0$.

Here $a = 2$, $b = -3$, $c = -2$. The formula gives

$$x = \frac{3 \pm \sqrt{9 + 16}}{4} = \frac{3 \pm 5}{4},$$

so $x = 2$ or $x = -\frac{1}{2}$.

YOUR TURN 4.5 Quadratic Equations

23. Solve each equation by factorisation.

(a) $x^2 - 5x + 6 = 0$

(b) $x^2 - x - 12 = 0$

(c) $x^2 + x - 20 = 0$

(d) $2x^2 - 5x - 3 = 0$

24. Solve each equation by completing the square, giving exact answers.

(a) $x^2 + 8x + 7 = 0$

(b) $x^2 - 2x - 5 = 0$

(c) $x^2 + 6x + 2 = 0$

(d) $2x^2 - 8x + 5 = 0$ (Hint: divide by 2 first)

25. Solve each equation with the quadratic formula, giving exact answers.

(a) $x^2 + 2x - 1 = 0$

(b) $x^2 + 6x + 4 = 0$

(c) $2x^2 - 3x - 2 = 0$

(d) $3x^2 + 2x - 1 = 0$

26. For each equation, state which of the three methods you would choose, and give a reason. You do not need to solve the equations.

(a) $x^2 - 7x + 12 = 0$

(b) $x^2 + 4x - 2 = 0$

(c) $5x^2 - 3x - 1 = 0$

(d) $x^2 - 9 = 0$

4.6 The Discriminant

The quadratic formula answers more than it is asked. Before you calculate a single root, the quantity under the square root sign already tells you how many real roots there are. That quantity has its own name.

KEY · IB The **discriminant** of $ax^2 + bx + c$ is

$$\Delta = b^2 - 4ac.$$

Its sign tells you how many real roots the equation has, without solving it.

The sign of Δ determines how many times the parabola meets the x -axis. The reason is that a negative number has no real square root, so $\pm\sqrt{\Delta}$ has no real value.

Discriminant	Real roots	Parabola and x -axis
$\Delta > 0$	two distinct	crosses twice
$\Delta = 0$	one repeated	touches once
$\Delta < 0$	none	does not meet it

EXAMPLE 4.20

Find the values of k for which $x^2 + kx + 9 = 0$ has exactly one (repeated) root.

One repeated root means $\Delta = 0$:

$$k^2 - 4(1)(9) = 0 \implies k^2 = 36 \implies k = \pm 6.$$

At these two values the parabola touches the x -axis at one point. Notice that the discriminant answers a question about the roots without finding them.

EXAMPLE 4.21

For the equation $3kx^2 + 2x + k = 0$, find the values of k that give two distinct real roots, two equal real roots, and no real roots.

Here the unknown constant appears in two places: $a = 3k$ and $c = k$, with $b = 2$. So

$$\Delta = b^2 - 4ac = 4 - 4(3k)(k) = 4 - 12k^2.$$

Now consider the three cases separately.

$$\Delta > 0 : 4 - 12k^2 > 0 \implies k^2 < \frac{1}{3} \implies -\frac{1}{\sqrt{3}} < k < \frac{1}{\sqrt{3}}.$$

$$\Delta = 0 : k^2 = \frac{1}{3} \implies k = \pm \frac{1}{\sqrt{3}}.$$

$$\Delta < 0 : k^2 > \frac{1}{3} \implies k < -\frac{1}{\sqrt{3}} \text{ or } k > \frac{1}{\sqrt{3}}.$$

One value of k must be checked separately. If $k = 0$ the equation becomes $2x = 0$. That is linear, not quadratic, and it has the single root $x = 0$. So the answer for two distinct real roots is $-\frac{1}{\sqrt{3}} < k < \frac{1}{\sqrt{3}}$, with $k \neq 0$.

CAUTION Before using the discriminant, check that the equation really is quadratic. If the coefficient of x^2 contains the unknown constant, the value that makes it zero must be excluded and treated separately.

TIP A question that says “two distinct roots”, “equal roots”, “a repeated root”, “touches the x -axis”, or “no real solutions” is asking about the discriminant. Recognising the wording tells you which method to use.

YOUR TURN 4.6 The Discriminant

27. Evaluate the discriminant and state the nature of the roots.

(a) $x^2 - 4x + 4$

(b) $2x^2 + 3x + 5$

(c) $3x^2 - 2x - 1$

(d) $x^2 + 5x + 7$

28. Find the value or values of k in each case.

(a) $x^2 + 6x + k = 0$ has equal roots

(b) $x^2 + kx + 16 = 0$ has equal roots

(c) $x^2 - 3x + k = 0$ has no real roots

(d) $kx^2 + 4x + k = 0$ has equal roots. State any value of k that must be excluded.

29. The equation $x^2 + (k + 1)x + 9 = 0$ has two distinct real roots. Find the set of values of k .

4.7 Quadratic Inequalities

A quadratic **inequality** is a different question. It does not ask where the parabola meets the axis. It asks where the parabola lies above the axis, or below it. The method is always the same three steps.

KEY To solve a quadratic inequality:

1. Collect every term on one side, so the other side is 0.
2. Find the roots of the matching equation. These are the only places where the sign can change.
3. Sketch the parabola and read the intervals where it lies on the side you want.

EXAMPLE 4.22

Solve $x^2 - x - 6 > 0$.

Factor: $x^2 - x - 6 = (x - 3)(x + 2)$, so the roots are $x = -2$ and $x = 3$. The parabola opens upward. It is therefore *above* the axis outside the roots, and below the axis between them. So

$$x^2 - x - 6 > 0 \quad \text{when} \quad x < -2 \quad \text{or} \quad x > 3.$$

A rough sketch of the parabola is faster and safer than memorising sign rules.

EXAMPLE 4.23

Solve $2x^2 \leq x + 6$.

First collect everything on one side:

$$2x^2 - x - 6 \leq 0.$$

Factorise: $(2x + 3)(x - 2) = 0$ gives roots $x = -\frac{3}{2}$ and $x = 2$. The parabola opens upward, so it lies *below* the axis between the roots. The sign $\leq$ includes the roots themselves:

$$-\frac{3}{2} \leq x \leq 2.$$

CAUTION *Never divide an inequality by a term containing x . Its sign is unknown, and dividing by a negative quantity reverses the inequality. Collect on one side and factorise instead.*

EXAMPLE 4.24

Solve $x^2 + x + 1 > 0$.

The discriminant is $1 - 4 = -3 < 0$, so the parabola never meets the x -axis. Since $a = 1 > 0$ it opens upward, so it lies entirely above the axis. The inequality holds for *every* real x .

When the discriminant is negative, a quadratic inequality has either every real number as its answer or none at all. The sign of a decides which.

YOUR TURN 4.7 Quadratic Inequalities

30. Solve each inequality.

(a) $x^2 - 9 < 0$

(b) $x^2 + 2x - 15 < 0$

(c) $x^2 - 4x + 3 \geq 0$

(d) $-x^2 + 4x > 0$

31. Solve each inequality. Two of them are special cases, so justify your answers.

(a) $2x^2 \leq x + 6$

(b) $x^2 \geq 16$

(c) $x^2 + x + 1 > 0$

(d) $x^2 + 4 < 0$

32. A rectangle has width x and length $x + 2$, both in centimetres.(a) Show that its area is $x^2 + 2x$.(b) Find the values of x for which the area is less than 15.

Chapter 4 · Lines & Quadratics

Reflections

TOK The same parabola appears in pure algebra, as the graph of x^2 , and in the physical world, as the path of every thrown object, the cross-section of a satellite dish, and the cable of certain bridges. Why should a curve defined by an abstract equation also describe falling stones? This is what the physicist Eugene Wigner called the “unreasonable effectiveness of mathematics.” Is mathematics invented to fit the world, or discovered as something already present in it?

IM Quadratics are among the oldest mathematics we have. Babylonian clay tablets from around 2000 BCE solve them to divide land and dig canals. The systematic method is younger: around 820 CE in Baghdad, al-Khwārizmī wrote *al-Kitāb al-mukhtaṣar fī ḥisāb al-jabr*, completing the square with geometric squares and rectangles. From its title, *al-jabr* (“restoring”), comes the word *algebra*, and from his name comes *algorithm*.

RW A parabola reflects every incoming parallel ray to a single point, called the focus. Satellite dishes and radio telescopes use this to collect weak signals at one point. Car headlights and torches use it in reverse: a bulb at the focus produces a parallel beam. Solar concentrators use it to focus sunlight and boil water. Quadratics also model optimisation everywhere: the maximum area for a fixed fence, the price that maximises profit, the angle at which a ball travels farthest.

EXT The discriminant divides quadratics into three cases. When $\Delta < 0$ the parabola does not meet the x -axis and there are no *real* roots, but the quadratic formula still returns an-

swers, now containing $\sqrt{-1}$. Keeping these “impossible” numbers instead of rejecting them leads to the **complex numbers** of Chapter 18, where every quadratic has exactly two roots and the discriminant only tells you whether those roots are real.

End-of-Chapter Problems — Chapter 4: Lines & Quadratics

1. Consider $f(x) = 2x - 1$.

(a) Find $f^{-1}(x)$. [2]

(b) Verify that $f(f^{-1}(x)) = x$. [2]

2. Consider $f(x) = 3 - 2x$.

(a) State the x - and y -intercepts. [2]

(b) Find $f^{-1}(x)$. [2]

3. Consider $f(x) = \frac{2x+1}{x-3}$, $x \neq 3$.

(a) State the domain. [1]

(b) Find $f^{-1}(x)$. [4]

4. Find the equation of the line through $(3, -1)$ parallel to $2x - y = 4$. [3]

5. Consider the points $A(-1, 2)$ and $B(3, 10)$.

(a) Find the midpoint of AB . [1]

(b) Find the gradient of AB . [1]

(c) Find the length of AB . [2]

6. The line $2x + 3y = 12$ is given.

(a) Find its x - and y -intercepts. [2]

(b) State its gradient. [2]

7. A line passes through $A(1, 2)$ and $B(5, 4)$.

(a) Find the gradient of AB . [1]

(b) Find the equation of AB . [2]

(c) Find the equation of the perpendicular bisector of AB . [4]

8. Solve the system $3x + 2y = 7$, $x - 2y = -3$. [3]

9. Solve the system

$$x + 2y + z = 4, \quad 2x + y - z = 2, \quad x - y + 2z = 2.$$

[6]

10. Consider the system

$$x + y + z = 3, \quad x + 2y + 3z = 4, \quad 2x + 3y + 4z = k.$$

- (a) Find the value of k for which the system has solutions. [3]
- (b) For this value of k , find the general solution. [3]
- (c) Describe the configuration of the three planes when $k = 10$. [1]

11. A quadratic has roots 2 and -5 .

- (a) Write it in factored form (with leading coefficient 1). [1]
- (b) Expand it into standard form. [2]

12. A parabola passes through $(0, 3)$, $(1, 0)$ and $(3, 0)$. Find its equation. [5]

13. Consider $f(x) = x^2 + 2x - 3$.

- (a) Factorise $f(x)$ and state its roots. [2]
- (b) Find the vertex. [2]
- (c) Sketch the graph, showing all intercepts. [2]

14. Consider $f(x) = x^2 - 6x + 5$.

- (a) Write $f(x)$ in vertex form. [2]
- (b) State the vertex and the axis of symmetry. [2]
- (c) Find the x - and y -intercepts. [3]

15. Solve the inequality $x^2 - 2x - 8 \leq 0$. [3]

16. The equation $x^2 - 4x + c = 0$ has no real roots. Find the set of values of c . [3]

17. Solve the inequality $x^2 + x - 6 > 0$. [3]

18. The equation $x^2 + kx + 4 = 0$ has two distinct real roots. Find the set of values of k . [4]

19. Find the values of k for which $x^2 + kx + (k + 3) = 0$ has equal roots. [4]

20. Two numbers have a sum of 10 and a product of 21. Find the numbers. [4]

21. The roots of $x^2 + bx + c = 0$ sum to 6 and differ by 4. Find b and c . [5]

22. Show that $x^2 + (k+1)x + k = 0$ has real roots for every real k , and find the value of k for which the roots are equal. [5]

23. The roots of $x^2 - px + q = 0$ are α and 2α . Show that $2p^2 = 9q$. [5]

24. Find the points of intersection of $y = x^2$ and $y = 2x + 3$. [4]

25. Consider the line $y = 2x - 5$ and the parabola $y = x^2 - 2x + 2$.

(a) Show that they do not intersect. [3]

(b) Explain what this means geometrically. [1]

26. A ball is thrown so that its height in metres after t seconds is $h(t) = 20t - 5t^2$.

(a) Find the time at which the ball lands. [2]

(b) Find the maximum height reached. [3]

27. Find the value of k for which the line $y = x + k$ is a tangent to the parabola $y = x^2$. [5]

28. A rectangle has length 3 more than its width w .

(a) Write its area A as a function of w . [2]

(b) Given the area is 40, find w . [3]

29. A rectangle has perimeter 20. Show that its area is $A = x(10 - x)$, where x is one side, and find the maximum possible area. [5]

30. Prove that the line $y = mx + c$ is a tangent to the parabola $y = x^2$ if and only if $c = -\frac{m^2}{4}$. [6]

31. GDC Consider $f(x) = x^3 - 4x + 1$.

(a) Using your GDC, sketch the graph of f for $-3 \leq x \leq 3$, labelling all key features. [2]

(b) Find the zeros of f , giving your answers to 3 significant figures. [2]

(c) Find the coordinates of the local maximum and local minimum points. [2]

(d) Solve $f(x) = 2x - 1$, giving your answers to 3 significant figures. [2]

32. A bridge has a parabolic arch. The arch is 40 m wide at road level and 10 m high at its centre. Take the road as the x -axis, with the centre of the arch at $x = 0$.

(a) Show that the arch is modelled by $y = 10 - \frac{x^2}{40}$. [4]

(b) Find the height of the arch 12 m from the centre. [2]

- (c) A lorry 8 m wide and 9.5 m high drives along the centre of the road. Determine whether it passes under the arch. [4]
- (d) Find the greatest width of a lorry 9 m high that can pass under the arch, giving your answer to three significant figures. [4]
- (e) The model gives a negative value of y when $|x| > 20$. Explain what this means for the model. [2]

33. Consider the line $y = mx + 3$ and the parabola $y = x^2 + 5x + 7$.

- (a) Show that at any point of intersection, $x^2 + (5 - m)x + 4 = 0$. [3]
- (b) Find the values of m for which the line is a tangent to the parabola. [4]
- (c) Find the values of m for which the line does not meet the parabola. [3]
- (d) For the smaller value of m found in part (b), find the coordinates of the point of tangency. [3]
- (e) State the values of m for which the line meets the parabola twice. [2]

34. **HL** Consider the system of equations

$$x + y + z = 2, \quad x + 2y + 3z = 5, \quad 2x + 3y + az = b,$$

where a and b are real constants.

- (a) Write the system as an augmented matrix and row reduce it. [4]
- (b) Find the value of a for which the system does not have a unique solution. [3]
- (c) For that value of a , find the value of b for which the system is consistent. [2]
- (d) For those values of a and b , find the general solution of the system. [4]
- (e) Describe how the three planes are arranged in each of the three cases: a not equal to the value in part (b); a equal to it with b consistent; and a equal to it with b not consistent. [3]

Challenge Questions

35. Tangents to a parabola. A line is a tangent to a curve when it meets the curve exactly once. This investigation turns that idea into a condition on the discriminant, then uses it to count the tangents from a point.

- (a) Show that the line $y = mx + c$ meets $y = x^2$ where $x^2 - mx - c = 0$, and hence that the line is a tangent exactly when $c = -\frac{m^2}{4}$. [4]
- (b) Find the two tangents to $y = x^2$ that pass through the point $(0, -4)$, and state where each one touches the parabola. [4]
- (c) Repeat part (a) for the parabola $y = ax^2$, with $a \neq 0$, and show that the tangency condition becomes $c = -\frac{m^2}{4a}$. [3]

- (d) Using part (c), find how many tangents to $y = x^2$ pass through the point $(0, k)$, for each value of k . State the three cases and explain each one from the picture. [5]
- (e) A student claims that every point in the plane has either two tangents to $y = x^2$ or none. Decide whether the claim is true, and justify your answer. [4]

36. A family of planes. Two planes that meet do so in a line. This investigation shows that infinitely many planes contain that same line, and uses the idea to decide when a third equation is consistent.

- (a) The planes $P_1 : x + y + z = 1$ and $P_2 : x - y + 2z = 2$ meet in a line. Find that line in the form (x, y, z) in terms of one parameter. [4]
- (b) For any constant λ , the equation $P_1 + \lambda P_2 = 0$ describes a plane. Show that every point of the line from part (a) lies on it, whatever λ is. [3]
- (c) Find the value of λ that produces the plane $2x + 3z = 3$, and hence explain why the system $P_1, P_2, 2x + 3z = 3$ has infinitely many solutions. [4]
- (d) Explain why the system $P_1, P_2, 2x + 3z = k$ has no solution for every $k \neq 3$. Reduce the augmented matrix to support your answer. [4]
- (e) A third plane is chosen at random. Which of the three outcomes in §4.3 is it most likely to produce, and why are the other two special? [3]

37. Sum and product of roots. A quadratic can be read from its roots without ever solving it. This investigation builds the relationships and then uses them to answer questions that would be slow by the formula.

- (a) Let the roots of $x^2 - Sx + P = 0$ be α and β . By expanding $(x - \alpha)(x - \beta)$, show that $\alpha + \beta = S$ and $\alpha\beta = P$. [3]
- (b) Show that $(\alpha - \beta)^2 = S^2 - 4P$, and hence write down the condition on S and P for the roots to be real. [3]
- (c) The roots of a quadratic sum to 6 and differ by 4. Use part (b) to find the quadratic, without finding the roots first. [3]
- (d) Show that $\alpha^2 + \beta^2 = S^2 - 2P$ and that $\frac{1}{\alpha} + \frac{1}{\beta} = \frac{S}{P}$. Verify both for $x^2 - 6x + 5 = 0$. [4]
- (e) Find the condition on S and P for the two roots to be reciprocals of each other, and give one quadratic with that property and irrational roots. [3]

