

Sequences & Series

2

SYLLABUS 1.2 1.3 1.4 1.8 1.9 1.10

A quantity that keeps doubling grows slowly at first, and then very fast. A sheet of paper is about 0.1 mm thick. Each fold doubles the thickness, so 42 folds would give 0.1×2^{42} mm, which is about 440,000 km. The Moon is about 384,000 km away. Nobody can fold paper 42 times, but the calculation is correct.

One more fold would reach the Moon and come back. Can you see why?

This chapter is about lists of numbers built by repeatedly adding a fixed amount, and lists built by repeatedly multiplying by a fixed factor. You will learn to find any term without writing out the terms before it, to add the first n terms in a single calculation, and to see why some sums that never end still have a finite total. The same ideas are used to value a savings plan, and, with a counting argument, to expand a bracket like $(a + b)^{12}$ without multiplying out twelve brackets.

2.1 Arithmetic Sequences

A theatre has 20 seats in the front row, 23 in the second, and 26 in the third. Each row has 3 more seats than the one before it. To find how many seats are in row 50, you should not have to count up one row at a time. There is a clear pattern: each row adds 3 seats. A regular pattern like this can be written as a formula, and finding that formula is the point of this section.

A sequence where each term is obtained by adding a fixed constant to the previous one is called **arithmetic** (also called an **arithmetic progression**, or **AP**). That constant is the **common difference** d .

DEF An **arithmetic sequence** has the form $u_1, u_1 + d, u_1 + 2d, \dots$, where u_1 is the first term and d is the common difference.

For any two consecutive terms: $d = u_{n+1} - u_n$.

The general term

The n -th term is the first term plus $(n - 1)$ steps of size d . Starting from u_1 , you take one step to reach u_2 , two steps to reach u_3 , and so on.

KEY · IB

$$u_n = u_1 + (n - 1)d$$

EXAMPLE 2.1

An arithmetic sequence has first term 7 and common difference -3 . Find u_{20} .

You need 19 steps of size -3 from $u_1 = 7$:

$$u_{20} = 7 + (20 - 1)(-3) = 7 - 57 = -50.$$

A negative common difference means the sequence is decreasing. The general term can be negative even when the first term is positive.

EXAMPLE 2.2

In an arithmetic sequence, $u_5 = 11$ and $u_9 = 27$. Find u_1 and d .

From u_5 to u_9 is exactly 4 steps:

$$4d = u_9 - u_5 = 27 - 11 = 16 \implies d = 4.$$

Then $u_1 = u_5 - 4d = 11 - 16 = -5$.

EXAMPLE 2.3

Find the number of terms in the arithmetic sequence $5, 9, 13, \dots, 101$.

Here $d = 4$. Set $u_n = 101$:

$$5 + (n - 1)(4) = 101 \implies 4(n - 1) = 96 \implies n = 25.$$

CAUTION The formula is $u_1 + (n - 1)d$, not $u_1 + nd$. For the first term ($n = 1$), you add zero differences, not one.

YOUR TURN 2.1 Arithmetic Sequences**1.** Find the term indicated.

(a) u_7 of 3, 7, 11, ...

[1] (b) u_{15} of the AP with $u_1 = 12$, $d = -4$
[1]

(c) u_{10} of $-5, -1, 3, \dots$

[1] (d) u_{25} of the AP with $u_1 = 100$,
 $d = -7$ [2]

2. Find the quantity indicated.

(a) d , given $u_1 = 5$ and $u_9 = 37$

[2] (b) u_1 , given $u_6 = 23$ and $d = 3$ [2]

(c) d , given $u_3 = 8$ and $u_{11} = 32$

[2] (d) u_1 , given $u_{12} = 50$ and $d = 4$ [2]

3. Find the number of terms in the AP 4, 7, 10, ..., 100.

[2]

4. Given $u_4 = 11$ and $u_{10} = 29$, find u_1 and d .

[3]

5. Show that the sequence 5, 11, 17, ... has a term equal to 101. State which term it is.

[3]

2.2 Arithmetic Series

When you need the total of the first n terms of an arithmetic sequence, adding them one at a time is always possible but rarely necessary. A pattern reduces the whole sum to one short calculation.

Consider adding all integers from 1 to 100. Write the sum forwards and backwards:

$$1 + 2 + 3 + \dots + 99 + 100$$

$$100 + 99 + 98 + \dots + 2 + 1$$

Every vertical pair sums to 101, and there are 100 such pairs. So twice the sum equals 100×101 , giving $S_{100} = 5050$.

The same pairing works for any arithmetic series: the first and last term always sum to the same value, $u_1 + u_n$. There are n such pairs when you write the series twice, so $2S_n = n(u_1 + u_n)$, and therefore:

KEY · IB

$$S_n = \frac{n}{2}(u_1 + u_n) = \frac{n}{2}(2u_1 + (n-1)d)$$

The first form is faster when you know u_n . Substituting $u_n = u_1 + (n-1)d$ gives the second, which works when only u_1 and d are known.

EXAMPLE 2.4

Find the sum of the first 30 terms of the arithmetic sequence 2, 7, 12, ...

Here $u_1 = 2$ and $d = 5$. The last term is not given, so use the second form:

$$S_{30} = \frac{30}{2}(2(2) + 29(5)) = 15(4 + 145) = 15 \times 149 = 2235.$$

EXAMPLE 2.5

Find the sum $4 + 7 + 10 + \dots + 100$.

First, $d = 3$. Find the number of terms: $4 + (n-1)(3) = 100 \Rightarrow n = 33$.

Then use the first form, since both u_1 and u_{33} are known:

$$S_{33} = \frac{33}{2}(4 + 100) = \frac{33 \times 104}{2} = 1716.$$

EXAMPLE 2.6

The sum of the first n terms of an arithmetic series is given by $S_n = 3n^2 + 5n$. Find u_1 , d , and the general term u_n .

$$u_1 = S_1 = 3 + 5 = 8.$$

$$u_2 = S_2 - S_1 = (12 + 10) - 8 = 14, \text{ so } d = u_2 - u_1 = 6.$$

$$\text{For the general term: } u_n = S_n - S_{n-1} = 3n^2 + 5n - 3(n-1)^2 - 5(n-1).$$

$$\text{Expanding: } u_n = 3n^2 + 5n - 3n^2 + 6n - 3 - 5n + 5 = 6n + 2.$$

Check: $u_1 = 8$ and $u_2 = 14$. Consistent.

CAUTION S_n is the total of the first n terms; u_n is just the n -th term. Confusing the two is the most common error in this topic. A question asking for a "term" wants u_n ; a question asking for a "sum" wants S_n .

Application: simple interest

The standard financial application of an arithmetic sequence is **simple interest**: the same fixed amount of interest is added each year, so the balance increases by the same amount each year.

EXAMPLE 2.7

\$2000 is deposited in an account paying 3% simple interest per year. Find the balance after 15 years.

The interest is $3\% \times 2000 = \$60$ every year, so the balances form an arithmetic sequence with $d = 60$. After 15 years, 15 payments of interest have been added:

$$2000 + 15 \times 60 = \$2900.$$

YOUR TURN 2.2 Arithmetic Series

6. Find the sum indicated.

(a) S_{12} for the AP 2, 5, 8, ...

[2]

(b) S_{20} for the AP with $u_1 = 100$,
 $d = -5$

[2]

(c) S_{15} for the AP 7, 11, 15, ...

[2]

(d) S_{25} for the AP with $u_1 = -12$, $d = 5$

[2]

7. Find the sum $6 + 10 + 14 + \dots + 90$.

[3]

8. Find the sum of all integers from 1 to 200 that are multiples of 6.

[3]

9. Find n if $S_n = 234$ for the AP 3, 6, 9, ...

[3]

10. An AP has $u_1 = 15$, $u_n = -9$ and $n = 13$. Find S_n .

[2]

11. Given $S_n = 4n^2 - n$, find u_1 , d and the general term u_n .

[4]

2.3 Geometric Sequences

A geometric sequence multiplies by the same number at each step. Start at 1 and double: 1, 2, 4, 8, 16, ... Many real quantities change in this way. Bacteria double in every generation. A ball bounces to 60% of the height it fell from. Light loses half its intensity for each extra filter. In every case, each new value is the previous value multiplied by a fixed number.




Figure 2.1 Adding a fixed amount grows in a straight line; multiplying by a fixed factor grows explosively.

A sequence where each term is obtained by multiplying the previous term by a fixed constant is called **geometric** (also called a **geometric progression**, or **GP**). That constant is the **common ratio** r .

DEF A **geometric sequence** has the form $u_1, u_1r, u_1r^2, \dots$, where u_1 is the first term and r is the common ratio.

For any two consecutive terms: $r = \frac{u_{n+1}}{u_n}$.

The general term

Starting from u_1 , each step multiplies by r . After $(n - 1)$ multiplications you reach u_n :

KEY · IB

$$u_n = u_1 \cdot r^{n-1}$$

EXAMPLE 2.8

A geometric sequence has $u_1 = 5$ and $r = 3$. Find u_6 .

$$u_6 = 5 \cdot 3^5 = 5 \times 243 = 1215.$$

EXAMPLE 2.9

A geometric sequence has $u_3 = 12$ and $u_6 = 96$. Find u_1 and r .

From u_3 to u_6 is 3 multiplicative steps: $u_6 = u_3 \cdot r^3$, so

$$r^3 = \frac{96}{12} = 8 \implies r = 2.$$

Then $u_1 = u_3 / r^2 = 12 / 4 = 3$.

EXAMPLE 2.10

Find the number of terms in the geometric sequence 2, 6, 18, ..., 4374.

Here $r = 3$. Set $u_n = 4374$:

$$2 \cdot 3^{n-1} = 4374 \implies 3^{n-1} = 2187 = 3^7 \implies n = 8.$$

TIP To test whether a sequence is arithmetic, check that differences are constant. To test whether it is geometric, check that ratios are constant. Never apply both tests to the same sequence expecting both to be true.

YOUR TURN 2.3 Geometric Sequences

12. Find the term indicated.

- | | | | |
|---|-----|---|-----|
| (a) u_6 of the GP 2, 6, 18, ... | [1] | (b) u_7 of the GP with $u_1 = 96$, $r = \frac{1}{2}$ | [2] |
| (c) u_5 of $\frac{1}{4}, \frac{1}{2}, 1, \dots$ | [2] | (d) u_8 of the GP with $u_1 = 3$, $r = -2$ | [2] |

13. Find the quantity indicated.

- | | | | |
|--|-----|--|-----|
| (a) r , given $u_1 = 5$ and $u_4 = 40$ | [2] | (b) u_1 , given $u_4 = 24$ and $r = 2$ | [2] |
| (c) r , given $u_1 = 81$ and $u_5 = 1$ | [3] | (d) u_1 , given $u_3 = 45$ and $r = 3$ | [2] |

14. Find the number of terms in the GP 3, 6, 12, ..., 768. [2]

15. Given $u_2 = 10$ and $u_5 = 1250$, find u_1 and r . [3]

16. The 3rd term of a GP is 12 and the 6th term is 96. Find the first term. [3]

2.4 Geometric Series

Adding the terms of a geometric sequence needs a different method. In an arithmetic series, the first and last terms have the same total as the second and second-to-last terms, so pairing works. In a geometric series these pair totals are all different, so pairing does not help. Instead there is an algebraic method: multiply the sum by r , line the two sums up, and subtract.

Let $S = u_1 + u_1r + u_1r^2 + \dots + u_1r^{n-1}$. Multiply by r :

$$rS = u_1r + u_1r^2 + \dots + u_1r^n.$$

Every term appears in both lines except u_1 in the first line and $u_1 r^n$ in the second. Subtract rS from S :

$$S(1 - r) = u_1 - u_1 r^n = u_1(1 - r^n).$$

Divide both sides by $(1 - r)$ to get the formula.

KEY · IB

$$S_n = \frac{u_1(1 - r^n)}{1 - r} = \frac{u_1(r^n - 1)}{r - 1}, \quad r \neq 1$$

Both forms are equivalent. When $0 < r < 1$, use the first form: both $1 - r^n$ and $1 - r$ are then positive, so no negative signs appear. When $r > 1$ the second form is tidier, for the same reason. If r is negative, either form works, but r^n changes sign with n , so take care with the arithmetic.

EXAMPLE 2.11

Find the sum of the first 10 terms of the geometric series $3 + 6 + 12 + \dots$

Here $u_1 = 3$ and $r = 2$:

$$S_{10} = \frac{3(2^{10} - 1)}{2 - 1} = 3 \times 1023 = 3069.$$

EXAMPLE 2.12

A geometric series has $u_1 = 80$ and $r = \frac{1}{2}$. Find the least value of n such that $S_n > 159$.

$$S_n = \frac{80(1 - (\frac{1}{2})^n)}{1 - \frac{1}{2}} = 160\left(1 - \frac{1}{2^n}\right).$$

Require $160(1 - \frac{1}{2^n}) > 159$, so $\frac{1}{2^n} < \frac{1}{160}$, giving $2^n > 160$.

Since $2^7 = 128 < 160$ and $2^8 = 256 > 160$, the least value is $n = 8$.

CAUTION The formula requires $r \neq 1$. If $r = 1$, every term equals u_1 and $S_n = nu_1$. Substituting $r = 1$ into the formula gives $0/0$, which is undefined.

Solving for n in geometric series often requires taking logarithms of both sides. See Ch. 1, §1.3.

YOUR TURN 2.4 Geometric Series

17. Find the sum indicated.

- | | | | |
|---|-----|--|-----|
| (a) S_8 for the GP $1, 3, 9, \dots$ | [2] | (b) S_{10} for the GP with $u_1 = 6$, $r = \frac{1}{3}$ | [2] |
| (c) S_6 for the GP $128, 64, 32, \dots$ | [2] | (d) S_6 for the GP with $u_1 = 1$, $r = -2$ | [2] |

18. The sum of the first three terms of a GP is 21 and the common ratio is 2. Find the first term. [2]
19. Find n such that $S_n = 63$ for the GP 1, 2, 4, ... [2]
20. Find the sum $3 + 6 + 12 + \dots + 1536$. [2]
21. Given $S_3 = 7$ and $S_6 = 63$, find u_1 and r . [4]

2.5 Sigma Notation

By now you have written several sums in the form $u_1 + u_2 + \dots + u_n$. There is a compact notation for this that appears throughout mathematics, physics, and statistics, and it is worth learning well.

The Greek letter Σ (capital sigma) is used as shorthand for “sum”. The expression

$$\sum_{k=1}^n u_k = u_1 + u_2 + u_3 + \dots + u_n$$

is read “the sum of u_k from $k = 1$ to n .” The variable k is the **index of summation**: it takes each integer value from the lower limit to the upper limit, the corresponding term is evaluated, and all terms are added.

DEF $\sum_{k=m}^n f(k) = f(m) + f(m+1) + f(m+2) + \dots + f(n)$, where $m \leq n$.

The index k is a **dummy variable**: the result depends only on f , m , and n , not on the name chosen for the index.

Properties of sigma notation

These three properties are just familiar rules of arithmetic, written in sigma form. They are used constantly when simplifying or manipulating sums.

KEY For a constant c and sequences $\{a_k\}$, $\{b_k\}$:

$$\sum_{k=1}^n c = nc$$

a constant summed n times

$$\sum_{k=1}^n c a_k = c \sum_{k=1}^n a_k$$

a constant factor comes outside the sum

$$\sum_{k=1}^n (a_k + b_k) = \sum_{k=1}^n a_k + \sum_{k=1}^n b_k$$

a sum splits over addition

EXAMPLE 2.13

Write $3 + 7 + 11 + 15 + 19$ in sigma notation.

The terms form an arithmetic sequence with $u_1 = 3$, $d = 4$: the k -th term is $u_k = 4k - 1$.

$$3 + 7 + 11 + 15 + 19 = \sum_{k=1}^5 (4k - 1).$$

EXAMPLE 2.14

Evaluate $\sum_{k=1}^6 5 \cdot 2^{k-1}$.

This is a geometric series with $u_1 = 5$ and $r = 2$:

$$\sum_{k=1}^6 5 \cdot 2^{k-1} = S_6 = \frac{5(2^6 - 1)}{2 - 1} = 5 \times 63 = 315.$$

EXAMPLE 2.15

Evaluate $\sum_{k=3}^{10} (2k + 1)$.

The lower limit is 3, not 1. Use the fact that the full sum minus the missing part gives the partial sum:

$$\sum_{k=3}^{10} (2k + 1) = \sum_{k=1}^{10} (2k + 1) - \sum_{k=1}^2 (2k + 1).$$

The first sum is an arithmetic series: $u_1 = 3$, $u_{10} = 21$, so $S_{10} = \frac{10}{2}(3 + 21) = 120$.

The second sum is just $3 + 5 = 8$.

Result: $120 - 8 = 112$.

CAUTION $\sum_{k=1}^n (a_k \cdot b_k) \neq \left(\sum_{k=1}^n a_k\right) \left(\sum_{k=1}^n b_k\right)$. You can split a sum over addition but not over multiplication. This is one of the most common misapplications of sigma notation.

YOUR TURN 2.5 Sigma Notation

22. Write each sum in sigma notation.

(a) $5 + 10 + 15 + 20 + 25$ [1] (b) $4 + 8 + 16 + 32 + 64$ [1]

(c) $3 + 6 + 9 + \dots + 30$ [2] (d) $1 + 4 + 9 + 16 + 25$ [2]

23. Evaluate $\sum_{k=1}^6 (3k - 2)$. [2]

24. Evaluate $\sum_{k=1}^5 2 \cdot 3^{k-1}$. [2]

25. Evaluate $\sum_{k=4}^8 (2k + 1)$. [2]

26. Find $\sum_{k=1}^n 5$ and $\sum_{k=1}^n 5k$ using the sigma properties. [2]

2.6 Infinite Geometric Series

Cut a pizza in half. Eat one half and keep the other. Cut what is left in half again, eat one piece, and keep the rest. If you could continue forever, how much pizza would you eat?

The pieces you eat are $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$. There are infinitely many of them, and every piece is larger than zero. Adding infinitely many positive numbers sounds like it must give an infinite total. But you began with one pizza, so you can never eat more than one pizza. The total can only be 1.

That is the surprise of this section: *an addition that never ends can still have a finite answer.*

Add the pieces one at a time and record the running total after each step. Each running total is a **partial sum**:

$$S_1 = 0.5, \quad S_2 = 0.75, \quad S_3 = 0.875, \quad S_4 = 0.9375, \quad \dots$$

Two things happen at once. Every partial sum is larger than the one before it, and every partial sum is still smaller than 1. Look at the distance from 1: it is 0.5, then 0.25, then 0.125. The distance halves at every step, so it never becomes zero, but it becomes as small as you wish. This value that the partial sums approach is called the **limit**, and here the limit is 1.




Figure 2.2 The partial sums of $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$ climb toward 1 but never pass it.

The pizza works because each piece is smaller than the piece before it. That is exactly what $|r| < 1$ means: every term is a fraction of the previous term, so the terms shrink toward zero.

Now change one number and the behaviour changes completely. Take $1 + 2 + 4 + 8 + \dots$, where $r = 2$. Each term is larger than the last, the partial sums are 1, 3, 7, 15, ..., and they pass every value you can name. There is no finite total.

A series whose partial sums approach a finite limit is said to **converge**. A series whose partial sums grow without any bound is said to **diverge**. A geometric series converges when $|r| < 1$ and diverges when $|r| \geq 1$.

To turn this into a formula, start from the finite sum and let n grow without bound. When $|r| < 1$, the power r^n shrinks toward 0:

$$S_n = \frac{u_1(1 - r^n)}{1 - r} \xrightarrow{n \rightarrow \infty} \frac{u_1(1 - 0)}{1 - r} = \frac{u_1}{1 - r}.$$

KEY · IB For a geometric series with $|r| < 1$:

$$S_\infty = \frac{u_1}{1 - r}$$

EXAMPLE 2.16

Find the sum to infinity of $6 + 4 + \frac{8}{3} + \dots$

The common ratio is $r = 4/6 = \frac{2}{3}$. Since $|r| < 1$ the series converges:

$$S_\infty = \frac{6}{1 - \frac{2}{3}} = \frac{6}{\frac{1}{3}} = 18.$$

EXAMPLE 2.17

Express $0.\overline{27} = 0.272727 \dots$ as an exact fraction.

Write as an infinite geometric series with $u_1 = \frac{27}{100}$ and $r = \frac{1}{100}$:

$$0.272727 \dots = \frac{27/100}{1 - 1/100} = \frac{27/100}{99/100} = \frac{27}{99} = \frac{3}{11}.$$

EXAMPLE 2.18

A geometric series has first term a and common ratio r . Given $S_\infty = 8$ and $u_2 = 2$, find a and r .

From S_∞ : $\frac{a}{1-r} = 8$, so $a = 8(1-r)$.

From $u_2 = ar = 2$: substitute a :

$$8(1-r)r = 2 \implies 8r^2 - 8r + 2 = 0 \implies 4r^2 - 4r + 1 = 0 \implies (2r - 1)^2 = 0.$$

So $r = \frac{1}{2}$, and $a = 8(1 - \frac{1}{2}) = 4$.

CAUTION Always verify $|r| < 1$ before applying the infinite sum formula. If $|r| \geq 1$, no finite sum to infinity exists.

YOUR TURN 2.6 Infinite Geometric Series

27. Find S_∞ for each series.

- | | | | |
|------------------------------------|-----|--|-----|
| (a) $12 + 4 + \frac{4}{3} + \dots$ | [2] | (b) the GP with $u_1 = 10$, $r = \frac{3}{5}$ | [2] |
| (c) $9 + 3 + 1 + \dots$ | [2] | (d) the GP with $u_1 = 8$, $r = -\frac{1}{4}$ | [2] |

28. Find the quantity indicated.

- | | | | |
|---|-----|--|-----|
| (a) u_1 , given $S_\infty = 20$ and $r = \frac{1}{5}$ | [2] | (b) r , given $u_1 = 6$ and $S_\infty = 9$ | [2] |
| (c) u_1 , given $S_\infty = 12$ and $r = \frac{1}{4}$ | [2] | (d) r , given $u_1 = 15$ and $S_\infty = 12$ | [2] |

29. Express $0.\overline{5} = 0.5555 \dots$ as an exact fraction. [2]

30. State whether $6 + 9 + 13.5 + \dots$ converges. Give a reason. [2]

31. A GP has $u_1 = 16$ and $u_3 = 4$. Find S_∞ . [3]

2.7 Financial Applications

A bank pays you 4% per year. In the first year you earn interest on your deposit. In the second year you earn interest on the deposit *and* on the interest from the first year. The amount added is different every year, so the balances do not form an arithmetic sequence. They form a geometric one: each year the balance is multiplied by 1.04.

This is **compound interest**, and it is why money left alone for a long time grows far more than most people expect. The balance after any number of years is simply a term of a geometric sequence.

Interest is often added more than once a year. If a present value PV earns interest at a nominal annual rate of $r\%$, compounded k times per year, then after n years:

KEY · IB

$$FV = PV \times \left(1 + \frac{r}{100k}\right)^{kn}$$

The quantity $\left(1 + \frac{r}{100k}\right)$ is the common ratio. The total number of compounding periods is kn .

EXAMPLE 2.19

\$5 000 is invested at 4% per year, compounded monthly. Find the value after 3 years.

Here $PV = 5000$, $r = 4$, $k = 12$, $n = 3$:

$$FV = 5000 \times \left(1 + \frac{4}{1200}\right)^{36} \approx \$5636.36.$$

EXAMPLE 2.20

How many complete years does it take for an investment to double at 6% per year, compounded annually?

Set $FV = 2 \cdot PV$:

$$(1.06)^n = 2 \implies n = \frac{\ln 2}{\ln 1.06} \approx 11.9.$$

After 12 complete years the investment has more than doubled.

TIP The Rule of 72: dividing 72 by the annual interest rate gives an approximate doubling time in years. At 6%, that is $72/6 = 12$ years, matching the exact calculation above.

Depreciation

Some things lose value instead of gaining it. A car, a laptop, or a machine may lose a fixed *percentage* of its value each year. This is called **depreciation**.

No new formula is needed. Use the same compound-interest formula with a **negative** rate r . A loss of 12% per year means $r = -12$, so the common ratio becomes

$$1 + \frac{r}{100} = 1 - \frac{12}{100} = 0.88,$$

a number less than 1. Multiplying by a ratio less than 1 at every step makes the values shrink, so the sequence of values is still geometric.

EXAMPLE 2.21

A car bought for \$28 000 loses 12% of its value each year. Find its value after 5 years.

Here $r = -12$ and $k = 1$, so each year the value is multiplied by 0.88:

$$FV = 28\,000 \times (0.88)^5 \approx \$14\,776.$$

After five years the car has lost nearly half its value. Notice that the value never reaches zero: each year removes 12% of a smaller amount than the year before.

Inflation and real value

There is a second geometric process working against your savings. **Inflation** is the yearly rise in prices. Interest increases the number in your account, while inflation reduces what that number can buy. If your account grows by 2% in a year and prices rise by 3% in the same year, you hold more money but you can buy less with it.

The **real value** of an investment is its future value adjusted for the price rises over the same period. It measures the gain in purchasing power rather than the gain in the number.

EXT Real value is not required by the syllabus, but it uses the same geometric reasoning. With interest producing a future value FV over n years, and prices rising at $i\%$ per year:

$$\text{real value} = \frac{FV}{\left(1 + \frac{i}{100}\right)^n}$$

Divide by the inflation factor because the same goods cost $\left(1 + \frac{i}{100}\right)^n$ times as much after n years.

EXAMPLE 2.22

\$10 000 is invested for 5 years at 4.5% per year, compounded annually. Prices rise at 2.5% per year. Find the future value and the real value of the investment.

The balance grows by a factor of 1.045 each year:

$$FV = 10\,000 \times (1.045)^5 \approx \$12\,461.82.$$

Now adjust for five years of price rises:

$$\text{real value} = \frac{12\,461.82}{(1.025)^5} \approx \$11\,014.43.$$

The account shows a gain of about \$2460, but in purchasing power the investor is only about \$1014 better off. An interest rate means little until you compare it with inflation.

TIP Your GDC's built-in Finance (TVM) solver handles compound interest, depreciation, and unknown rates or times directly: enter the known values of N , $I\%$, PV , FV , and the compounding frequency, and solve for the missing one. In examinations, state the values you entered.

Solving for n in financial problems requires exponential equations. See Ch. 1, §1.3.

YOUR TURN 2.7 Financial Applications

32. GDC Find the future value of each investment.

- | | |
|---|--|
| <p>(a) \$3 000 at 4% p.a., compounded annually, for 5 years [2]</p> | <p>(b) \$8 000 at 6% p.a., compounded monthly, for 3 years [2]</p> |
| <p>(c) \$500 at 3.6% p.a., compounded monthly, for 10 years [2]</p> | <p>(d) \$2 000 at 5% p.a., compounded quarterly, for 4 years [2]</p> |

33. GDC Find the annual interest rate if \$5 000 grows to \$5 828.84 in 3 years, compounded annually. [3]

34. GDC How many complete years does it take \$2 000 to double at 9% p.a., compounded annually? [3]

35. GDC Find the present value needed to accumulate \$15 000 in 8 years at 5% p.a., compounded quarterly. [3]

36. A car valued at \$25 000 depreciates at 15% per year. Find its value after 4 years. [2]

2.8 Counting Principles HL

Before the chapter's final result, a short look at a question that sounds simple: *how many ways?* The answer is what makes the binomial theorem work, and it starts with one principle.

If one choice can be made in m ways and a second, independent choice in n ways, then the pair of choices can be made in $m \times n$ ways. This is the **multiplication principle**, and it extends to more choices: a menu offering 4 starters, 5 mains, and 3 desserts allows $4 \times 5 \times 3 = 60$ different meals.

Arranging n distinct objects in a row is the principle applied n times: n choices for the first position, $n - 1$ for the second, and so on down to 1. The total is n **factorial**:

KEY

$$n! = n \times (n - 1) \times (n - 2) \times \cdots \times 2 \times 1, \quad 0! = 1$$

Permutations and combinations

Two standard questions arise when selecting r objects from n distinct ones. If the *order* of selection matters, the count is a **permutation**. If only the final collection matters, it is a **combination**.

The two counts are closely related. Suppose you choose 3 letters from A, B, C, D, E. The selection $\{A, B, C\}$ can be written in $3! = 6$ different orders: ABC, ACB, BAC, BCA, CAB, CBA. A permutation count treats these six as six different results. A combination count treats them as one. So the combination count is the permutation count divided by $r!$:

$$\binom{n}{r} = \frac{{}^nP_r}{r!}.$$

KEY · IB HL Selecting r from n distinct objects:

$${}^nP_r = \frac{n!}{(n-r)!} \quad (\text{ordered}), \quad \binom{n}{r} = {}^nC_r = \frac{n!}{r!(n-r)!} \quad (\text{unordered})$$

EXAMPLE 2.23

Eight runners compete. In how many ways can gold, silver, and bronze be awarded?

The three medals are distinguishable, so order matters:

$${}^8P_3 = 8 \times 7 \times 6 = 336.$$

EXAMPLE 2.24

A committee of 5 is chosen from 6 boys and 5 girls. How many committees contain exactly 2 girls?

Choose the girls and the boys separately, then multiply:

$$\binom{5}{2} \binom{6}{3} = 10 \times 20 = 200.$$

A committee has no internal order, so combinations are correct throughout, and the multiplication principle combines the two independent choices.

CAUTION Ask one question before every counting problem: *does order matter?* Medals, passwords, and rankings: yes, permutations. Committees, hands of cards, and subsets: no, combinations.

TIP Your GDC computes both: nPr and nCr . Arrangements with repeated identical objects and circular arrangements are not required in this course.

YOUR TURN 2.8 Counting Principles HL

37. In how many ways can 6 different books be arranged on a shelf? [1]
38. A 4-digit security code uses digits from 1 to 9 with no digit repeated. How many such codes are there? [2]
39. Find the number of ways of choosing a committee of 3 from 12 people. [1]
40. From 7 men and 6 women, find the number of ways of choosing a group of 2 men and 2 women. [2]
41. Solve $\binom{n}{2} = 45$. [3]

2.9 The Binomial Theorem

Expanding $(a + b)^2$ is quick. Expanding $(a + b)^3$ takes longer. Expanding $(a + b)^{12}$ by multiplying out twelve brackets is not realistic. Fortunately the answer follows a pattern, so no bracket multiplication is needed at all.

Look at the small cases:

$$(a + b)^1 = a + b, \quad (a + b)^2 = a^2 + 2ab + b^2, \quad (a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3.$$

Two features are regular. The power of a falls from n to 0 while the power of b rises from 0 to n , and in every term the two powers add to n . The coefficients form a row of **Pascal's triangle**, in which each entry is the sum of the two entries above it:

$$\begin{array}{ccccccc} & & & & 1 & & & \\ & & & & & 1 & & \\ & & & 1 & & 2 & & 1 \\ & & 1 & & 3 & & 3 & & 1 \\ & 1 & & 4 & & 6 & & 4 & & 1 \\ 1 & & 5 & & 10 & & 10 & & 5 & & 1 \end{array}$$

Counting explains why. Multiplying out $(a + b)^n$ means choosing a or b from each of the n brackets. A term $a^{n-r}b^r$ arises once for every way of choosing which r brackets contribute a b , and that count is exactly $\binom{n}{r}$. The coefficients of the expansion are the combinations of the previous section.

KEY · IB For $n \in \mathbb{N}$:

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$

$$\text{where } \binom{n}{r} = \frac{n!}{r!(n-r)!}.$$

Pascal's triangle is fastest for small n . When n is large, or when a question asks for one coefficient only, use the single term $\binom{n}{r}a^{n-r}b^r$ shown in the middle of the expansion above and choose the r you need.

TIP The booklet shows that single term inside the expansion, but it is worth naming it. The **general term** is

$$T_{r+1} = \binom{n}{r}a^{n-r}b^r,$$

so $r = 0$ gives the first term, $r = 1$ the second, and r gives term number $r + 1$. The subscript is $r + 1$ because r counts how many factors of b the term contains, and the first term contains none. In an examination, write down this term, find the r that produces the power you want,

then substitute.

EXAMPLE 2.25

Expand $(x + 2)^4$.

Row 4 of the triangle gives coefficients 1, 4, 6, 4, 1, with the powers of 2 increasing as the powers of x decrease:

$$(x + 2)^4 = x^4 + 4x^3(2) + 6x^2(4) + 4x(8) + 16 = x^4 + 8x^3 + 24x^2 + 32x + 16.$$

EXAMPLE 2.26

Find the coefficient of x^3 in $(2x - 3)^5$.

The general term is $\binom{5}{r}(2x)^{5-r}(-3)^r$. For x^3 , take $5 - r = 3$, so $r = 2$:

$$\binom{5}{2}(2x)^3(-3)^2 = 10 \times 8x^3 \times 9 = 720x^3.$$

The coefficient is 720. Keep the sign inside the bracket with $b = -3$: the $(-3)^r$ produces the alternating signs automatically.

EXAMPLE 2.27

Find the term independent of x in $\left(x^2 + \frac{2}{x}\right)^6$.

Track the power of x in the general term:

$$T_{r+1} = \binom{6}{r}(x^2)^{6-r}\left(\frac{2}{x}\right)^r = \binom{6}{r}2^r x^{12-3r}.$$

Independent of x means $12 - 3r = 0$, so $r = 4$:

$$\binom{6}{4}2^4 = 15 \times 16 = 240.$$

TIP $\binom{n}{r}$ can also be read from a GDC table. To solve $\binom{6}{r} = 20$, generate $\binom{6}{r}$ for $r = 0, \dots, 6$ and read off $r = 3$. Symmetry, $\binom{n}{r} = \binom{n}{n-r}$, halves the search.

YOUR TURN 2.9 The Binomial Theorem

42. Expand or write down the terms requested.

- (a) Expand $(x + 3)^4$. [2] (b) Write down the first three terms, in ascending powers of x , of $(1 + 2x)^7$. [2]
- (c) Expand $(2x - 1)^4$. [2] (d) Write down the first three terms, in ascending powers of x , of $(1 - 3x)^6$. [2]
43. Find the coefficient of x^2 in the expansion of $(3x - 2)^5$. [3]
44. Find the term independent of x in $\left(x + \frac{1}{x^2}\right)^9$. [3]
45. GDC Find the values of r for which $\binom{8}{r} = 56$. [2]
46. Find the middle term in the expansion of $(x - 2y)^6$. [3]

2.10 The Binomial Series HL

So far n has been a positive whole number, because $(a + b)^n$ meant n brackets multiplied together. But expressions such as $\sqrt{1+x} = (1+x)^{1/2}$ and $\frac{1}{1+x} = (1+x)^{-1}$ also have exponents. Can they be expanded too?

They can, and the same coefficient pattern still works for *any* rational n . Two things change. The expansion never ends, and it is only valid when x is small enough.

KEY · IB HL For $n \in \mathbb{Q}$:

$$(a + b)^n = a^n \left(1 + n\frac{b}{a} + \frac{n(n-1)}{2!} \left(\frac{b}{a}\right)^2 + \dots \right)$$

In practice you first factor the expression into the form $(1 + \text{something})^n$, which turns the booklet formula into the version you actually compute with:

$$(1 + x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots, \quad |x| < 1.$$

Why does the expansion stop for whole numbers but not otherwise? Look at the coefficients $n(n-1)(n-2)\dots$. If $n = 3$, the third factor is $n - 3 = 0$, so every later coefficient is zero and the series ends: that is the finite theorem of §2.9. If $n = \frac{1}{2}$ or $n = -1$, no factor is ever zero, so the terms continue forever.

The condition $|x| < 1$ should look familiar. Put $n = -1$:

$$(1 + x)^{-1} = 1 - x + x^2 - x^3 + \dots$$

This is a geometric series with first term 1 and common ratio $-x$, and it converges exactly when $|-x| < 1$. The two infinite series of this chapter are the same idea seen from two directions.

EXAMPLE 2.28

Expand $(1 - 3x)^{1/3}$ up to and including the term in x^2 , and state the values of x for which the expansion is valid.

Apply the series with $n = \frac{1}{3}$ and x replaced by $-3x$:

$$(1 - 3x)^{1/3} = 1 + \frac{1}{3}(-3x) + \frac{\frac{1}{3}(-\frac{2}{3})}{2}(-3x)^2 + \dots = 1 - x - x^2 - \dots$$

Validity requires $|-3x| < 1$, that is $|x| < \frac{1}{3}$.

When the bracket does not already start with 1, factor out the larger term first, as the booklet formula does. The expansion is then valid for $|\frac{b}{a}| < 1$.

EXAMPLE 2.29

Expand $\frac{1}{2+x}$ in ascending powers of x up to x^2 , stating the validity.

Factor out the 2:

$$(2+x)^{-1} = \frac{1}{2} \left(1 + \frac{x}{2}\right)^{-1} = \frac{1}{2} \left(1 - \frac{x}{2} + \frac{x^2}{4} - \dots\right) = \frac{1}{2} - \frac{x}{4} + \frac{x^2}{8} - \dots$$

valid for $|\frac{x}{2}| < 1$, that is $|x| < 2$.

EXAMPLE 2.30

By expanding $(1 - x)^{1/2}$ up to x^2 and choosing a suitable value of x , find an approximation to $\sqrt{2}$.

The expansion is

$$(1 - x)^{1/2} = 1 - \frac{x}{2} - \frac{x^2}{8} - \dots$$

Now pick x so that $1 - x$ connects to 2. Take $x = 0.02$: then $1 - x = 0.98 = \frac{49}{50}$, so

$$\sqrt{0.98} = \frac{7}{\sqrt{50}} = \frac{7\sqrt{2}}{10} \implies \sqrt{2} = \frac{10}{7}\sqrt{0.98}.$$

The series gives $\sqrt{0.98} \approx 1 - 0.01 - 0.00005 = 0.98995$, hence

$$\sqrt{2} \approx \frac{10}{7}(0.98995) \approx 1.41421,$$

agreeing with the true value to five decimal places, from just three terms. The key is small x : the closer x is to 0, the faster the series converges.

CAUTION Always state the interval of validity alongside an HL binomial expansion; the series is meaningless outside it. And the expansion needs the form $(1 + \bullet)^n$: factor first, then expand.

At HL these expansions return as Maclaurin series in Ch. 17, where the same coefficients arise from repeated differentiation.

YOUR TURN 2.10 The Binomial Series HL

47. Expand each up to and including the term in x^3 , stating the validity.

(a) $(1 + x)^{-1}$ [2] (b) $(1 - x)^{-2}$ [3]

48. Expand each up to and including the term in x^2 , stating the validity.

(a) $(1 + 4x)^{1/2}$ [3] (b) $(1 + x)^{1/3}$ [3]

49. Expand $\frac{1}{3 - x}$ in ascending powers of x up to x^2 , stating the validity. [3]

50. Expand $(8 + x)^{1/3}$ in ascending powers of x up to x^2 , stating the validity. [3]

51. Use the expansion of $(1 + x)^{1/2}$ up to the term in x^2 , with $x = 0.1$, to estimate $\sqrt{1.1}$ to four decimal places. [3]

Reflections

TOK Fold a sheet of paper 42 times and the stack reaches the Moon. The arithmetic is exact, but no one has ever folded a sheet more than about eight times, so the object the calculation describes has never existed. Can a statement be mathematically true and physically impossible at the same time? What is the calculation actually telling us about?

TOK The sum $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots = 1$ is an equality, yet no finite number of steps ever reaches 1. We accept the answer because we define the sum to be the limit of the partial sums. Is that a discovery about infinity, or a decision we made so that the addition has an answer?

RW Around 450 BCE Zeno argued that a runner can never finish a race. First he must reach the halfway point, then half of what remains, and so on without end. Each step of the argument looks correct, and it troubled mathematicians for two thousand years. The answer is one line of this chapter: $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots = 1$. Infinitely many steps can have a finite total, provided the steps shrink fast enough.

IM The triangle of binomial coefficients was known long before Blaise Pascal wrote about it in 1654. Around the 2nd century BCE the Indian scholar Pingala counted the possible patterns of short and long syllables in poetry, a problem that produces exactly these numbers; a later Indian commentary set them out as a triangle and called it *Meru Prastara*, the staircase of Mount Meru. The Persian mathematicians al-Karaji and Omar Khayyam used it around 1000–1100 CE, and it is still called the Khayyam triangle in Iran. The Chinese mathematician Yang Hui published it in 1261, and it is called Yang Hui's triangle in China. Tartaglia used it in Italy in the 1500s, where it is called Tartaglia's triangle. The same pattern was therefore found independently in several places over eighteen centuries, and each culture named it after a different person. What does that tell you about who a mathematical result belongs to?

EXT A geometric series with $|r| < 1$ is the simplest example of a **power series**, a sum of increasing powers of x :

$$\sum_{k=0}^{\infty} x^k = \frac{1}{1-x}, \quad |x| < 1.$$

This one identity produces many others. Differentiate each term, or integrate each term, and you obtain the series for $\ln(1+x)$, for $\arctan x$, and for many further functions. In calculus, the topic of this chapter is *the seed from which the analysis of functions grows*.

End-of-Chapter Problems — Chapter 2: Sequences & Series

1. An arithmetic sequence has first term a and common difference d . The 5th term is 17 and the sum of the first 10 terms is 155.
 - (a) Write down two equations in a and d . [2]
 - (b) Find a and d . [2]
 - (c) Find the first negative term of the sequence. [2]

2. An arithmetic sequence has $u_3 = 7$ and $u_9 = 25$.
 - (a) Find u_1 and d . [3]
 - (b) Find S_{20} . [2]
 - (c) Find the least value of n such that $S_n > 600$. [3]

3. An arithmetic series has $u_1 = 2$ and $S_{10} = 155$.
 - (a) Find the common difference. [2]
 - (b) Find the sum $\sum_{k=11}^{20} u_k$. [4]

4. An arithmetic sequence has 20 terms. The sum of the odd-positioned terms $u_1 + u_3 + u_5 + \cdots + u_{19}$ is 160, and the sum of the even-positioned terms $u_2 + u_4 + u_6 + \cdots + u_{20}$ is 140. Find the first term u_1 and the common difference d . [6]

5. The sum of the first n terms of a series is given by $S_n = n^2 + 3n$.
 - (a) Find u_1 and u_2 . [2]
 - (b) Show that the series is arithmetic and state the common difference. [3]
 - (c) Find the value of n such that $u_n = 43$. [2]

6. An AP has $u_1 = 120$ and $d = -3$.
 - (a) Find the first term that is negative. [3]
 - (b) Find the value of n that gives the maximum partial sum S_n , and find that maximum. [4]

7. A geometric sequence has first term a and common ratio r , where $r > 0$. Given $u_3 + u_4 = 12$ and $u_5 + u_6 = 3$, find a and r . [5]

8. The first three terms of a geometric sequence are k , 6, and $k + 5$.

- (a) Show that $k^2 + 5k - 36 = 0$, and hence find the possible values of k . [4]
 (b) For the value of k that gives a convergent series, find S_∞ . [3]

9.

- (a) Show that $\sum_{k=1}^n r^{k-1} = \frac{1-r^n}{1-r}$ for $r \neq 1$ by multiplying S by r and subtracting. [3]
 (b) Hence find $\sum_{k=1}^{10} \left(\frac{2}{3}\right)^{k-1}$, giving your answer as an exact fraction. [3]

10. A geometric series has $u_1 = 3$ and $S_\infty = 12$.

- (a) Find the common ratio. [2]
 (b) Show that $S_n = 12\left(1 - \left(\frac{3}{4}\right)^n\right)$. [2]
 (c) Find the least value of n such that $S_n > 11.9$. [3]

11.

- (a) Find $\sum_{k=1}^n (4k - 3)$ in terms of n . [3]
 (b) Find the value of n such that $\sum_{k=1}^n (4k - 3) = 780$. [3]

12. Consider the geometric series $1 + 2x + 4x^2 + 8x^3 + \dots$

- (a) Write down the common ratio. [1]
 (b) Find the values of x for which the sum to infinity exists. [2]
 (c) For these values, find S_∞ as a function of x . [2]

13. A geometric sequence has all positive terms. The sum of the first two terms is 10 and the sum to infinity is 25. Find u_1 and r . [5]**14.** The sum to infinity of a geometric series is $\frac{3}{2}$ times the sum of the first 4 terms. Find the common ratio. [5]**15.** The repeating decimal $0.\overline{142857} = 0.142857142857\dots$ can be written as an infinite geometric series.

- (a) Write down u_1 and the common ratio r . [2]
 (b) Find the exact fractional value of $0.\overline{142857}$. [3]
 (c) Verify your answer by converting the fraction from part (b) back to a decimal using long division. [1]

16. GDC A machine is bought for \$60 000 and depreciates at 18% per year.

- (a) Find its value after 4 years. [2]
 (b) Find the least number of complete years until the value falls below one quarter of the purchase price. [3]

17. GDC \$8000 is invested for 6 years at 5% per year, compounded quarterly. Over the same period, inflation runs at 3% per year.

- (a) Find the future value of the investment. [2]
 (b) Find the real value of the investment after 6 years. [2]
 (c) Comment on what your answers say about the investment's true growth. [1]

18. GDC An investment account pays 5% per year, compounded annually. A sum P is invested.

- (a) Write an expression for the value of the account after n years. [1]
 (b) Find the value of P if the account is worth \$20 000 after 10 years. [3]
 (c) Find the number of complete years for the account to triple in value. [3]

19. GDC The value of a car decreases by 18% each year. A new car costs \$32 000.

- (a) Find the value of the car after 3 years. [2]
 (b) Find the year in which the value first falls below \$10 000. [3]
 (c) Find the total loss in value over the first 5 years. [3]

20. A school committee of 6 is to be chosen from 9 teachers and 7 students.

- (a) Find the number of committees containing exactly 4 students. [2]
 (b) Find the number of committees containing at least 4 students. [3]

21.

- (a) Show that $\binom{n}{2} = \frac{n(n-1)}{2}$, and explain the connection with the sum $1+2+\cdots+(n-1)$. [3]
 (b) Hence find n such that $\binom{n}{2} = 190$. [2]

22. In the expansion of $(1+kx)^8$, the coefficient of x^2 is 112.

- (a) Find the possible values of k . [3]
 (b) For $k > 0$, find the coefficient of x^3 . [2]

23. Consider the expansion of $\left(x + \frac{2}{x^2}\right)^{12}$.

- (a) Find the term independent of x . [4]

- (b) Find the coefficient of x^3 . [2]

24.

- (a) Expand $(1 + 2x)^{-2}$ in ascending powers of x up to and including the term in x^3 , and state the values of x for which the expansion is valid. [4]
(b) Use your expansion with $x = 0.01$ to estimate $\frac{1}{1.02^2}$ to 5 decimal places. [2]

25.

- (a) Show that $(4 + x)^{1/2} = 2 \left(1 + \frac{x}{4}\right)^{1/2}$, and expand it in ascending powers of x up to the term in x^2 , stating the validity. [4]
(b) Hence find an approximation to $\sqrt{4.2}$, giving five decimal places. [2]

26. A ball is dropped from a height of 10 m. After each bounce it reaches 70% of the height it fell from.

- (a) Find the height reached after the 3rd bounce. [2]
(b) Find the total vertical distance travelled by the ball before it comes to rest. [4]

27. The first, second, and fifth terms of an arithmetic sequence with first term 3 are, in that order, the first, second, and third terms of a geometric sequence.

- (a) Given the arithmetic sequence is non-constant, show that its common difference is $d = 6$ and the geometric sequence has common ratio $r = 3$. [4]
(b) Determine whether the geometric series $3 + 9 + 27 + \dots$ has a sum to infinity, giving a reason. [2]

28. A sequence is defined by $u_1 = 3$ and $u_{n+1} = 2u_n + 1$ for $n \geq 1$.

- (a) Write down u_2 , u_3 , and u_4 . [2]
(b) Let $v_n = u_n + 1$. Show that v_n is a geometric sequence. [3]
(c) Find an explicit formula for u_n . [2]

29. Consider the two arithmetic sequences 4, 7, 10, ... and 3, 7, 11, ... Both sequences contain the value 7.

- (a) Find the general term of each sequence. [2]
(b) Find the next three values that appear in both sequences. [4]
(c) Show that the common terms form an arithmetic sequence and state its common difference. [2]

30. A student can invest \$3000 in one of two accounts. Account A pays 5% per year *sim-*

ple interest; account B pays 4% per year, compounded annually.

- (a) Write down an expression for the value of each account after n years. [3]
- (b) Find the value of each account after 10 years. [2]
- (c) Using a table or graph on your GDC, find the least whole number of years after which account B is worth more than account A. [3]

31. GDC A savings plan pays \$500 into an account at the start of each year. The account earns 4% per year, compounded annually.

- (a) Show that after 3 years the total value is $500(1.04)^3 + 500(1.04)^2 + 500(1.04)$. [2]
- (b) Write this as a geometric series and find the total after 10 years. [4]
- (c) Find the number of complete years until the total exceeds \$8 000. [3]

32. GDC A sum of L is borrowed. Interest is added at a rate i per month, and a fixed repayment M is made at the end of each month, so the debt after one month is $L(1+i) - M$.

- (a) Explain why the debt after one month is $L(1+i) - M$. [2]
- (b) Show that the debt after two months is $L(1+i)^2 - M[(1+i) + 1]$. [3]
- (c) Hence show that the debt after n months is $L(1+i)^n - M\left(\frac{(1+i)^n - 1}{i}\right)$. [4]
- (d) A loan of \$20 000 is taken at 6% per year compounded monthly, and is to be cleared in exactly 5 years. Find the monthly repayment. [4]
- (e) Find the total amount repaid, and hence the total interest paid. [3]

33. Consider the expansion of $\left(2x - \frac{3}{x}\right)^{10}$.

- (a) Write down an expression for the general term of the expansion. [2]
- (b) Find the term that is independent of x . [4]
- (c) Find the coefficient of x^4 . [3]
- (d) Explain why the expansion contains no term in x^3 . [2]
- (e) By substituting a suitable value of x , find the sum of all the coefficients. [2]
- (f) State how many terms the expansion has, and which powers of x occur. [2]

34. An arithmetic sequence has first term a and common difference d , where $a > 0$ and $d \neq 0$. Its 1st, 4th and 13th terms form a geometric sequence.

- (a) Show that $3d = 2a$. [4]
- (b) Given that $a = 3$, find d and write down the first four terms of the arithmetic sequence. [3]
- (c) Find the common ratio of the geometric sequence. [2]
- (d) Find the sum of the first 20 terms of the arithmetic sequence. [3]
- (e) Find the sum of the first 8 terms of the geometric sequence. [3]

Challenge Questions

35. Turning products into sums. This investigation shows why a logarithm converts a geometric sequence into an arithmetic one.

- Let $u_n = \log_2(3^n)$. Show that $\{u_n\}$ is arithmetic and state its common difference. [3]
- Find the exact value of $\sum_{k=1}^{20} u_k$. [3]
- Find the smallest value of n such that $u_n > 30$. [3]
- Now let $v_n = \log_b(a^n)$, where $a > 0$ and $b > 1$ are fixed. Show that $\{v_n\}$ is arithmetic for every such a and b , and state its common difference. [3]
- Let $\{w_n\}$ be any geometric sequence with all terms positive and common ratio r . Show that $\{\log_b w_n\}$ is arithmetic, and state its common difference. [4]
- Part (e) is the reason logarithmic scales are used for quantities that grow by multiplication. Explain the connection in your own words, and give one example of such a quantity. [3]

36. Counting multiples. Each set of multiples below is an arithmetic sequence, so its total is an arithmetic series. Throughout, work with the integers from 1 to 1000.

- Find the sum of those divisible by 3, and the sum of those divisible by 5. [3]
- Explain why adding your two answers counts some integers twice, and find the sum of those divisible by both. [3]
- Hence find the sum of those divisible by 3 or by 5. [2]
- Find the sum of those divisible by 3 or by 5, but not by both. [3]
- Let p and q be distinct primes and let N be a positive integer. Show that the sum of the integers from 1 to N divisible by p or q but not both is

$$S_p + S_q - 2S_{pq}, \quad \text{where } S_m = m \frac{t(t+1)}{2}, \quad t = \left\lfloor \frac{N}{m} \right\rfloor.$$

- Use the result of part (e) with $p = 7$, $q = 11$ and $N = 1000$. Verify your answer by a second method. [3]

37. An infinite geometric series has $S_\infty = 4$ and $u_2 = \frac{3}{4}$.

- Find two possible pairs of values for a (first term) and r . [4]
- For each solution, a second series is formed by squaring every term. Show each squared series is geometric and find its sum to infinity. [4]
- For the solution with $r < \frac{1}{2}$, find the sum to infinity of the series formed by taking the positive square root of every term. [3]

