

GEOMETRY & TRIGONOMETRY

CHAPTER 10

Vectors

10

SYLLABUS

3.12–3.18

HL ONLY

A stool with three legs never wobbles. A stool with four legs, on the same uneven floor, almost always does.

This is not bad carpentry. It is geometry. Three points always lie in one flat surface, whatever their heights, so three feet always find the floor together. A fourth point does not have to lie in that surface, and when it does not, the stool rocks between two pairs of legs. Photographers and surveyors know this, which is why a tripod has three legs and not four.

That fact cannot be written down in ordinary arithmetic. It is not about how big anything is; it is about position, direction, and flatness. To state it, and to prove it, we need a language in which a direction is a thing you can calculate with. That language is the subject of this chapter.

A **vector** is a quantity with both a size and a direction. Speed is a number, but velocity is a vector. Distance is a number, but displacement is a vector. A pilot flying due north through a wind from the west travels neither due north nor at their airspeed, because their motion and the wind point different ways and both must be accounted for.

Once a quantity has a direction, ordinary addition stops working. Adding three and three no longer has to give six. If the two point the same way you get six, if they point opposite ways you get zero, and any value in between is possible. The answer depends on the angle.

This chapter builds that new arithmetic, then uses it on geometry. A line in space becomes a single equation. The angle between two directions comes straight out of a product. A flat plane at any tilt is fixed by one perpendicular arrow, and by the last section we can say exactly what three points and a plane have to do with each other.

The result is a language for describing space. Physics is written in it, computer graphics runs on it, and navigation depends on it, from the pilot's crosswind to the satellites overhead.

By the end of this chapter you will be able to add and scale vectors, use the scalar product to find angles and test for perpendicularity, and use the vector product to find areas and normals. You will write the equation of a line and of a plane, decide how two lines or three planes are arranged, and find shortest distances. The course builds all this because a vector

carries a size and a direction in one object. That makes it the natural tool for measuring position and movement in three dimensions, where coordinates alone become unmanageable.

10.1 Vectors: Magnitude and Direction

A vector has exactly two properties: a **magnitude**, which is its length, and a **direction**. Where it sits in space does not matter. Two arrows of the same length pointing the same way are the same vector.

An arrow drawn to represent a vector is called a **directed line segment**. “Directed” because it has an arrowhead, and “line segment” because it is a piece of a line with two ends. Examination questions use this phrase, so it is worth knowing that it means an arrow and nothing more.

Vectors must be kept apart from ordinary numbers, which in this context are called **scalars**. Three notations are in common use, and they all mean the same thing. Printed books, including this one, use a bold letter **a**. By hand you cannot write bold type, so you write either a letter with an arrow above it, $\vec{a}$, or an underlined letter, $\underline{a}$. An arrow is also used for a vector named by its two endpoints: $\overrightarrow{AB}$ means the vector from the point A to the point B .

The most useful representation is the **column vector**. It lists how far to travel along each axis to get from the tail to the head.

$$\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} = a_1 \hat{\mathbf{i}} + a_2 \hat{\mathbf{j}} + a_3 \hat{\mathbf{k}},$$

Here $\hat{\mathbf{i}}$, $\hat{\mathbf{j}}$ and $\hat{\mathbf{k}}$ are the **base vectors**: arrows of length 1 along the x -, y - and z -axes. The hat is there to record that each one has length 1, a point returned to in the next subsection. The formula booklet prints them without hats, as $\mathbf{i}$, $\mathbf{j}$, $\mathbf{k}$; the two mean the same thing. The column form and the base-vector form also mean the same thing, so use whichever is convenient.

The algebra of arrows

Vectors are added **tip to tail**. Draw the second vector starting where the first one ends. The sum is the arrow from the start of the first to the end of the second. In components it is simple: everything happens coordinate by coordinate.

KEY $\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} a_1 + b_1 \\ a_2 + b_2 \\ a_3 + b_3 \end{pmatrix}, \quad k \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} = \begin{pmatrix} ka_1 \\ ka_2 \\ ka_3 \end{pmatrix}.$




Figure 10.1 Adding tip to tail. Draw $\mathbf{b}$ starting where $\mathbf{a}$ ends; the sum runs from the start of $\mathbf{a}$ to the end of $\mathbf{b}$.

Scalar multiplication by k scales the length by $|k|$ and reverses the direction when k is negative. This gives a clean test: two nonzero vectors are **parallel** exactly when one is a scalar multiple of the other, $\mathbf{a} = k\mathbf{b}$.

Two particular vectors are worth naming.

The **negative** of $\mathbf{v}$, written $-\mathbf{v}$, is the vector of the same length pointing the opposite way. It is the case $k = -1$ of scalar multiplication. Subtraction is then just addition of the negative:

$$\mathbf{a} - \mathbf{b} = \mathbf{a} + (-\mathbf{b}).$$

The **zero vector** $\mathbf{0}$ has magnitude 0 and no direction at all. It is what you get from $\mathbf{v} + (-\mathbf{v})$, or from $0\mathbf{v}$. It is the only vector whose direction is undefined, and it is written in bold to keep it apart from the number 0.

Magnitude and unit vectors

Three words are used for the same quantity. The **magnitude** of a vector, its **length**, and its **modulus** all mean the size of the arrow, and all are written $|\mathbf{v}|$. This book uses whichever reads more naturally in the sentence. The value comes straight from Pythagoras, extended into three dimensions.

KEY · IB The magnitude of $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$ is $|\mathbf{v}| = \sqrt{v_1^2 + v_2^2 + v_3^2}$.

Dividing a vector by its own length leaves the direction untouched but sets the length to 1. The result is a **unit vector**, written $\hat{\mathbf{v}}$, the pure direction stripped of any magnitude, ready to be scaled to any length you like.

KEY $\hat{\mathbf{v}} = \frac{\mathbf{v}}{|\mathbf{v}|}$ is the unit vector in the direction of $\mathbf{v}$.

The base vectors are unit vectors, which is what their hats record: $|\hat{\mathbf{i}}| = |\hat{\mathbf{j}}| = |\hat{\mathbf{k}}| = 1$, one unit along each axis. So writing $\mathbf{a} = a_1\hat{\mathbf{i}} + a_2\hat{\mathbf{j}} + a_3\hat{\mathbf{k}}$ says that $\mathbf{a}$ is built by taking a_1 steps of length 1 along x , then a_2 along y , then a_3 along z .

EXAMPLE 10.1

Find a vector of magnitude 6 in the direction of $\mathbf{v} = \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$.

First the length: $|\mathbf{v}| = \sqrt{2^2 + (-1)^2 + 2^2} = \sqrt{9} = 3$. The unit vector is $\hat{\mathbf{v}} = \frac{1}{3}\mathbf{v}$, and scaling it to length 6 means multiplying by 6:

$$6\hat{\mathbf{v}} = \frac{6}{3} \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 4 \\ -2 \\ 4 \end{pmatrix}.$$

Reduce to a unit vector first, then rescale. Those two steps answer every question of the form “find a vector of magnitude m in the direction of”.

YOUR TURN 10.1 Vectors: Magnitude and Direction

1. Find the magnitude of each vector.

- | | | | |
|---|-----|--|-----|
| (a) $\begin{pmatrix} 3 \\ 4 \\ 0 \end{pmatrix}$ | [1] | (b) $\begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$ | [1] |
| (c) $\begin{pmatrix} 0 \\ 3 \\ 4 \end{pmatrix}$ | [1] | (d) $\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ | [2] |

2. Given $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 0 \\ -1 \\ 4 \end{pmatrix}$, evaluate each of the following.

- | | | | |
|--------------------------------|-----|--------------------------------|-----|
| (a) $\mathbf{a} + \mathbf{b}$ | [1] | (b) $\mathbf{a} - \mathbf{b}$ | [1] |
| (c) $\mathbf{a} + 2\mathbf{b}$ | [2] | (d) $3\mathbf{a} - \mathbf{b}$ | [2] |

3. Unit vectors and scaling.

- (a) Find the unit vector in the direction of $\begin{pmatrix} 0 \\ 3 \\ 4 \end{pmatrix}$. [2]
 (b) Find the unit vector in the direction of $\begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$. [2]
 (c) Find a vector of magnitude 6 in the direction of $\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$. [3]
 (d) Explain why every unit vector has magnitude exactly 1, whatever its direction. [2]

4. Decide whether each pair of vectors is parallel, and justify your answer.

- (a) $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\begin{pmatrix} 2 \\ 4 \\ 6 \end{pmatrix}$ [1] (b) $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\begin{pmatrix} 2 \\ 4 \\ 5 \end{pmatrix}$ [2]

10.2 Position Vectors and Geometry

Fix the origin O . The **position vector** of a point A is the arrow $\overrightarrow{OA}$ from the origin to A ; its components are simply the coordinates of A . Position vectors are the bridge between the free-floating algebra of arrows and the fixed points of a geometry problem.

The arrow from one point to another is found by subtraction. Almost everything that follows rests on this one fact.

KEY $\overrightarrow{AB} = \mathbf{b} - \mathbf{a}$, the position vector of the head minus that of the tail.

The distance between A and B is then the magnitude of that displacement, $|\overrightarrow{AB}| = |\mathbf{b} - \mathbf{a}|$, and the midpoint of AB has position vector $\frac{1}{2}(\mathbf{a} + \mathbf{b})$.

EXAMPLE 10.2

Points $A(1, 2, -1)$ and $B(4, 0, 3)$ are given. Find $\overrightarrow{AB}$ and the distance AB .

$$\overrightarrow{AB} = \mathbf{b} - \mathbf{a} = \begin{pmatrix} 4 \\ 0 \\ 3 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} = \begin{pmatrix} 3 \\ -2 \\ 4 \end{pmatrix}, \quad AB = \sqrt{3^2 + (-2)^2 + 4^2} = \sqrt{29}.$$

Head minus tail gives the arrow; the magnitude of that arrow is the distance. Every three-dimensional distance problem is this one line.

Three points A, B, C are **collinear** (on one straight line) when $\overrightarrow{AB}$ and $\overrightarrow{AC}$ are parallel, that is, one is a scalar multiple of the other. This is the vector version of “same direction from a shared point”, and the equation of a line in the next section is built from it.

Proving geometry with vectors

Vectors give a way of proving geometric facts by algebra, with no diagram-chasing and no coordinates. The method is always the same three steps.

KEY Proving a geometric property with vectors.

1. Choose one point as the origin, and name the position vectors of the others with single letters.
2. Write every point in the figure in terms of those letters.
3. Show the required relation by algebra.

Choosing the origin well is what keeps the algebra short. Put it at a vertex that many edges meet.

EXAMPLE 10.3

Prove that the diagonals of a parallelogram bisect each other.

Let the parallelogram be $OABC$, with the origin at O . Write $\overrightarrow{OA} = \mathbf{a}$ and $\overrightarrow{OC} = \mathbf{c}$.

Because $OABC$ is a parallelogram, $\overrightarrow{CB} = \overrightarrow{OA} = \mathbf{a}$, so the fourth vertex is

$$\overrightarrow{OB} = \mathbf{a} + \mathbf{c}.$$

The two diagonals are OB and AC . Find the midpoint of each.

Midpoint of OB : $\frac{1}{2}(\mathbf{0} + \mathbf{a} + \mathbf{c}) = \frac{1}{2}(\mathbf{a} + \mathbf{c})$.

Midpoint of AC : $\frac{1}{2}(\mathbf{a} + \mathbf{c})$.

The two midpoints are the same point. A single point lying halfway along both diagonals is exactly what “bisect each other” means. ■

Notice that no coordinates were used, and the result holds for every parallelogram, in a plane or in space.

EXAMPLE 10.4

M and N are the midpoints of the sides OA and OB of a triangle. Prove that MN is parallel to AB and half its length.

Take the origin at O , with $\overrightarrow{OA} = \mathbf{a}$ and $\overrightarrow{OB} = \mathbf{b}$. The midpoints are then

$$\overrightarrow{OM} = \frac{1}{2}\mathbf{a}, \quad \overrightarrow{ON} = \frac{1}{2}\mathbf{b}.$$

Now find the two displacement vectors, each as head minus tail:

$$\overrightarrow{MN} = \frac{1}{2}\mathbf{b} - \frac{1}{2}\mathbf{a} = \frac{1}{2}(\mathbf{b} - \mathbf{a}), \quad \overrightarrow{AB} = \mathbf{b} - \mathbf{a}.$$

So $\overrightarrow{MN} = \frac{1}{2}\overrightarrow{AB}$.

One vector is a scalar multiple of the other, so the two segments are parallel, and the scalar is $\frac{1}{2}$, so MN is half the length of AB . ■

The whole proof is one line of algebra. This result is known as the midpoint theorem.

YOUR TURN 10.2 Position Vectors and Geometry

5. Points $A(1, 2, -1)$ and $B(4, 0, 3)$ are given.

- | | | | |
|----------------------------------|-----|----------------------------------|-----|
| (a) Find $\overrightarrow{AB}$. | [1] | (b) Find the distance AB . | [2] |
| (c) Find the midpoint of AB . | [2] | (d) Find $\overrightarrow{BA}$. | [1] |

6. Further position-vector work.

- | | |
|---|-----|
| (a) A is $(1, 1, 1)$ and $\overrightarrow{AB} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}$. Find the coordinates of B . | [2] |
| (b) B is the midpoint of AC , with $A(2, 1, 3)$ and $B(4, 5, 3)$. Find C . | [3] |
| (c) Show that $A(1, 1, 1)$, $B(2, 3, 5)$ and $C(4, 7, 13)$ are collinear. | [3] |

7. Proving geometry with vectors. In each part, take the origin at O and write $\overrightarrow{OA} = \mathbf{a}$, $\overrightarrow{OB} = \mathbf{b}$.

- | | |
|--|-----|
| (a) $OACB$ is a parallelogram. Write $\overrightarrow{OC}$ in terms of $\mathbf{a}$ and $\mathbf{b}$. | [2] |
| (b) Find the midpoint of the diagonal OC , and the midpoint of the diagonal AB . | [3] |
| (c) Hence prove that the diagonals of a parallelogram bisect each other. | [2] |
| (d) M and N are the midpoints of OA and OB . Prove that $\overrightarrow{MN} = \frac{1}{2}\overrightarrow{AB}$, and state what this shows about the two segments. | [4] |

10.3 The Scalar Product

There are two ways to multiply vectors, and they answer two different questions. The first, the **scalar product** (or dot product), takes two vectors and returns a number, and that number carries the angle between them. It has two forms, and they always agree.

KEY · IB **Scalar product.** For $\mathbf{a}, \mathbf{b}$ with angle θ between them,

$$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3 \quad \text{and} \quad \mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta.$$

The second form is worth pausing on, because the $\cos \theta$ is not decoration. Ordinary multiplication assumes both quantities are measured along the same line: 3×4 makes sense because both numbers sit on one number line. Two vectors usually point different ways, so multiplying their lengths would measure nothing in particular.

The repair is to bring them onto the same line first. Keep $\mathbf{a}$ as it is, and use only the part of $\mathbf{b}$ that lies along $\mathbf{a}$ — the shadow $\mathbf{b}$ casts on $\mathbf{a}$, which has length $|\mathbf{b}| \cos \theta$. Multiplying those

two gives

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| \times |\mathbf{b}| \cos \theta,$$

the length of $\mathbf{a}$ times how much of $\mathbf{b}$ goes in $\mathbf{a}$'s direction. Doing it the other way round, casting $\mathbf{a}$ onto $\mathbf{b}$, gives $|\mathbf{b}| \times |\mathbf{a}| \cos \theta$, the same number. That is why the order does not matter.




Figure 10.2 The shadow of $\mathbf{b}$ on $\mathbf{a}$. Dropping the tip of $\mathbf{b}$ perpendicularly onto $\mathbf{a}$ leaves a length of $|\mathbf{b}| \cos \theta$, and the scalar product multiplies that by $|\mathbf{a}|$.

The scalar product then obeys the rules you would expect of a multiplication.

- **Commutative:** $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$. The order does not matter.
- **Distributive:** $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$. Brackets expand as usual.
- **Scalars move outside:** $(k\mathbf{a}) \cdot \mathbf{b} = k(\mathbf{a} \cdot \mathbf{b})$. A constant factor can be taken out.
- **Dotted with itself:** $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$, since $\theta = 0$ and $\cos 0 = 1$. The result is the squared length, and this is the line used most often in proofs.

The base vectors show both extremes at once. Each has length 1 and lies along its own axis, so dotting one with itself gives $1 \times 1 \times \cos 0 = 1$. Different axes are at right angles, so dotting two different base vectors gives 0.

$$\hat{\mathbf{i}} \cdot \hat{\mathbf{i}} = \hat{\mathbf{j}} \cdot \hat{\mathbf{j}} = \hat{\mathbf{k}} \cdot \hat{\mathbf{k}} = 1, \quad \hat{\mathbf{i}} \cdot \hat{\mathbf{j}} = \hat{\mathbf{j}} \cdot \hat{\mathbf{k}} = \hat{\mathbf{k}} \cdot \hat{\mathbf{i}} = 0.$$

This is exactly why the component form works. Expanding $(a_1\hat{\mathbf{i}} + a_2\hat{\mathbf{j}} + a_3\hat{\mathbf{k}}) \cdot (b_1\hat{\mathbf{i}} + b_2\hat{\mathbf{j}} + b_3\hat{\mathbf{k}})$ produces nine terms, and every term pairing two *different* base vectors is wiped out by that zero. Only the three matching terms survive, leaving $a_1b_1 + a_2b_2 + a_3b_3$.

The component form is easy to compute; the cosine form carries the geometry. Setting the two equal is what lets you find the angle between any two directions, and it is used throughout the chapter.

KEY · IB

$$\cos \theta = \frac{a_1b_1 + a_2b_2 + a_3b_3}{|\mathbf{a}| |\mathbf{b}|}.$$

EXAMPLE 10.5

Find the angle between $\mathbf{a} = \begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$.

Compute the three ingredients: $\mathbf{a} \cdot \mathbf{b} = 2 + 2 + 4 = 8$, $|\mathbf{a}| = \sqrt{4 + 1 + 4} = 3$, $|\mathbf{b}| = 3$. Then

$$\cos \theta = \frac{8}{9} \implies \theta = \arccos \frac{8}{9} \approx 27.3^\circ.$$

Dot product over product of lengths: the recipe never changes, whether the vectors come from a triangle, a pair of lines, or a pair of planes.

The most important special case is a right angle. Since $\cos 90^\circ = 0$, the scalar product of perpendicular vectors is zero, and the converse holds too. This gives a test for perpendicularity that needs no angles at all, just a multiply-and-add.

KEY $\mathbf{a} \perp \mathbf{b} \iff \mathbf{a} \cdot \mathbf{b} = 0$ (for nonzero $\mathbf{a}, \mathbf{b}$).

EXAMPLE 10.6

Find the angle between $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix}$.

The dot product is $\mathbf{a} \cdot \mathbf{b} = (1)(2) + (2)(0) + (2)(-1) = 0$. Because it vanishes, the vectors are perpendicular: $\theta = 90^\circ$. Notice there was no need to find either magnitude, a zero dot product settles the angle instantly.

YOUR TURN 10.3 The Scalar Product

8. Evaluate each scalar product.

- (a) $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ -1 \\ 2 \end{pmatrix}$ [1] (b) $\begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ [1]
 (c) $\begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ [1] (d) $\begin{pmatrix} 3 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ -3 \end{pmatrix}$ [1]

9. Find the angle between each pair of vectors.

- (a) $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ [2] (b) $\begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ [3]
 (c) $\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$ [3]

10. Perpendicularity.

- (a) Given $|\mathbf{a}| = 3$, $|\mathbf{b}| = 4$ and an angle of 60° between them, find $\mathbf{a} \cdot \mathbf{b}$. [2]
 (b) Find k so that $\begin{pmatrix} 2 \\ k \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix} = 0$. [2]
 (c) Find k so that $\begin{pmatrix} 1 \\ k \\ 2 \end{pmatrix}$ is perpendicular to $\begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix}$. [3]

(d) Explain why $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$ for every vector $\mathbf{a}$. [2]

11. A rhombus has sides $\mathbf{a}$ and $\mathbf{b}$ drawn from one vertex, so $|\mathbf{a}| = |\mathbf{b}|$.

(a) Write the two diagonals in terms of $\mathbf{a}$ and $\mathbf{b}$. [2]

(b) Expand $(\mathbf{a} + \mathbf{b}) \cdot (\mathbf{a} - \mathbf{b})$. [3]

(c) Hence prove that the diagonals of a rhombus are perpendicular. [3]

(d) Verify your result for $\mathbf{a} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$. [3]

10.4 The Vector Equation of a Line

A line is fixed by two pieces of information: one point it passes through, and a direction it runs along. Start at a known point $\mathbf{a}$ and move any distance along a direction $\mathbf{b}$. As the **parameter** λ takes every real value, this reaches every point on the line.

KEY · IB **Vector equation of a line:** $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$, where $\mathbf{a}$ is the position vector of a point on the line and $\mathbf{b}$ is a direction vector.

Writing $\mathbf{r} = (x, y, z)$ and reading off each coordinate gives the **parametric form**; eliminating λ between them gives the **Cartesian form**.

KEY · IB **Parametric:** $x = x_0 + \lambda l$, $y = y_0 + \lambda m$, $z = z_0 + \lambda n$.

Cartesian: $\frac{x - x_0}{l} = \frac{y - y_0}{m} = \frac{z - z_0}{n}$, where $\mathbf{b} = \begin{pmatrix} l \\ m \\ n \end{pmatrix}$.




Figure 10.3 A line built from a point and a direction. The vector $\mathbf{a}$ reaches the known point A ; travelling $\lambda\mathbf{b}$ from there lands on a general point R , whose position vector $\mathbf{r}$ closes the triangle. Negative λ runs back the other way, so the whole line is covered. The direction $\mathbf{b}$ has no fixed home, so the free copy drawn above is the same vector.

The direction vector carries the geometry of how two lines relate. The **angle between two lines** is the angle between their direction vectors, found by the scalar product (take the acute value). Two lines are **parallel** when their directions are scalar multiples. And in three dimensions there is a possibility with no two-dimensional analogue: lines that are neither parallel nor crossing.

Two lines in space fall into exactly one of four cases. Two questions settle which: are the directions parallel, and do the lines share a point?

Case	Directions	Points shared	Description
Coincident	parallel	all of them	the same line written two ways
Parallel	parallel	none	different lines, same direction
Intersecting	not parallel	exactly one	they cross at a single point
Skew	not parallel	none	they pass without meeting, like an overpass above a road

Only the last case is new. Two lines in a plane must either be parallel or cross, but in three dimensions they can miss each other entirely.

To classify a pair of lines, first compare directions. If they are scalar multiples, check whether a point of one lies on the other: yes means coincident, no means parallel. If the directions differ, set the two equations equal and try to solve for the parameters: a consistent solution means they intersect (and gives the point); an inconsistent system means they are skew.

EXAMPLE 10.7

Do the lines $\mathbf{r}_1 = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ and $\mathbf{r}_2 = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ intersect?

The directions are not multiples of each other, so the lines are not parallel. Setting components equal:

$$1 + \lambda = 2 + \mu, \quad \lambda = 3 - \mu, \quad 2 + \lambda = 1 + 2\mu.$$

From the first two, $\lambda - \mu = 1$ and $\lambda + \mu = 3$, so $\lambda = 2$, $\mu = 1$. Check in the third: $2 + 2 = 4$ and $1 + 2(1) = 3$. Since $4 \neq 3$, the system is inconsistent, so the lines are **skew**. Two equations fix the parameters; the third decides whether the lines really meet.

Lines in motion: kinematics

So far the parameter has had no meaning attached to it. Give it one and the line equation describes a moving object. Read λ as *time* t , and $\mathbf{r} = \mathbf{a} + t\mathbf{b}$ says: start at $\mathbf{a}$, and move with constant **velocity** $\mathbf{b}$. The direction of $\mathbf{b}$ is the heading; its magnitude $|\mathbf{b}|$ is the **speed**. This is precisely the pilot from the opening page, with velocity the tip-to-tail sum of airspeed and wind.

EXAMPLE 10.8

An aircraft passes the point $(10, 5)$ (coordinates in km) at $t = 0$ and flies with constant velocity $\begin{pmatrix} 150 \\ 200 \end{pmatrix} \text{ km h}^{-1}$. Find its speed, its position at $t = 2$ hours, and whether its path passes through the airport at $(310, 405)$.

The speed is the magnitude of the velocity: $\sqrt{150^2 + 200^2} = 250 \text{ km h}^{-1}$. The position at time t is

$$\mathbf{r} = \begin{pmatrix} 10 \\ 5 \end{pmatrix} + t \begin{pmatrix} 150 \\ 200 \end{pmatrix}, \quad \text{so at } t = 2: \quad \mathbf{r} = \begin{pmatrix} 310 \\ 405 \end{pmatrix}.$$

That *is* the airport, reached at exactly $t = 2$. To test any point, solve the first coordinate for t , then check the others. One value of t must satisfy every equation at once. This is the same test used for intersecting lines.

YOUR TURN 10.4 The Vector Equation of a Line

12. Writing the equation of a line.

- Write a vector equation of the line through $A(1, 2, 3)$ with direction $\begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$. [1]
- Find a direction vector of the line through $A(1, 0, 2)$ and $B(4, 3, 5)$. [2]
- Write a vector equation of that line. [2]

13. Forms of the equation.

- Write $\mathbf{r} = \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}$ in parametric form. [2]
- Write $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ 4 \end{pmatrix}$ in Cartesian form. [2]
- Does $(3, 3, 2)$ lie on $\mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$? Justify your answer. [3]

14. An aircraft passes $(10, 5)$ at $t = 0$, with velocity $\begin{pmatrix} 150 \\ 200 \end{pmatrix} \text{ km h}^{-1}$.

- Find its speed. [2]
- Write its position vector at time t . [2]
- Find its position after 2 hours. [2]

10.5 The Vector Product

The second way to multiply vectors returns not a number but a *vector*. The **vector product** (or cross product) $\mathbf{a} \times \mathbf{b}$ points perpendicular to *both* of the vectors it is built from, and its length measures the area they span. Where the dot product was about alignment, the cross product is about the plane the two vectors sweep out.

KEY-IB **Vector product.** For $\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$, $\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$,

$$\mathbf{a} \times \mathbf{b} = \begin{pmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{pmatrix}, \quad |\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta \hat{\mathbf{n}},$$

where θ is the angle between $\mathbf{a}$ and $\mathbf{b}$, and $\hat{\mathbf{n}}$ is the unit vector perpendicular to both, with direction given by the right-hand rule. Taking magnitudes of the second form gives $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$, since $|\hat{\mathbf{n}}| = 1$.

The direction is fixed by the **right-hand rule**. Four properties are worth keeping in mind.

- **Perpendicular to both.** The result is at right angles to the plane of $\mathbf{a}$ and $\mathbf{b}$. This is what makes it the tool for building normal vectors in the next section.
- **Not commutative.** It is **anti-commutative**: $\mathbf{b} \times \mathbf{a} = -(\mathbf{a} \times \mathbf{b})$. The order matters, and reversing it reverses the answer.
- **Distributive, and scalars come out.** $\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = \mathbf{u} \times \mathbf{v} + \mathbf{u} \times \mathbf{w}$ and $(k\mathbf{v}) \times \mathbf{w} = k(\mathbf{v} \times \mathbf{w})$.
- **Zero for parallel vectors.** If $\mathbf{a}$ and $\mathbf{b}$ are parallel then $\sin \theta = 0$, so $\mathbf{a} \times \mathbf{b} = \mathbf{0}$. In particular $\mathbf{v} \times \mathbf{v} = \mathbf{0}$, the mirror image of $\mathbf{v} \cdot \mathbf{v} = |\mathbf{v}|^2$.

The base vectors again show the pattern most clearly. Each is perpendicular to the other two, so crossing two different ones gives a unit vector along the third axis, and the right-hand rule fixes which way it points. Crossing one with itself gives $\mathbf{0}$, since the angle is 0 and $\sin 0 = 0$.

$$\hat{\mathbf{i}} \times \hat{\mathbf{j}} = \hat{\mathbf{k}}, \quad \hat{\mathbf{j}} \times \hat{\mathbf{k}} = \hat{\mathbf{i}}, \quad \hat{\mathbf{k}} \times \hat{\mathbf{i}} = \hat{\mathbf{j}}, \quad \hat{\mathbf{i}} \times \hat{\mathbf{i}} = \hat{\mathbf{j}} \times \hat{\mathbf{j}} = \hat{\mathbf{k}} \times \hat{\mathbf{k}} = \mathbf{0}.$$

Read the first three in the cyclic order $\hat{\mathbf{i}} \rightarrow \hat{\mathbf{j}} \rightarrow \hat{\mathbf{k}} \rightarrow \hat{\mathbf{i}}$: going forwards gives a plus, and going backwards gives a minus, so $\hat{\mathbf{j}} \times \hat{\mathbf{i}} = -\hat{\mathbf{k}}$.

One array, two products

Both products come out of the same nine numbers. Multiply every component of $\mathbf{a}$ by every component of $\mathbf{b}$ and lay the results in a square.

	b_1	b_2	b_3
a_1	a_1b_1	a_1b_2	a_1b_3
a_2	a_2b_1	a_2b_2	a_2b_3
a_3	a_3b_1	a_3b_2	a_3b_3

The **diagonal** is the scalar product. Those are the three terms where a base vector meets itself, and $\hat{\mathbf{i}} \cdot \hat{\mathbf{i}} = 1$ keeps them; everything off the diagonal is killed by $\hat{\mathbf{i}} \cdot \hat{\mathbf{j}} = 0$. Adding the shaded entries gives $a_1b_1 + a_2b_2 + a_3b_3$.

The **off-diagonal** entries are the vector product, taken in opposite pairs. Here the diagonal is what dies, since $\hat{\mathbf{i}} \times \hat{\mathbf{i}} = \mathbf{0}$, and each pair of mirror-image entries subtracts to give one component:

$$\underbrace{a_2b_3 - a_3b_2}_{\text{the pair missing 1}}, \quad \underbrace{a_3b_1 - a_1b_3}_{\text{the pair missing 2}}, \quad \underbrace{a_1b_2 - a_2b_1}_{\text{the pair missing 3}}.$$

So the component of $\mathbf{a} \times \mathbf{b}$ in a given direction is built from the two rows and columns that leave that direction out — which is exactly why the answer is perpendicular to both. The two products are the same array read two ways: down the diagonal, or across it.

Area of a parallelogram and a triangle

The magnitude has a clean geometric meaning: it is the area of the parallelogram with $\mathbf{a}$ and $\mathbf{b}$ as sides. To see why, read the two factors of $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$ separately. Take $|\mathbf{a}|$ as the base. Then $|\mathbf{b}| \sin \theta$ is the perpendicular height, because $\mathbf{b}$ is the slanted side and $\sin \theta$ keeps only the part of it at right angles to the base. So the magnitude is base $\times$ height, which is the area.

KEY · IB Area of a parallelogram $= |\mathbf{a} \times \mathbf{b}|$, where $\mathbf{a}$ and $\mathbf{b}$ are two adjacent sides.

A triangle is half of such a parallelogram, so its area is $\frac{1}{2}|\mathbf{a} \times \mathbf{b}|$; remember the half, because the booklet prints only the parallelogram.




Figure 10.4 The parallelogram spanned by $\mathbf{a}$ and $\mathbf{b}$. Its area is $|\mathbf{a} \times \mathbf{b}|$, and a triangle on the same two sides has half that area.

EXAMPLE 10.9

Find the area of the triangle with vertices $A(0, 0, 0)$, $B(2, 1, 0)$, $C(1, 0, 2)$.

Take $\overrightarrow{AB} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$ and $\overrightarrow{AC} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$. Their cross product is

$$\overrightarrow{AB} \times \overrightarrow{AC} = \begin{pmatrix} (1)(2) - (0)(0) \\ (0)(1) - (2)(2) \\ (2)(0) - (1)(1) \end{pmatrix} = \begin{pmatrix} 2 \\ -4 \\ -1 \end{pmatrix}.$$

Its magnitude is $\sqrt{4 + 16 + 1} = \sqrt{21}$, so the triangle's area is $\frac{1}{2}\sqrt{21}$. The cross product does the work here: one calculation and one square root replace any trigonometry.

YOUR TURN 10.5 The Vector Product

15. Evaluate each vector product.

- | | | | |
|--|-----|--|-----|
| (a) $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \times \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ | [1] | (b) $\begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} \times \begin{pmatrix} 0 \\ 3 \\ 0 \end{pmatrix}$ | [1] |
| (c) $\begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} \times \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$ | [2] | (d) $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \times \begin{pmatrix} 0 \\ 5 \\ 6 \end{pmatrix}$ | [3] |

16. Areas and magnitudes.

- | | |
|---|-----|
| (a) Given $ \mathbf{a} = 3$, $ \mathbf{b} = 5$ and an angle of 30° , find $ \mathbf{a} \times \mathbf{b} $. | [2] |
| (b) Find the area of the parallelogram with sides $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix}$. | [2] |
| (c) Find the area of the triangle with vertices $A(0, 0, 0)$, $B(2, 1, 0)$, $C(1, 0, 2)$. | [3] |
| (d) Explain why $\mathbf{v} \times \mathbf{v} = \mathbf{0}$ for every vector $\mathbf{v}$. | [2] |

10.6 Planes

A plane is a flat, endless sheet, and like a line it can be built from a point and some directions. A line needed only one direction, because a line is one-dimensional: one number, the parameter, is enough to say where you are on it. A plane is two-dimensional, so it needs two directions and two numbers. Anchor at $\mathbf{a}$, then travel λ lots of $\mathbf{b}$ and μ lots of $\mathbf{c}$. The pair (λ, μ) acts as a coordinate system inside the sheet, and every point of the plane has exactly one such pair. The two directions must not be parallel: if they were, every combination of them would still point along a single line, and the sheet would collapse to that line.

KEY-IB Vector equation of a plane: $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b} + \mu\mathbf{c}$.

Two parameters make this form awkward to use. It is hard to tell whether a point lies on the plane, or whether two such equations describe the same plane. A better description uses not

the directions the plane contains but the single direction it does *not*: the **normal vector** $\mathbf{n}$, perpendicular to the whole sheet. Every displacement within the plane is perpendicular to $\mathbf{n}$, and the scalar product states exactly that.

KEY · IB Normal (scalar-product) form: $\mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n}$,

Cartesian form: $n_1x + n_2y + n_3z = d$, where $\mathbf{n} = \begin{pmatrix} n_1 \\ n_2 \\ n_3 \end{pmatrix}$ and $d = \mathbf{a} \cdot \mathbf{n}$.

The picture behind this is worth holding on to. Take any point R on the plane. The displacement $\mathbf{r} - \mathbf{a}$ from the known point A to R lies flat in the sheet, so it is perpendicular to $\mathbf{n}$ and its scalar product with $\mathbf{n}$ is zero. Expanding $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$ gives the form above.




Figure 10.5 A plane held by one point and one perpendicular direction. Every displacement inside the sheet, such as $\mathbf{r} - \mathbf{a}$, is at right angles to the normal $\mathbf{n}$, so $(\mathbf{r} - \mathbf{a}) \cdot \mathbf{n} = 0$ for every point R of the plane, and for no other point.

A Cartesian plane equation can now be read directly: the coefficients of x, y, z are the normal vector. To find a plane through three points, use the cross product of two edge vectors to build $\mathbf{n}$, then fix d from any one point.

EXAMPLE 10.10

Find the Cartesian equation of the plane with normal $\mathbf{n} = \begin{pmatrix} 3 \\ -2 \\ 2 \end{pmatrix}$ through $A(4, 8, -1)$.

With $\mathbf{n} = (3, -2, 2)$ the equation is $3x - 2y + 2z = d$. Substitute A to find d :

$$d = 3(4) - 2(8) + 2(-1) = 12 - 16 - 2 = -6, \quad \text{so} \quad 3x - 2y + 2z = -6.$$

The normal supplies the coefficients and a single point supplies the constant. That is the whole method.

When no normal is given, the cross product produces one.

EXAMPLE 10.11

Find the Cartesian equation of the plane through $A(2, 0, 0)$, $B(0, 3, 0)$, $C(0, 0, 6)$.

Two edge vectors span the plane: $\overrightarrow{AB} = \begin{pmatrix} -2 \\ 3 \\ 0 \end{pmatrix}$ and $\overrightarrow{AC} = \begin{pmatrix} -2 \\ 0 \\ 6 \end{pmatrix}$. Their cross product is normal to both, hence to the plane:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{pmatrix} (3)(6) - (0)(0) \\ (0)(-2) - (-2)(6) \\ (-2)(0) - (3)(-2) \end{pmatrix} = \begin{pmatrix} 18 \\ 12 \\ 6 \end{pmatrix} = 6 \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}.$$

Any scalar multiple of a normal is a normal, so use the cleaner $(3, 2, 1)$. Through A : $d = 3(2) = 6$, giving $3x + 2y + z = 6$. Check with B : $2(3) = 6$; and C : $1(6) = 6$. Both lie on it. Scaling the normal down before computing d keeps every later number small.

That example settles the stool from the opening page. Any three points not in a straight line give two edge vectors; their cross product is a normal, and a normal with a point fixes a plane. So three points always have exactly one plane through them, and three feet always reach the floor. A fourth point comes with no such guarantee, and the fourth leg is the one left hanging.

Angles with a plane

Two planes meet at an angle, and that angle is the angle between their normals, computed by the ordinary scalar product. To see why, look along the line where the planes meet, so that each plane is seen edge-on and appears as a line. Turning one arm of an angle through 90° , and then the other arm through 90° , leaves the angle unchanged. The normals are exactly the two arms turned in this way, so they carry the same angle.




Figure 10.6 Two planes seen end-on, looking straight along the line where they meet. Each plane shows as a line. The angle between the planes and the angle between their normals are the same angle.

Taking the absolute value of the dot product guarantees the acute answer.

KEY Angle between two planes (normals $\mathbf{n}_1, \mathbf{n}_2$): $\cos \theta = \frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{|\mathbf{n}_1| |\mathbf{n}_2|}$.

EXAMPLE 10.12

Find the acute angle between the planes $x + 2y - z = 3$ and $2x - y + z = 1$.

Read the normals straight off the coefficients: $\mathbf{n}_1 = (1, 2, -1)$ and $\mathbf{n}_2 = (2, -1, 1)$. Then $\mathbf{n}_1 \cdot \mathbf{n}_2 = 2 - 2 - 1 = -1$, and both magnitudes are $\sqrt{6}$, so

$$\cos \theta = \frac{|-1|}{\sqrt{6} \cdot \sqrt{6}} = \frac{1}{6} \Rightarrow \theta \approx 80.4^\circ.$$

The dot product came out negative, which means the angle between the normals is obtuse. Taking the absolute value gives the acute angle instead, which is what the question asks for.

A line and a plane need one extra step. The angle we want is between the line and its projection *in* the plane: the shadow the line would cast if light fell straight down onto the sheet. But the scalar product gives the angle φ between the line and the *normal*. These two angles add to 90° , so subtract from 90° to get the one we want.




Figure 10.7 The angle between a line and a plane is measured to the line's projection in the plane, marked θ , not to the normal. Since the normal stands at 90° to the projection, the two angles at the point of entry satisfy $\theta + \varphi = 90^\circ$.

KEY Angle between a line (direction $\mathbf{d}$) and a plane (normal $\mathbf{n}$): $\sin \theta = \frac{|\mathbf{d} \cdot \mathbf{n}|}{|\mathbf{d}| |\mathbf{n}|}$.

Using $\sin$ instead of $\cos$ is what builds the “ 90° minus” into the formula. The line's angle to the plane is the complement of its angle to the normal.

Intersections: line and plane, plane and plane

To find where a line meets a plane, put the line's parametric coordinates into the plane's Cartesian equation. That gives one equation in λ . Solve it, then substitute back to get the point.

EXAMPLE 10.13

Find where the line $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$ meets the plane $3x - 2y + 2z = 11$.

Parametrically, $x = 1 + 2\lambda$, $y = \lambda$, $z = 2 - \lambda$. Substituting into the plane,

$$3(1 + 2\lambda) - 2\lambda + 2(2 - \lambda) = 11 \Rightarrow 2\lambda + 7 = 11 \Rightarrow \lambda = 2,$$

so the point is $(5, 2, 0)$. Check: $3(5) - 2(2) + 2(0) = 11$. One substitution, one linear equation, one point. If instead the λ terms had cancelled, the line would be parallel to the plane. There is then no solution if the constants disagree, and the whole line lies in the plane if they agree.

Two non-parallel planes do not meet at a point; they meet along a whole **line**, like the spine of an open book. That line runs within both planes, so its direction is perpendicular to both normals, and the cross product builds it directly.

EXAMPLE 10.14

Find a vector equation of the line of intersection of the planes $x + y + z = 6$ and $2x - y + z = 3$. The direction is perpendicular to both normals:

$$\mathbf{d} = \mathbf{n}_1 \times \mathbf{n}_2 = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \times \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix}.$$

For a point on the line, pick a coordinate and set it to zero. With $z = 0$: $x + y = 6$ and $2x - y = 3$, so $x = 3$, $y = 3$. Hence

$$\mathbf{r} = \begin{pmatrix} 3 \\ 3 \\ 0 \end{pmatrix} + t \begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix}.$$

Check the direction: $(2, 1, -3) \cdot (1, 1, 1) = 0$ and $(2, 1, -3) \cdot (2, -1, 1) = 0$. Perpendicular to both normals, as the geometry demands. Setting $z = 0$ fails only if the line never crosses the plane $z = 0$. In that case set $x = 0$ or $y = 0$ instead.

Three planes

Three planes form a system of three linear equations, the same systems solved by row reduction in Ch. 4. Nothing new has to be calculated here. What is new is that every algebraic outcome now has a shape, and the shape is often easier to remember than the algebra.

There are five arrangements worth knowing. The first two have solutions; the last three do not, and they are told apart by how many of the normals are parallel.






Unique solution	Infinitely many solutions	No solutions (inconsistent system)		
		No normals parallel	Two normals parallel	Three normals parallel
				
Three planes meet at a point	Three planes meet along a line	Three planes form a prism	One plane cutting two parallel planes	Three parallel planes

Figure 10.8 The five arrangements of three planes. Only the first has a single solution; the second has a whole line of them. The last three have none, and the number of parallel normals says which one you are looking at.

One further case is possible but rarely asked: if the three equations are multiples of one another, they all describe the same plane, and every point of that plane is a solution.

Row reduction tells you which case you are in without any picture. A row of zeros with a zero on the right means a dependent system, so the planes share a line. A row of zeros with a non-zero on the right means an inconsistent one, so there is no common point. The pictures only tell you what that answer looks like.

EXAMPLE 10.15

The planes $x + y + z = 6$ and $2x - y + z = 3$ from the previous example are joined by a third, $3x + 2z = c$. Describe the configuration of the three planes for $c = 9$ and for $c \neq 9$.

Adding the first two equations gives $3x + 2z = 9$, so every point on their line of intersection satisfies $3x + 2z = 9$. Indeed, check the line $\mathbf{r} = (3, 3, 0) + t(2, 1, -3)$ in the third plane: $3(3 + 2t) + 2(-3t) = 9 + 6t - 6t = 9$, for every t .

So when $c = 9$ the third plane contains the entire line, and all three planes share it, like an open book with three pages on one spine. When $c \neq 9$ the third plane is parallel to that line but misses it. The three planes then bound a triangular prism: each pair meets in a line, the three lines are parallel, and no point lies on all three. In the language of Ch. 4, the system is dependent when $c = 9$ and inconsistent otherwise; the geometry is what those words look like.

YOUR TURN 10.6 Planes

17. Equations of planes.

- Write the Cartesian equation of the plane with normal $\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ through the origin. [1]
- State a normal vector to the plane $2x - y + 3z = 5$. [1]
- Find the Cartesian equation of the plane with normal $\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ through $(2, 0, 1)$. [2]
- Does $(1, 1, 1)$ lie on the plane $x + y + z = 3$? [1]

18. Planes from points, and angles.

- Find the Cartesian equation of the plane through $A(2, 0, 0)$, $B(0, 3, 0)$ and $C(0, 0, 6)$. [4]
- Find the acute angle between the planes $x + 2y - z = 3$ and $2x - y + z = 1$. [4]

19. Intersections.

- Find where the line $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$ meets the plane $3x - 2y + 2z = 11$. [4]
- Find a direction vector of the line of intersection of $x + y + z = 6$ and $2x - y + z = 3$. [3]
- Find the angle between the line in part (a) and the plane $3x - 2y + 2z = 11$. [4]

10.7 Distances

Two objects that never meet are still some distance apart, and asking for that distance always means the same thing: the *shortest* one. There is a single idea behind every case in this section, and it needs no new formula.

KEY **The shortest link is perpendicular.** Of all the segments joining two objects, the shortest is the one at right angles to what it joins. Perpendicular means a scalar product of zero, so the geometry turns straight into an equation.

That gives a method rather than a formula to memorise. Nothing here is printed in the formula booklet, so working it out is the point.

KEY **Finding a shortest distance.**

1. Write the unknown point on each object in terms of its parameter.
2. Set the scalar product of the joining vector with each direction it must be perpendicular to equal to zero.
3. Solve for the parameter or parameters, which locates the feet of the perpendicular.
4. Find the length of the joining vector.

From a point to a line

A general point on the line is $F = \mathbf{a} + \lambda \mathbf{b}$. As λ varies, F slides along the line and the length PF changes. It is shortest exactly when $\overrightarrow{PF}$ is perpendicular to the line, so $\overrightarrow{PF} \cdot \mathbf{b} = 0$. That is one equation in one unknown.




Figure 10.9 Every dashed segment joins P to the line. The shortest of them meets the line at right angles, at the foot of the perpendicular F .

EXAMPLE 10.16

Find the distance from $P(0, 1, 3)$ to the line $\mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$.

A general point on the line is $F = (1 + \lambda, 2\lambda, -1 + 2\lambda)$, so

$$\overrightarrow{PF} = \begin{pmatrix} 1 + \lambda \\ 2\lambda - 1 \\ 2\lambda - 4 \end{pmatrix}.$$

Now make it perpendicular to the direction:

$$\overrightarrow{PF} \cdot \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix} = (1 + \lambda) + 2(2\lambda - 1) + 2(2\lambda - 4) = 9\lambda - 9 = 0,$$

so $\lambda = 1$ and the foot is $F(2, 2, 1)$. Then $\overrightarrow{PF} = (2, 1, -2)$ and the distance is $\sqrt{4 + 1 + 4} = 3$. Notice what the method gives you for free: not just the distance but the point on the line that is closest to P , which many questions ask for as well.

From a point to a plane

For a plane the perpendicular direction is already known: it is the normal $\mathbf{n}$. So no scalar product needs to be set to zero. Instead, walk from P along $\mathbf{n}$ until you land on the plane. The general point on that walk is $\mathbf{p} + t\mathbf{n}$, and putting it into $\mathbf{r} \cdot \mathbf{n} = d$ gives one equation in t . The distance travelled is then $|t| |\mathbf{n}|$.

EXAMPLE 10.17

Find the distance from $P(1, 1, 1)$ to the plane $2x - y + 2z = 12$.

The normal is $\mathbf{n} = (2, -1, 2)$, so a point on the walk from P is

$$\mathbf{p} + t\mathbf{n} = (1 + 2t, 1 - t, 1 + 2t).$$

This lies on the plane when

$$2(1 + 2t) - (1 - t) + 2(1 + 2t) = 12 \implies 9t + 3 = 12 \implies t = 1.$$

The foot is $(3, 0, 3)$, and the distance is $|t| |\mathbf{n}| = 1 \times 3 = 3$. Check the foot on the plane: $6 - 0 + 6 = 12$.

Between two skew lines

Skew lines never meet, yet there is still a shortest gap between them, and the same idea finds it. Take a general point on each, P_1 on the first and P_2 on the second. The joining vector $\overrightarrow{P_1P_2}$ is shortest when it is perpendicular to *both* directions at once. That is two scalar products set to zero, so two equations in the two unknowns λ and μ .

EXAMPLE 10.18

Find the shortest distance between $l_1 : \mathbf{r} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ and $l_2 : \mathbf{r} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$.

General points are $P_1 = (\lambda, \lambda, 1 + \lambda)$ and $P_2 = (1 + \mu, 1, \mu)$, so

$$\overrightarrow{P_1P_2} = \begin{pmatrix} 1 + \mu - \lambda \\ 1 - \lambda \\ \mu - 1 - \lambda \end{pmatrix}.$$

Perpendicular to both directions gives two equations:

$$\overrightarrow{P_1P_2} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = 1 - 3\lambda + 2\mu = 0, \quad \overrightarrow{P_1P_2} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = 2\mu - 2\lambda = 0.$$

The second gives $\mu = \lambda$; the first then gives $1 - \lambda = 0$, so $\lambda = \mu = 1$. The feet are $P_1(1, 1, 2)$ and $P_2(2, 1, 1)$, and

$$\overrightarrow{P_1P_2} = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \quad \text{distance} = \sqrt{2}.$$

Two ordinary simultaneous equations, and the answer comes with both endpoints of the shortest segment attached.

The remaining cases

Nothing else needs a new method. Every other pairing either meets, and so has distance zero, or is parallel, and then any convenient point reduces it to one of the three cases above.

The two objects	What to do
Two parallel lines	Take any point on one line, then use point to line
Two intersecting lines	They meet, so the distance is 0
Two skew lines	Two scalar products set to zero, as above
A line parallel to a plane	Take any point on the line, then use point to plane
A line meeting a plane	They meet, so the distance is 0
Two parallel planes	Take any point on one plane, then use point to plane
Two non-parallel planes	They meet along a line, so the distance is 0

The first step in any of these questions is therefore not calculation but classification. Decide

whether the objects meet, and if they do not, decide which of the three methods the pair reduces to.

EXAMPLE 10.19

Find the distance between the parallel planes $2x - y + 2z = 12$ and $2x - y + 2z = 3$.

Both have normal $(2, -1, 2)$, so they are parallel and never meet. Take any point on the second plane; setting $y = z = 0$ gives $x = \frac{3}{2}$, so $A(\frac{3}{2}, 0, 0)$ lies on it. Now walk from A along the normal to the first plane:

$$2\left(\frac{3}{2} + 2t\right) - (-t) + 2(2t) = 12 \implies 9t + 3 = 12 \implies t = 1,$$

so the distance is $|t| |\mathbf{n}| = 3$. Any other starting point on the second plane gives the same answer, which is what parallel means.

YOUR TURN 10.7 Distances

20. Distance from a point to a line. Throughout, $l : \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$.

- (a) Find the foot of the perpendicular from $P(4, 0, 2)$ to l , and hence the distance from P to l . [4]
- (b) Find the distance from $Q(1, 4, 4)$ to l . [3]
- (c) The line m has equation $\mathbf{r} = \begin{pmatrix} 4 \\ 0 \\ 2 \end{pmatrix} + t \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$. Explain why m is parallel to l , and write down the distance between them. [2]
- (d) Check your answer to part (c) by repeating the method with the point on m where $t = 1$. [3]

21. Distance from a point to a plane. Throughout, $\Pi : x + 2y + 2z = 14$.

- (a) Find the foot of the perpendicular from $P(1, 1, 1)$ to Π , and hence the distance from P to Π . [4]
- (b) Find the distance from the origin to Π . [3]
- (c) Find the distance between Π and the parallel plane $x + 2y + 2z = 5$. [3]
- (d) Show that the line $\mathbf{r} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix}$ is parallel to Π , and write down the distance between the line and the plane. [3]

22. Distance between two lines. Throughout, $l_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ and $l_2 : \mathbf{r} = \begin{pmatrix} 0 \\ 2 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$.

- (a) Show that l_1 and l_2 are skew. [4]
- (b) Write down general points P_1 on l_1 and P_2 on l_2 , and find the values of λ and μ for which $\overrightarrow{P_1P_2}$ is perpendicular to both direction vectors. [5]

- (c) Hence state the two feet of the common perpendicular and the shortest distance between l_1 and l_2 . [3]
- (d) The lines $n_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $n_2 : \mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$ are not skew. Apply the same method to them and explain what your answer means. [4]

Chapter 10 · Vectors

Reflections

TOK Both operations are called products, yet neither behaves like multiplication. The dot product takes two vectors and gives back a number. The cross product changes sign when you swap the order, and returns zero for two vectors that are not themselves zero.

So is either one really multiplication, or have we borrowed a comfortable word for something new? And why these definitions in particular? Nothing forced $|\mathbf{a}||\mathbf{b}|\cos\theta$ on us. It was chosen because it answered a question people wanted answered. How much of mathematics is discovered, and how much is chosen and then found to be useful?

TOK The same plane can be a drawing, a vector equation, or the single line $2x - y + 3z = 7$. The symbols are easier to calculate with. The picture is easier to believe.

If the two disagree in your head, which one do you trust, and what happens in four dimensions, where the picture is no longer available to you?

RW Every three-dimensional game and animated film runs on this chapter. The dot product decides how brightly a surface is lit, by comparing the direction of the light with the direction the surface faces: a value near 1 points at the light and is bright, a value near 0 faces edge-on and stays dark. The cross product builds the normals that say which way each surface points in the first place. At the same time the physics engine adds force and velocity vectors tip to tail, thousands of times a second, which is why a thrown object curves the way your eye expects.

EXT The cross product only exists because we live in three dimensions.

In two dimensions there is no room for a vector perpendicular to two others. In four dimensions or more there is too much room: no single perpendicular direction is picked out, because a whole family of them works. Three dimensions is the only case where two vectors determine exactly one perpendicular line. That is why $\mathbf{a} \times \mathbf{b}$ can be a vector at all.

End-of-Chapter Problems

1. Given $\mathbf{a} = \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 1 \\ 3 \\ -1 \end{pmatrix}$:
 - (a) find $\mathbf{a} + 2\mathbf{b}$; [2]
 - (b) find $|\mathbf{a}|$; [1]
 - (c) find the unit vector in the direction of $\mathbf{a}$. [2]

2. For $A(1, -1, 2)$ and $B(3, 2, 4)$:
 - (a) find $\overrightarrow{AB}$; [1]
 - (b) find $|\overrightarrow{AB}|$; [2]
 - (c) find the midpoint of AB . [2]

3. The vectors $\mathbf{a} = \begin{pmatrix} 3 \\ -1 \\ 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 1 \\ 2 \\ k \end{pmatrix}$ are perpendicular. Find k . [3]

4. Find the angle between $\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$. [4]

5. Consider the line $l: \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}$.
 - (a) Find the point on l when $\lambda = 2$. [2]
 - (b) Determine whether $P(5, 2, 3)$ lies on l . [2]

6. Find, in degrees, the acute angle between two lines with direction vectors $\begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ and $\begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix}$. [4]

7. Given $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix}$, find $\mathbf{a} \times \mathbf{b}$ and $|\mathbf{a} \times \mathbf{b}|$. [4]

8. Find the area of the triangle with vertices $A(1, 0, 0)$, $B(0, 1, 0)$, $C(0, 0, 1)$. [4]

9. A plane Π has normal $\begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}$ and passes through $A(1, 2, -1)$.
 - (a) Find the Cartesian equation of Π . [3]
 - (b) Determine whether $B(2, 1, 0)$ lies on Π . [1]

10. Find the Cartesian equation of the plane through $A(1, 0, 0)$, $B(0, 1, 0)$ and $C(0, 0, 1)$. [4]

11. Given $\mathbf{a} = \begin{pmatrix} 4 \\ -3 \\ 0 \end{pmatrix}$:
 - (a) find $|\mathbf{a}|$; [1]

- (b) find the unit vector $\hat{\mathbf{a}}$; [2]
(c) find the vector of magnitude 10 in the direction of $\mathbf{a}$. [2]

12. Points $A(2, 1, 3)$ and $B(4, 5, 3)$ are given. Find the point C such that B is the mid-point of AC . [3]

13. The line $l : \mathbf{r} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ meets the plane $z = 4$. Find the point of intersection. [4]

14. Find the angle between the vectors $\begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$. [4]

15. Show that the points $A(1, 2, 3)$, $B(2, 4, 5)$ and $C(4, 8, 9)$ are collinear. [4]

16. Given $|\mathbf{a}| = 5$, $|\mathbf{b}| = 3$ and $\mathbf{a} \cdot \mathbf{b} = 9$, find the angle between $\mathbf{a}$ and $\mathbf{b}$. [4]

17. The lines $l_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$ and $l_2 : \mathbf{r} = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ intersect. Find the point of intersection. [5]

18. The plane Π has equation $2x + y - 2z = 6$.

- (a) State a normal vector to Π . [1]
(b) Find where the line $\mathbf{r} = \lambda \begin{pmatrix} 2 \\ 1 \\ -2 \end{pmatrix}$ meets Π . [4]

19. Given $\mathbf{a} = \begin{pmatrix} 1 \\ -2 \\ 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}$:

- (a) find $\mathbf{a} \cdot \mathbf{b}$; [2]
(b) hence state the angle between $\mathbf{a}$ and $\mathbf{b}$. [2]

20. Find the area of the triangle with vertices $A(0, 0, 0)$, $B(3, 0, 0)$, $C(0, 4, 0)$ using the vector product. [4]

21. Find a vector equation of the line through $A(2, -1, 3)$ and $B(4, 1, 7)$. [4]

22. Find the acute angle between the planes $x + y + z = 1$ and $x - y + z = 3$. [5]

23. Given $\mathbf{a} = \begin{pmatrix} 3 \\ 4 \\ 0 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 0 \\ 0 \\ 5 \end{pmatrix}$, find $\mathbf{a} \times \mathbf{b}$ and $|\mathbf{a} \times \mathbf{b}|$. [4]

24. Find the value of t for which $\begin{pmatrix} 1 \\ t \\ 2 \end{pmatrix}$ and $\begin{pmatrix} 2 \\ -1 \\ t \end{pmatrix}$ are perpendicular. [4]

25. At time $t = 0$ seconds a drone is at the point $(2, -1, 10)$ (coordinates in metres) and moves with constant velocity $\begin{pmatrix} 3 \\ 4 \\ 0 \end{pmatrix} \text{ m s}^{-1}$.

- (a) Write down a vector equation for the drone's position at time t . [2]
 (b) Find the drone's speed. [2]
 (c) Determine whether the drone passes through the point $(17, 19, 10)$, and if so, at what time. [3]

26. Find a vector equation of the line of intersection of the planes $x + y - z = 2$ and $2x - y + z = 1$. [6]

27. Consider the planes $\Pi_1 : x + y + z = 3$, $\Pi_2 : x - y + z = 1$ and $\Pi_3 : x + z = c$, where $c \in \mathbb{R}$.

- (a) Find a vector equation of the line of intersection of Π_1 and Π_2 . [4]
 (b) Show that when $c = 2$ the three planes intersect in this line. [2]
 (c) Describe geometrically the configuration of the three planes when $c \neq 2$. [2]

28. Show that the lines $l_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ and $l_2 : \mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}$ intersect, and find the point of intersection. [6]

29. Find the shortest distance from the point $P(1, 2, 3)$ to the plane $2x - y + 2z = 0$, by finding the foot of the perpendicular from P . [6]

30. Find a unit vector perpendicular to both $\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 2 \\ 0 \\ -1 \end{pmatrix}$. [5]

31. The points $A(1, 0, 0)$, $B(0, 2, 0)$ and $C(0, 0, 3)$ are given.

- (a) Find the Cartesian equation of the plane ABC . [3]
 (b) Find the area of triangle ABC . [3]

32. Find the angle that the line $l : \mathbf{r} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1 \\ 2 \end{pmatrix}$ makes with the plane $x + 2y + 2z = 5$. [6]

33. The points $A(2, 0, 1)$, $B(1, 3, 2)$ and $C(4, 1, 0)$ are given, and O is the origin.

- (a) Find $\overrightarrow{AB}$ and $\overrightarrow{AC}$. [2]
 (b) Find $\overrightarrow{AB} \times \overrightarrow{AC}$. [3]
 (c) Find the exact area of triangle ABC . [2]
 (d) Find the Cartesian equation of the plane through A , B and C . [4]
 (e) Find the exact shortest distance from O to that plane. [3]
 (f) Hence find the volume of the tetrahedron $OABC$. [2]

34. Consider the lines

$$l_1 : \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}, \quad l_2 : \mathbf{r} = \begin{pmatrix} 4 \\ 1 \\ 4 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}.$$

- (a) Show that l_1 and l_2 are not parallel. [2]
 (b) Determine whether the lines intersect, and if so find the point of intersection. [5]
 (c) Find the acute angle between the lines. [3]
 (d) Find the Cartesian equation of the plane containing both lines. [4]
 (e) Explain why such a plane exists only because of your answer to part (b). [2]

35. GDC Two aircraft fly at the same altitude. At time t hours their position vectors, in km, are

$$\mathbf{r}_A = \begin{pmatrix} 0 \\ 0 \\ 2 \end{pmatrix} + t \begin{pmatrix} 300 \\ 400 \\ 0 \end{pmatrix}, \quad \mathbf{r}_B = \begin{pmatrix} 600 \\ 0 \\ 2 \end{pmatrix} + t \begin{pmatrix} 0 \\ 500 \\ 0 \end{pmatrix}.$$

- (a) Find the speed of each aircraft. [3]
 (b) Find $\overrightarrow{AB}$ at time t . [3]
 (c) Show that the distance d between them satisfies $d^2 = 100\,000t^2 - 360\,000t + 360\,000$. [4]
 (d) Find the time at which the aircraft are closest, and the least distance between them. [4]

Challenge Questions

36. The shortest distance from a point to a line. Section 10.7 works this out one case at a time. Here you build a formula that does every case, then a second formula, and show the two agree.

Let l be the line $\mathbf{r} = \mathbf{a} + \lambda\mathbf{d}$, and P a point not on it.

- (a) Explain why the shortest distance from P to l is measured along the perpendicular. [2]
 (b) Let $F = \mathbf{a} + \lambda\mathbf{d}$ be the foot of that perpendicular. Explain why $\overrightarrow{PF} \cdot \mathbf{d} = 0$, and use this to show that

$$\lambda = \frac{(\mathbf{p} - \mathbf{a}) \cdot \mathbf{d}}{|\mathbf{d}|^2}.$$

- (c) For $l: \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$ and $P(4, 3, 1)$, find λ , the foot F , and the distance. [5]
 (d) A second formula is $\text{distance} = \frac{|\overrightarrow{AP} \times \mathbf{d}|}{|\mathbf{d}|}$, where A is any point on l . Check that it agrees with part (c). [4]
 (e) Explain why the second formula works, using $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \theta$. (Hint: what is $|\overrightarrow{AP}| \sin \theta$ in the diagram?) [4]

37. How far apart are two skew lines? Two skew lines never meet, yet there is still a shortest distance between them. This investigation finds it.

Let $l_1: \mathbf{r} = \mathbf{a}_1 + \lambda\mathbf{d}_1$ and $l_2: \mathbf{r} = \mathbf{a}_2 + \mu\mathbf{d}_2$ be skew.

- (a) Show that $l_1: \mathbf{r} = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}$ and $l_2: \mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}$ are skew. [4]
 (b) The shortest join between the two lines is perpendicular to both. Explain why its direction must be parallel to $\mathbf{n} = \mathbf{d}_1 \times \mathbf{d}_2$. [3]

- (c) Hence explain why the shortest distance is the length of the projection of $\mathbf{a}_2 - \mathbf{a}_1$ onto $\mathbf{n}$, that is

$$\text{distance} = \frac{|(\mathbf{a}_2 - \mathbf{a}_1) \cdot \mathbf{n}|}{|\mathbf{n}|}.$$

[4]

- (d) Use the formula on the two lines in part (a), giving an exact answer.

[5]

- (e) What does the formula give when the two lines intersect? Explain why your answer is what it should be.

[3]

38. The triple product and volume. Combining both products gives $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})$, a single number called the *scalar triple product*. It measures volume.

Throughout, take $\mathbf{u} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$, $\mathbf{v} = \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$ and $\mathbf{w} = \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$.

- (a) Find $\mathbf{v} \times \mathbf{w}$, then $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})$.

[4]

- (b) The parallelepiped with edges $\mathbf{u}$, $\mathbf{v}$, $\mathbf{w}$ has base area $|\mathbf{v} \times \mathbf{w}|$. Explain why its volume is

$|\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|$. (*Hint: volume = base area $\times$ perpendicular height*)

[4]

- (c) Evaluate $\mathbf{v} \cdot (\mathbf{w} \times \mathbf{u})$ and $\mathbf{u} \cdot (\mathbf{w} \times \mathbf{v})$. State what each result shows about the triple product.

[4]

- (d) Three vectors are *coplanar* when they all lie in one plane. Explain why this happens exactly when the triple product is zero, and test

$\begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}$, $\begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$, $\begin{pmatrix} 2 \\ 5 \\ 2 \end{pmatrix}$.

[4]

- (e) A tetrahedron on the same three edges has one sixth of the parallelepiped's volume. Find the volume of the tetrahedron with vertices at the origin and at the three points $\mathbf{u}$, $\mathbf{v}$, $\mathbf{w}$ above.

[3]

