



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/31

Paper 3 Pure Mathematics 3

May/June 2026

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.



[4]

[illegible]



- [5]



- 3** The equation of a curve is $y = \frac{2x^2 + 7x}{3x^2 - 1}$.

Find an equation of the normal to the curve at the point where $x = 1$. Give your answer in the form $ax + by + c = 0$, where a , b and c are integers.

[6]

This image shows a full page of blank handwriting practice paper. It features multiple sets of horizontal lines. Each set consists of a solid top line, a dashed middle line, and a solid bottom line, providing a guide for letter height and placement. The paper is white with light gray or blue lines. There are faint, diagonal watermarks across the page that read "PapaCambridge" and "papacambridge.com".



- 4 The square roots of $-71 - (20\sqrt{3})i$ can be expressed in the Cartesian form $x + iy$, where x and y are real and exact.

By first forming a quartic equation, find the square roots of $-71 - (20\sqrt{3})i$. Give your answer in exact Cartesian form. [5]

[5]





5 (a) Show that $(\cos x + \sin x)(\operatorname{cosec} x - \sec x) \equiv 2 \cot 2x$.

[4]

This image shows a full page of blank handwriting practice paper. It features multiple sets of horizontal lines. Each set consists of a solid top line, a dashed midline, and a solid bottom line, providing a guide for letter height and placement. The paper is white with light gray or blue lines. There are faint, diagonal watermarks across the page that read "PapaCambridge" and "papacambridge.com".



7

(b) Hence, solve the equation $(\cos x + \sin x)(\operatorname{cosec} x - \sec x) = 3$ for $0 \leq x \leq \pi$.

[3]



6 A curve has the equation $y = e^{(x^2+4x)} \cos^2 x$, where $\cos x \neq 0$.

(a) Show that the x -coordinates of the stationary points of this curve satisfy the equation $\tan x = x + 2$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) By sketching a suitable pair of graphs, show that the equation $\tan x = x + 2$ has only one root in the interval $-\pi < x < \pi$.

[2]





(c) Show by calculation that this root lies between 1.2 and 1.3.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Use the iterative formula $x_{n+1} = \tan^{-1}(x_n + 2)$ to calculate this root correct to 3 decimal places. Give the result of each iteration to 5 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



7



The shaded region on the Argand diagram shows points representing the complex numbers, z , defined by two inequalities. The shaded region is bounded by the circle and the line parallel to the imaginary axis shown in the diagram and includes its boundaries.

- (a) Find two inequalities in terms of z that define the shaded region. [3]

.....

.....

.....

.....

- (b) Calculate the greatest value of $|z|$ for points in the shaded region. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





8 Find the exact value of $\int_0^1 3x^2 e^{3x} dx$.

[illegible]



- 9 The points A and B have position vectors $\begin{pmatrix} 2 \\ -5 \\ 3 \end{pmatrix}$ and $\begin{pmatrix} -6 \\ 4 \\ 1 \end{pmatrix}$ respectively.

The line t passes through the points A and B .

- (a) Find a vector equation for t .

[2]

.....

.....

.....

.....

.....

.....

.....

.....

The line s has equation $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \\ 4 \end{pmatrix} + \mu \begin{pmatrix} -1 \\ 2 \\ -1 \end{pmatrix}$.

- (b) Determine whether s and t are parallel, intersect or are skew.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





DO NOT WRITE IN THIS MARGIN

[illegible]



- 10** The polynomial $ax^3 - 11x^2 + bx + 4$, where a and b are constants, is denoted by $p(x)$. It is given that $(x - 2)$ is a factor of $p(x)$, and that when $p'(x)$ is divided by $(x - 2)$ the remainder is 24.

(a) Find the value of a and the value of b .

[5]

This image shows a full page of primary-ruled paper. It features multiple sets of horizontal dotted lines, each set consisting of two parallel dotted lines with a dashed midline between them. This pattern repeats down the entire page, providing a guide for letter height and placement. The background is white, and there are no margins or other markings present.



(b) (i) When a and b have these values, factorise $p(x)$ completely.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence, solve the inequality $p(x) \geq 0$.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





- 11 (a)** By differentiating $\frac{\cos 4x}{\sin 4x}$, show that if $y = \cot 4x$, then $\frac{dy}{dx} = -4 \operatorname{cosec}^2 4x$. [3]

[illegible]

- (b)** The variables x and y satisfy the differential equation

$$e^{(3x+2)} \frac{dy}{dx} = \sin^2 4y.$$

It is given that $y = \frac{1}{16}\pi$ when $x = -\frac{2}{3}$.

Solve the differential equation, and find the value of y in the range $-\frac{1}{8}\pi \leq y \leq 0$ when $x = 1$. [6]

[illegible]



DO NOT WRITE IN THIS MARGIN



If you use the following page to complete the answer to any question, the question number must be clearly shown.

[illegible]



DO NOT WRITE IN THIS MARGIN





Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.

