



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/61

Paper 6 Probability & Statistics 2

May/June 2026

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages.



- DO NOT WRITE IN THIS MARGIN

[5]

[illegible]



- (a) State suitable null and alternative hypotheses for the test. [1]

.....

.....

.....

.....

.....

- (b) State what is meant by a Type II error in this context. [1]

[illegible]

- (c) Given that the true percentage of faulty talking dolls is now 12%, use a binomial distribution to find the probability of a Type II error. [3]

[illegible]



3 In the work done by a typesetting department, incorrect characters appear randomly and independently at a constant average rate of 5 incorrect characters per 100 pages.

(a) A book that is typeset in the department contains 500 pages. The number of incorrect characters in the book is denoted by X .

(i) State the distribution of X , including the values of any parameters. [1]

.....

.....

.....

.....

(ii) State an appropriate approximating distribution for X , including the values of any parameters.

Justify your choice of approximating distribution.

[2]

[illegible]

(iii) Use your approximating distribution to find $P(X > 35)$. [3]

[illegible]



- (b) Another book that is typeset in the department contains n pages. The probability that the book contains fewer than 18 incorrect characters is less than 0.01.

Use an approximating distribution to find the largest possible value of n .

[5]





- 4 The score on one spin of a 5-sided spinner is denoted by the random variable X with probability distribution as shown in the table.

x	1	2	3	4	5
$\mathbf{P}(X = x)$	0.1	0.3	0.4	0.1	0.1

- (a) Show that $E(X) = 2.8$. [1]

It is given that $\text{Var}(X) = 1.16$.

The spinner is spun 300 times. The score on each spin is noted and the mean, $\bar{X}$, of the 300 scores is found.

- (b) Given that $P(2.8 - k < \bar{X} < 2.8 + k) = 0.2$, find the value of k . [4]

This image shows a full page of blank handwriting practice paper. It features multiple sets of horizontal lines. Each set consists of a solid top line, a dashed middle line, and a solid bottom line, providing a guide for letter height and placement. The paper is white and contains no other markings or text.



- (c) Explain why it was necessary to use the Central Limit Theorem in your answer to 4(b).

[1]

- (d) For another similar spinner, it is claimed that $E(X) = 2.8$. Fred suspects that this value is incorrect. The spinner is spun 300 times, and the mean of the 300 scores is found to be 2.92.

Given that the variance of the score on one spin of this spinner is also 1.16, test Fred's suspicion at the 5% significance level. [5]

[5]





- [2]

- [4]



- (b) The random variable W has the distribution $B(200, 0.02)$. It is given that, using the binomial distribution, $P(W = 2) = 0.146$, correct to 3 decimal places.

Show that a suitable approximating distribution gives the same value of $P(W = 2)$ correct to 2 decimal places, but not correct to 3 decimal places.

[4]

[illegible]



6 A random variable X has probability density function given by

$$f(x) = \begin{cases} \frac{8}{7}\lambda e^{-\lambda x} & 0 < x < \ln 2, \\ 0 & \text{otherwise,} \end{cases}$$

where λ is a constant.

(a) Show that $\lambda = 3$.

[4]

[illegible]



(b) Use integration by parts to find $E(X)$.

[4]





Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

[illegible]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.

