

Calorimetry

solara

November 19, 2024

Heat transfer: relation between mass, time and energy.

Heating a mass of water of 1 kg from 20°C to 30°C demands 1/2 less energy than would a mass of 2 kg from 20°C to 30°C ; and 1/2 less energy than needed to raise 1 kg from 20°C to 40°C.

We observe a double proportionality between energy, mass, & the difference in temperature ΔT .

$$Q = Cm\Delta T$$

$$Q_{heat} = C [J/kg/^\circ C] * m [kg] * \Delta T [^\circ C]$$

C is the coefficient of heat capacity. It's the quantity of heat needed to raise one kg of matter by 1°C. If the value of C is high, the material will need more energy to be heated. A high C characterizes a high resistance to temperature change. (thermal inertia)

To cool down 1 kg of water by 1°C, you have to subtract C .

$$C_{water} = 4180 J * kg^{-1} * ^\circ C^{-1}$$

$$4180 J = 1 kcal$$

$$C_{oil} \sim 2000 J * kg^{-1} * ^\circ C^{-1}$$

$$C_{metals} \sim \leq 1000 J * kg^{-1} * ^\circ C^{-1}$$

The high thermal inertia of water:

—→ stabilizes earth's climates. Oceanic climates have a shorter amplitude than continental ones. (like Belgium, the Netherlands and the UK)

—→ stabilizes the temperature of the human body. $C_{human} \approx C_{water}$.