

5110

Total Patients

249

Total Stroke Cases

4.87

Stroke Rate %

29.20

Average BMI

106.15

Average Glucose Le...

498

Hypertension Patie...

108


Male Stroke Patients

141

Female Stroke Pati...

STROKE RISK ANALYSIS


Stroke Occurrence by Residence Type



age_group

All

Stroke occurrence by Marital Status



Average Glucose level of Stroke patients




Stroke cases by Smoking Status



age_group


All

Stroke Distribution by Gender




Stroke Distribution By Hypertension

Stroke Status ● No Stroke ● Stroke Occured




Stroke distribution by heart disease

Stroke Status ● No Stroke ● Stroke Occured




Stroke occurrence by Age Group

Stroke Status ● No Stroke ● Stroke Occured



Average BMI by Stroke Status



Stroke Analysis

The project analysed factors associated with stroke occurrence using healthcare data. The findings showed that stroke risk was higher among senior citizens, patients with hypertension, heart disease, obesity, high glucose levels, and individuals with a smoking history.

Although females recorded more total stroke cases, males showed a slightly higher stroke rate percentage. Seniors accounted for the majority of stroke cases, making age one of the strongest risk factors. Patients with hypertension and heart disease were significantly more likely to experience stroke compared to others.

Overall, the analysis suggests that lifestyle habits and existing health conditions play a major role in stroke risk and highlights the importance of early detection and preventive healthcare.

|