

# Inverse Landau damping using VP


PHYS 783 - Hasith Perera

# Introduction

- I wanted to look for inverse Landau damping for a change !
- Just to recall from class the condition for inverse Landau damping is as follows

$$\omega_i = \frac{\pi}{2} \omega_{pe} \frac{4\pi e^2}{k^2 m_e} \left. \frac{dF_{e0}}{dv_z} \right|_{v_z = \omega_r/k}$$

- Need a **positive velocity gradient** in the phase space density around the resonant frequency  $v_{\phi}$




# VP - Summary


- This is a 1D-1V Vlasov- Poisson solver
- The electric potential  $\phi(x)$  is solved using a **Green's function** method.
- Multiple species are supported.\*
- Krook collision operator is available. (I didn't test or use it in this project)
- Phase space densities are evolved using a 3rd order **Adams-Bashforth method**
- All Simulation run times for the simulation shown here are < 2 mins on a single core. (No parallelization is used)

# Standing wave - attempt 1

- Initial tests were done to make sure I can see landau damping using a **standing wave initialization**
- A bump on the tail was initialized at  $\pm 3.5 v_{th}$  close to the resonance
  - I see a slower decay when the bumps are added (dashed line)
  - . But not a growth in the field




A



A cut at  $t = 0$ ,  
 $v_{\phi} = 2.86$ ,  $u = \pm 3.5$

D




Dashed lines are when the bumps are added

# Standing wave(contd)


- Let look for the signature in the integrated phase space density.
- Plot A: landau damping simulation
- Plot B: Blue and Green shows how the bumps on the tail evolves
  - I don't fully understand what's going on there
  - It could be the amplitude is too large and the system is showing non-linear effects
- Plot C: Tried a low amplitude run

There is a lot going on need to simplify the simulation even more




# Single propagating wave - attempt 2

- Initialization uses a the fluid equilibrium for an electrostatic wave
- It looks usable
- **There were some issues** on the potential calculation (could be user error too. I am using this for the first time here ! )




- 
- Initialize a bump (with 2 new species)
  - Plot compares a pure landau damped simulation vs a simulation with a bump implemented via simulating 4 species

Species with the same charge and mass are not fully talking to each other !



# Single propagating wave - attempt 3



- 2 species (e and ions) + a bump on the tail externally added at  $u=3.5 v_{th}$ ,  $t = 0$
- Good news
  - At large time the change in phase space density shows **higher energy particles being moved to low energy**





# Energy transfer

- Top plot: Weak Landau damping
- Bottom: With the added bump on the tail



# Field Particle Correlation

- FPC term plotted here

$$\frac{q_s v^2}{2} \frac{\partial \delta f_s(x, v, t)}{\partial v} E(x, t)$$

- Plot A: FPC for landau damping only
- Plot B: FPC for bump on the tail
- When the bump was added notice the signature close to the resonant velocity switched directions.
- But also at later in time we can find some particles that gained energy, This would explain why the field lost energy

