

Automatic Time Step Selection for Numerical Solution of Neutron Diffusion Problems

Aleksandr Avvakumov ¹
Petr Vabishchevich ^{2,3}

Valery Strizhov ²
Aleksandr Vasilev ³

¹National Research Center "Kurchatov Institute", Moscow, Russia

²Nuclear Safety Institute, Russian Academy of Sciences, Moscow, Russia

³North-Eastern Federal University, Yakutsk, Russia

7th FDM:T&A, June 11-16, 2018 Lozenetz

Introduction

bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla
bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla
bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla
bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla

Problem description

Consider second-order parabolic equation

$$\frac{\partial u}{\partial t} - \sum_{\alpha=1}^m \frac{\partial}{\partial x_{\alpha}} \left(k(\mathbf{x}, t) \frac{\partial u}{\partial x_{\alpha}} \right) + s(\mathbf{x}, t)u = f(\mathbf{x}, t), \quad \mathbf{x} \in \Omega, \quad 0 < t \leq T,$$

where $\underline{k} \leq k(\mathbf{x}) \leq \bar{k}$, $\mathbf{x} \in \Omega$, $\underline{k} > 0$.

Boundary condition

$$u(\mathbf{x}, t) = g(\mathbf{x}, t), \quad \mathbf{x} \in \partial\Omega, \quad 0 < t \leq T.$$

Initial condition

$$u(\mathbf{x}, 0) = u^0(\mathbf{x}), \quad \mathbf{x} \in \Omega.$$

Operator notation

Cauchy problem

$$\frac{du}{dt} + A(t)u = f(t), \quad 0 < t \leq T,$$

with initial condition

$$u(0) = u_0.$$

Assume $A(t) \geq 0$ in H then

$$\|u(t)\| \leq \|u_0\| + \int_0^t \|f(\theta)\| d\theta.$$

Solution evaluation

Introduce irregular time grid

$$t^0 = 0, \quad t^{n+1} = t^n + \tau^{n+1}, \quad n = 0, 1, \dots, N-1, \quad t^N = T.$$

Implicit scheme are used

$$\frac{y^{n+1} - y^n}{\tau^{n+1}} + A^{n+1}y^{n+1} = f^{n+1}, \quad n = 0, 1, \dots, N-1,$$

and initial condition $y^0 = u^0$.

Layerwise estimate ($A^{n+1} \geq 0$)

$$\|y^{n+1}\| \leq \|y^n\| + \tau^{n+1}\|f^{n+1}\|.$$

Difference estimate

$$\|y^{n+1}\| \leq \|u^0\| + \sum_{k=0}^n \tau^{k+1}\|f^{k+1}\|.$$

Solution error

For $z^n = y^n - u^n$:

$$\frac{z^{n+1} - z^n}{\tau^{n+1}} + A^{n+1} z^{n+1} = \psi^{n+1}, \quad n = 0, 1, \dots, N-1,$$

$$z^0 = 0.$$

Approximation error

$$\psi^{n+1} = f^{n+1} - \frac{u^{n+1} - u^n}{\tau^{n+1}} - A^{n+1} u^{n+1}.$$

Layerwise estimate

$$\|z^{n+1}\| \leq \sum_{k=0}^n \tau^{k+1} \|\psi^{k+1}\|.$$

Difference estimate

$$\|z^{n+1}\| \leq \delta t^{n+1}.$$

Time step selection algorithm

- 1 Predictable time step: $\tilde{\tau}^{n+1} = \gamma \tau^n$ (eg $\gamma = 1.25$)
- 2 Predictive solution \tilde{y}^{n+1} : an explicit scheme, $\tilde{t}^{n+1} = t^n + \tilde{\tau}^{n+1}$
- 3 Estimation of approximation error: by found \tilde{y}^{n+1} from an implicit scheme
- 4 Step selection τ^{n+1} : $\|\psi^{n+1}\| \approx \delta$
- 5 Solution on a new time layer y^{n+1} : an implicit scheme, $t^{n+1} = t^n + \tau^{n+1}$

Neutron diffusion equation

One group diffusion approximation with one group delayed neutron sources

$$\frac{1}{\nu} \frac{\partial \phi}{\partial t} - \nabla \cdot D \nabla \phi + \Sigma_a \phi = (1 - \beta) \nu \Sigma_f \phi + \lambda c,$$

$$\frac{\partial c}{\partial t} + \lambda c = \beta \nu \Sigma_f \phi.$$

Boundary condition

$$D \frac{\partial \phi}{\partial n} + \gamma \phi = 0.$$

Initial conditions

$$\phi(0) = \phi^0, c(0) = c^0.$$

Calculated formulas

The approximation error

$$\begin{aligned}\tilde{\psi}^{n+1} &= (A^{n+1} - A^n)\varphi^n + A^{n+1}(\tilde{\varphi}^{n+1} - \varphi^n) \\ &= \tilde{\tau}^{n+1} \left(\frac{A^{n+1} - A^n}{\tilde{\tau}^{n+1}} \varphi^n + A^{n+1} \frac{\tilde{\varphi}^{n+1} - \varphi^n}{\tilde{\tau}^{n+1}} \right),\end{aligned}$$

Error $\tilde{\psi}^{n+1}$ compare with $\tilde{\tau}^{n+1}$, and ψ^{n+1} with step τ^{n+1} :

$$\bar{\tau}^{n+1} = \gamma_{n+1} \tau^n, \quad \gamma_{n+1} = \frac{\delta}{\|\tilde{\psi}^{n+1}\|} \gamma.$$

Where $\varphi^n = \{\varphi^n, s^n\}$, $\psi^{n+1} = \{\psi_1^{n+1}, \psi_2^{n+1}\}$, $\tilde{\psi}^{n+1} = \{\tilde{\psi}_1^{n+1}, \tilde{\psi}_2^{n+1}\}$,

$$A = \begin{pmatrix} -\nabla \cdot D \nabla + \Sigma_a - (1 - \beta) \nu \Sigma_f - \lambda & 0 \\ 0 & \lambda - \beta \nu \Sigma_f \end{pmatrix}.$$

The needed time step

$$\tau^{n+1} \leq \bar{\tau}^{n+1}, \quad \tau^{n+1} \leq \tilde{\tau}^{n+1}, \quad \tau^{n+1} = \max \{ \tau^0, \min \{ \gamma_{n+1}, \gamma \} \tau^n \}.$$

The approximation error has the first order in time

$$\tilde{\psi}^{n+1} = \mathcal{O}(\tilde{\tau}_{n+1}).$$


In view of this, we set

$$\|\tilde{\psi}^{n+1}\| \leq \|(A^{n+1} - A^n)\varphi^n + A^{n+1}(\tilde{\varphi}^{n+1} - \varphi^n)\|.$$

Calculated formula for time step


$$\gamma_{n+1} = \frac{\delta}{\|(A^{n+1} - A^n)\varphi^n + A^{n+1}(\tilde{\varphi}^{n+1} - \varphi^n)\|} \gamma.$$

IAEA-2D benchmark



- One group of instantaneous neutrons
- One group of delayed neutrons
- Modeling effect of immersion or extraction of control rods

Nuclear power




Nuclear power for immersion (left) and extraction (right).

$$P(t) = a \sum_{g=1}^G \int_{\Omega} \Sigma_{fg} \phi_g d\mathbf{x},$$

where a – normalization factor.

Error




Error for immersion (left) and extraction (right).

$$\epsilon_P(t) = |P_{ref} - P|,$$

where P_{ref} – reference solution.

Time step



Time steps for immersion (left) and extraction (right).

Counting time and number of steps

	immersion			extraction		
δ	$\max(\epsilon_P)$	n	t , sec	$\max(\epsilon_P)$	n	t , sec
$4 \cdot 10^{-5}$	0.450	136	16	0.590	241	35
$2 \cdot 10^{-5}$	0.241	159	20	0.290	373	62
$1 \cdot 10^{-5}$	0.125	270	37	0.120	773	145

Reference solution: fixed time step 10^{-5} , number of steps – 50000, counting time – 2130 .

Conclusion

bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla
bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla
bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla
bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla bla

Thank you for your attention!