

# Automatic Time Step Selection for Numerical Solution of Neutron Diffusion Problems

Aleksandr Avvakumov <sup>1</sup>    Valery Strizhov <sup>2</sup>  
Petr Vabishchevich <sup>2,3</sup>    Aleksandr Vasilev <sup>3</sup>

<sup>1</sup>National Research Center “Kurchatov Institute“, Moscow, Russia

<sup>2</sup>Nuclear Safety Institute, Russian Academy of Sciences, Moscow, Russia

<sup>3</sup>North-Eastern Federal University, Yakutsk, Russia

7th FDM:T&A, June 11-16, 2018 Lozenetz

# Introduction

# Problem description

Consider second-order parabolic equation

$$\frac{\partial u}{\partial t} - \sum_{\alpha=1}^m \frac{\partial}{\partial x_{\alpha}} \left( k(\mathbf{x}, t) \frac{\partial u}{\partial x_{\alpha}} \right) + s(\mathbf{x}, t)u = f(\mathbf{x}, t), \quad \mathbf{x} \in \Omega, \quad 0 < t \leq T,$$

where  $\underline{k} \leq k(\mathbf{x}) \leq \bar{k}$ ,  $\mathbf{x} \in \Omega$ ,  $\underline{k} > 0$ .

# Problem description

Consider second-order parabolic equation

$$\frac{\partial u}{\partial t} - \sum_{\alpha=1}^m \frac{\partial}{\partial x_{\alpha}} \left( k(\mathbf{x}, t) \frac{\partial u}{\partial x_{\alpha}} \right) + s(\mathbf{x}, t)u = f(\mathbf{x}, t), \quad \mathbf{x} \in \Omega, \quad 0 < t \leq T,$$

where  $\underline{k} \leq k(\mathbf{x}) \leq \bar{k}$ ,  $\mathbf{x} \in \Omega$ ,  $\underline{k} > 0$ .

Boundary condition

$$u(\mathbf{x}, t) = g(\mathbf{x}, t), \quad \mathbf{x} \in \partial\Omega, \quad 0 < t \leq T.$$

Initial condition

$$u(\mathbf{x}, 0) = u^0(\mathbf{x}), \quad \mathbf{x} \in \Omega.$$

# Operator notation

Cauchy problem

$$\frac{du}{dt} + A(t)u = f(t), \quad 0 < t \leq T,$$

with initial condition

$$u(0) = u_0.$$

# Operator notation

Cauchy problem

$$\frac{du}{dt} + A(t)u = f(t), \quad 0 < t \leq T,$$

with initial condition

$$u(0) = u_0.$$

Assume  $A(t) \geq 0$  in  $H$  then

$$\|u(t)\| \leq \|u_0\| + \int_0^t \|f(\theta)\| d\theta.$$

# Solution evaluation

Introduce irregular time grid

$$t^0 = 0, \quad t^{n+1} = t^n + \tau^{n+1}, \quad n = 0, 1, \dots, N-1, \quad t^N = T.$$

# Solution evaluation

Introduce irregular time grid

$$t^0 = 0, \quad t^{n+1} = t^n + \tau^{n+1}, \quad n = 0, 1, \dots, N-1, \quad t^N = T.$$

Implicit scheme are used

$$\frac{y^{n+1} - y^n}{\tau^{n+1}} + A^{n+1}y^{n+1} = f^{n+1}, \quad n = 0, 1, \dots, N-1,$$

and initial condition  $y^0 = u^0$ .



# Solution evaluation

Introduce irregular time grid

$$t^0 = 0, \quad t^{n+1} = t^n + \tau^{n+1}, \quad n = 0, 1, \dots, N-1, \quad t^N = T.$$

Implicit scheme are used

$$\frac{y^{n+1} - y^n}{\tau^{n+1}} + A^{n+1}y^{n+1} = f^{n+1}, \quad n = 0, 1, \dots, N-1,$$

and initial condition  $y^0 = u^0$ .

Layerwise estimate ( $A^{n+1} \geq 0$ )

$$\|y^{n+1}\| \leq \|y^n\| + \tau^{n+1}\|f^{n+1}\|.$$

# Solution evaluation

Introduce irregular time grid

$$t^0 = 0, \quad t^{n+1} = t^n + \tau^{n+1}, \quad n = 0, 1, \dots, N-1, \quad t^N = T.$$

Implicit scheme are used

$$\frac{y^{n+1} - y^n}{\tau^{n+1}} + A^{n+1} y^{n+1} = f^{n+1}, \quad n = 0, 1, \dots, N-1,$$

and initial condition  $y^0 = u^0$ .

Layerwise estimate ( $A^{n+1} \geq 0$ )

$$\|y^{n+1}\| \leq \|y^n\| + \tau^{n+1} \|f^{n+1}\|.$$

Difference estimate

$$\|y^{n+1}\| \leq \|u^0\| + \sum_{k=0}^n \tau^{k+1} \|f^{k+1}\|.$$

# Solution error

For  $z^n = y^n - u^n$ :

$$\frac{z^{n+1} - z^n}{\tau^{n+1}} + A^{n+1} z^{n+1} = \psi^{n+1}, \quad n = 0, 1, \dots, N-1,$$

$$z^0 = 0.$$

# Solution error

For  $z^n = y^n - u^n$ :

$$\frac{z^{n+1} - z^n}{\tau^{n+1}} + A^{n+1} z^{n+1} = \psi^{n+1}, \quad n = 0, 1, \dots, N-1,$$

$$z^0 = 0.$$

Approximation error

$$\psi^{n+1} = f^{n+1} - \frac{u^{n+1} - u^n}{\tau^{n+1}} - A^{n+1} u^{n+1}.$$

# Solution error

For  $z^n = y^n - u^n$ :

$$\frac{z^{n+1} - z^n}{\tau^{n+1}} + A^{n+1} z^{n+1} = \psi^{n+1}, \quad n = 0, 1, \dots, N-1,$$

$$z^0 = 0.$$

Approximation error

$$\psi^{n+1} = f^{n+1} - \frac{u^{n+1} - u^n}{\tau^{n+1}} - A^{n+1} u^{n+1}.$$

Layerwise estimate

$$\|z^{n+1}\| \leq \sum_{k=0}^n \tau^{k+1} \|\psi^{k+1}\|.$$

# Solution error

For  $z^n = y^n - u^n$ :

$$\frac{z^{n+1} - z^n}{\tau^{n+1}} + A^{n+1} z^{n+1} = \psi^{n+1}, \quad n = 0, 1, \dots, N-1,$$

$$z^0 = 0.$$

Approximation error

$$\psi^{n+1} = f^{n+1} - \frac{u^{n+1} - u^n}{\tau^{n+1}} - A^{n+1} u^{n+1}.$$

Layerwise estimate

$$\|z^{n+1}\| \leq \sum_{k=0}^n \tau^{k+1} \|\psi^{k+1}\|.$$

Difference estimate

$$\|z^{n+1}\| \leq \delta t^{n+1}.$$

# Time step selection algorithm

- 1 Predictable time step:  $\tilde{\tau}^{n+1} = \gamma \tau^n$  (eg  $\gamma = 1.25$ )

# Time step selection algorithm

- 1 Predictable time step:  $\tilde{\tau}^{n+1} = \gamma \tau^n$  (eg  $\gamma = 1.25$ )
- 2 Predictive solution  $\tilde{y}^{n+1}$ : an explicit scheme,  $\tilde{t}^{n+1} = t^n + \tilde{\tau}^{n+1}$



# Time step selection algorithm

- 1 Predictable time step:  $\tilde{\tau}^{n+1} = \gamma \tau^n$  (eg  $\gamma = 1.25$ )
- 2 Predictive solution  $\tilde{y}^{n+1}$ : an explicit scheme,  $\tilde{t}^{n+1} = t^n + \tilde{\tau}^{n+1}$
- 3 Estimation of approximation error: by found  $\tilde{y}^{n+1}$  from an implicit scheme

# Time step selection algorithm

- 1 Predictable time step:  $\tilde{\tau}^{n+1} = \gamma \tau^n$  (eg  $\gamma = 1.25$ )
- 2 Predictive solution  $\tilde{y}^{n+1}$ : an explicit scheme,  $\tilde{t}^{n+1} = t^n + \tilde{\tau}^{n+1}$
- 3 Estimation of approximation error: by found  $\tilde{y}^{n+1}$  from an implicit scheme
- 4 Step selection  $\tau^{n+1}$ :  $\|\psi^{n+1}\| \approx \delta$

# Time step selection algorithm

- 1 Predictable time step:  $\tilde{\tau}^{n+1} = \gamma \tau^n$  (eg  $\gamma = 1.25$ )
- 2 Predictive solution  $\tilde{y}^{n+1}$ : an explicit scheme,  $\tilde{t}^{n+1} = t^n + \tilde{\tau}^{n+1}$
- 3 Estimation of approximation error: by found  $\tilde{y}^{n+1}$  from an implicit scheme
- 4 Step selection  $\tau^{n+1}$ :  $\|\psi^{n+1}\| \approx \delta$
- 5 Solution on a new time layer  $y^{n+1}$ : an implicit scheme,  $t^{n+1} = t^n + \tau^{n+1}$

# Neutron diffusion equation

One group diffusion approximation with one group delayed neutron sources

$$\frac{1}{\nu} \frac{\partial \phi}{\partial t} - \nabla \cdot D \nabla \phi + \Sigma_a \phi = (1 - \beta) \nu \Sigma_f \phi + \lambda c,$$

$$\frac{\partial c}{\partial t} + \lambda c = \beta \nu \Sigma_f \phi.$$

# Neutron diffusion equation

One group diffusion approximation with one group delayed neutron sources

$$\frac{1}{v} \frac{\partial \phi}{\partial t} - \nabla \cdot D \nabla \phi + \Sigma_a \phi = (1 - \beta) \nu \Sigma_f \phi + \lambda c,$$

$$\frac{\partial c}{\partial t} + \lambda c = \beta \nu \Sigma_f \phi.$$

Boundary condition

$$D \frac{\partial \phi}{\partial n} + \gamma \phi = 0.$$

Initial conditions

$$\phi(0) = \phi^0, c(0) = c^0.$$

## Calculated formulas

Denote vectors and matrix  $\varphi = \{\varphi, s\}$ ,  $\psi = \{\psi_1, \psi_2\}$ ,

$$A = \begin{pmatrix} -\nabla \cdot D \nabla + \Sigma_a - (1 - \beta) \nu \Sigma_f - \lambda & 0 \\ 0 & \lambda - \beta \nu \Sigma_f \end{pmatrix}.$$

## Calculated formulas

Denote vectors and matrix  $\varphi = \{\varphi, s\}$ ,  $\psi = \{\psi_1, \psi_2\}$ ,

$$A = \begin{pmatrix} -\nabla \cdot D \nabla + \Sigma_a - (1 - \beta) \nu \Sigma_f - \lambda & 0 \\ 0 & \lambda - \beta \nu \Sigma_f \end{pmatrix}.$$

The approximation error

$$\begin{aligned} \tilde{\psi}^{n+1} &= (A^{n+1} - A^n) \varphi^n + A^{n+1} (\tilde{\varphi}^{n+1} - \varphi^n) \\ &= \tilde{\tau}^{n+1} \left( \frac{A^{n+1} - A^n}{\tilde{\tau}^{n+1}} \varphi^n + A^{n+1} \frac{\tilde{\varphi}^{n+1} - \varphi^n}{\tilde{\tau}^{n+1}} \right). \end{aligned}$$

## Calculated formulas

Denote vectors and matrix  $\varphi = \{\varphi, s\}$ ,  $\psi = \{\psi_1, \psi_2\}$ ,

$$A = \begin{pmatrix} -\nabla \cdot D \nabla + \Sigma_a - (1 - \beta) \nu \Sigma_f - \lambda & 0 \\ 0 & \lambda - \beta \nu \Sigma_f \end{pmatrix}.$$

The approximation error

$$\begin{aligned} \tilde{\psi}^{n+1} &= (A^{n+1} - A^n) \varphi^n + A^{n+1} (\tilde{\varphi}^{n+1} - \varphi^n) \\ &= \tilde{\tau}^{n+1} \left( \frac{A^{n+1} - A^n}{\tilde{\tau}^{n+1}} \varphi^n + A^{n+1} \frac{\tilde{\varphi}^{n+1} - \varphi^n}{\tilde{\tau}^{n+1}} \right). \end{aligned}$$

Error  $\tilde{\psi}^{n+1}$  compare with  $\tilde{\tau}^{n+1}$ , and  $\psi^{n+1}$  with step  $\tau^{n+1}$ :

$$\tilde{\tau}^{n+1} = \gamma_{n+1} \tau^n, \quad \gamma_{n+1} = \frac{\delta}{\|\tilde{\psi}^{n+1}\|} \gamma.$$



## The needed time step

$$\tau^{n+1} \leq \bar{\tau}^{n+1}, \quad \tau^{n+1} \leq \tilde{\tau}^{n+1}, \quad \tau^{n+1} = \max \{ \tau^0, \min \{ \gamma_{n+1}, \gamma \} \tau^n \}.$$

The needed time step

$$\tau^{n+1} \leq \bar{\tau}^{n+1}, \quad \tau^{n+1} \leq \tilde{\tau}^{n+1}, \quad \tau^{n+1} = \max \{ \tau^0, \min \{ \gamma_{n+1}, \gamma \} \tau^n \}.$$

The approximation error has the first order in time

$$\tilde{\psi}^{n+1} = \mathcal{O}(\tilde{\tau}_{n+1}).$$

The needed time step

$$\tau^{n+1} \leq \bar{\tau}^{n+1}, \quad \tau^{n+1} \leq \tilde{\tau}^{n+1}, \quad \tau^{n+1} = \max \{ \tau^0, \min \{ \gamma_{n+1}, \gamma \} \tau^n \}.$$

The approximation error has the first order in time

$$\tilde{\psi}^{n+1} = \mathcal{O}(\tilde{\tau}_{n+1}).$$

In view of this, we set

$$\|\tilde{\psi}^{n+1}\| \leq \|(A^{n+1} - A^n)\varphi^n + A^{n+1}(\tilde{\varphi}^{n+1} - \varphi^n)\|.$$

The needed time step

$$\tau^{n+1} \leq \bar{\tau}^{n+1}, \quad \tau^{n+1} \leq \tilde{\tau}^{n+1}, \quad \tau^{n+1} = \max \{ \tau^0, \min \{ \gamma_{n+1}, \gamma \} \tau^n \}.$$

The approximation error has the first order in time

$$\tilde{\psi}^{n+1} = \mathcal{O}(\tilde{\tau}_{n+1}).$$


In view of this, we set

$$\|\tilde{\psi}^{n+1}\| \leq \|(A^{n+1} - A^n)\varphi^n + A^{n+1}(\tilde{\varphi}^{n+1} - \varphi^n)\|.$$

Calculated formula for time step


$$\gamma_{n+1} = \frac{\delta}{\|(A^{n+1} - A^n)\varphi^n + A^{n+1}(\tilde{\varphi}^{n+1} - \varphi^n)\|} \gamma.$$

# IAEA-2D benchmark




- Without reflector

# IAEA-2D benchmark



- Without reflector
- One group of *instantaneous* and *delayed* neutrons

# IAEA-2D benchmark



- Without reflector
- One group of *instantaneous* and *delayed* neutrons
- Modeling effect of *immersion* or *extraction* of control rods

# Scenario

Define the scenario of process:

- 1 The spectral problem is solved (initial condition);



# Scenario

Define the scenario of process:

- 1 The spectral problem is solved (initial condition);
- 2 Calculation for the non-stationary model in the range 0 to 0.5 sec;

# Scenario

Define the scenario of process:

- 1 The spectral problem is solved (initial condition);
- 2 Calculation for the non-stationary model in the range 0 to 0.5 sec;
- 3 At a moment of 0.1 sec the value  $\Sigma_a$  for the zone 3 changes to  $\pm 0.000625$ .

# Software



gmsh

# Software



gmsh

SLEPc

# Software



gmsh



FENICS  
PROJECT

# Software




gmsh



FENICS  
PROJECT



# Nuclear power




Nuclear power for immersion (left) and extraction (right).

$$P(t) = a \sum_{g=1}^G \int_{\Omega} \Sigma_{fg} \phi_g d\mathbf{x},$$

where  $a$  – normalization factor.

# Error




Error for immersion (left) and extraction (right).

$$\epsilon_P(t) = |P_{ref} - P|,$$

where  $P_{ref}$  – reference solution.



# Time step



Time steps for immersion (left) and extraction (right).

# Counting time and number of steps

	immersion			extraction		
$\delta$	$\max(\epsilon_P)$	$n$	$t$ , sec	$\max(\epsilon_P)$	$n$	$t$ , sec
$4 \cdot 10^{-5}$	0.450	136	16	0.590	241	35
$2 \cdot 10^{-5}$	0.241	159	20	0.290	373	62
$1 \cdot 10^{-5}$	0.125	270	37	0.120	773	145

Reference solution: fixed time step  $10^{-5}$ , number of steps – 50000, counting time – 2130 sec.

# Conclusion

- An algorithm for automatic time step selection for numerical solution of neutron diffusion problems has been developed.

# Conclusion

- An algorithm for automatic time step selection for numerical solution of neutron diffusion problems has been developed.
- The solution is obtained using guaranteed stable implicit schemes, and the step choice is performed with the use of the solution obtained by an explicit scheme.

# Conclusion

- An algorithm for automatic time step selection for numerical solution of neutron diffusion problems has been developed.
- The solution is obtained using guaranteed stable implicit schemes, and the step choice is performed with the use of the solution obtained by an explicit scheme.
- Calculation results obtained for a neutron diffusion problems demonstrate reliability of the proposed algorithm for time step choice.

# Conclusion

- An algorithm for automatic time step selection for numerical solution of neutron diffusion problems has been developed.
- The solution is obtained using guaranteed stable implicit schemes, and the step choice is performed with the use of the solution obtained by an explicit scheme.
- Calculation results obtained for a neutron diffusion problems demonstrate reliability of the proposed algorithm for time step choice.

Thank you for your attention!