

EXERCISES 3 (INTERSECTION THEORY)

Exercise 1. Let A be a domain with fraction field K , and L a finite field extension of K . Show that the integral closure of A in L is of finite type over A under any of the following assumptions:

- (i) A is integrally closed (in K), and L/K is separable.
- (ii) $A = k[t_1, \dots, t_n]$ and L/K purely inseparable.
- (iii) $A = k[t_1, \dots, t_n]$.
- (iv) A is a k -algebra of finite type.

Exercise 2. Let A be a domain with fraction field K . Show that A is integrally closed (in K) if and only if $A_{\mathfrak{p}}$ is integrally closed (in K) for every $\mathfrak{p} \in \text{Spec } A$.

A variety is called *normal* if the local rings $\mathcal{O}_{X,x}$ are integrally closed (in $K(X)$) for all points x of X .

Exercise 3. Let X be an integral variety, and $L/k(X)$ a finite field extension.

- (i) Show that there is a finite morphism of varieties $n: X' \rightarrow X$ with $k(X') = L$ as $k(X)$ -algebras and X' normal, with the following property. For any dominant morphism $f: Y \rightarrow X$ with Y a normal and integral variety, and any morphism of $k(X)$ -algebras $L \subset k(Y)$, there is a unique dominant morphism $g: Y \rightarrow X'$ such that $f = n \circ g$. The morphism n is the *normalisation of X in L* .
- (ii) Let $f: Y \rightarrow X$ be a finite dominant morphism with Y an integral and normal variety, and $k(Y) = L$ as $k(X)$ -algebras. Show that f is the normalisation of X in L .

Exercise 4. Let A be a noetherian local domain, which is integrally closed. Assume that the maximal ideal of A is an associated prime of the module A/aA , for some $a \in A$. Show that A has dimension one.

Exercise 5. Let X be a normal, integral, variety. Let D, D' be two effective Cartier divisors without common irreducible components. Show that $D \cap D' \rightarrow D$ is an effective Cartier divisor.