

Numerical Methods for PDE: Hyperbolic PDE

Note that ψ_m is an eigenvector for the forward difference operator D_h^+ , the backward difference operator D_h^- and the centered difference operator D_h . For example,

$$D_h^- \psi_m(x) = \frac{\psi_m(x) - \psi_m(x-h)}{h} = \frac{1 - e^{-2\pi i m h}}{h} \psi_m(x).$$

It is also an eigenvector for the forward and backward shift operators S_h^+ and S_h^- defined as

$$S_h^+ v(x) = v(x+h), \quad S_h^- v(x) = v(x-h),$$

where

$$\begin{aligned} S_h^+ \psi_m(x) &= e^{2\pi i m(x+h)} = e^{2\pi i m h} \psi_m(x), \\ S_h^- \psi_m(x) &= e^{2\pi i m(x-h)} = e^{-2\pi i m h} \psi_m(x). \end{aligned}$$

Now, consider forward-backward difference method in the operator form:

$$u^{j+1} = (1 - \lambda)u^j + \lambda S_h^- u^j = ((1 - \lambda)I + \lambda S_h^-)u^j.$$

Thus, for stability of the method, we need that the eigenvalues are less than or equal to one in magnitude, that is

$$|1 - \lambda + \lambda e^{-2\pi i m h}| \leq 1.$$

As $1 - \lambda + \lambda e^{-2\pi i m h}$ describes a circle centered at $1 - \lambda$ of radius $|\lambda|$, we see that the method is stable if and only if $0 \leq \lambda \leq 1$.

For the forward-centered method for advection equation, the eigen-values are

$$-\frac{\lambda}{2} e^{2\pi i m h} + 1 + \frac{\lambda}{2} e^{-2\pi i m h} = 1 - i\lambda \sin(2\pi m h).$$

Numerical Methods for PDE: Hyperbolic PDE



For the forward-centered method for advection equation, the eigenvalues are

$$-\frac{\lambda}{2}e^{2\pi imh} + 1 + \frac{\lambda}{2}e^{-2\pi imh} = 1 - i\lambda \sin(2\pi mh).$$

As every eigenvalue has magnitude greater than one, the method is indeed unconditionally unstable.

Numerical Methods for PDE: Hyperbolic PDE

For the forward-centered method for advection equation, the eigenvalues are

$$-\frac{\lambda}{2}e^{2\pi imh} + 1 + \frac{\lambda}{2}e^{-2\pi imh} = 1 - i\lambda \sin(2\pi mh).$$

As every eigenvalue has magnitude greater than one, the method is indeed unconditionally unstable.

Similarly, for forward-forward difference method, the eigenvalues are

$$-\lambda e^{2\pi imh} + 1 + \lambda$$

that lie on the circle centered at $1 + \lambda$ of radius $|\lambda|$. This confirms our earlier observation that, for every $\lambda > 0$, the method is unstable.

Numerical Methods for PDE: Hyperbolic PDE

For the forward-centered method for advection equation, the eigenvalues are

$$-\frac{\lambda}{2}e^{2\pi imh} + 1 + \frac{\lambda}{2}e^{-2\pi imh} = 1 - i\lambda \sin(2\pi mh).$$

As every eigenvalue has magnitude greater than one, the method is indeed unconditionally unstable.

Similarly, for forward-forward difference method, the eigenvalues are

$$-\lambda e^{2\pi imh} + 1 + \lambda$$

that lie on the circle centered at $1 + \lambda$ of radius $|\lambda|$. This confirms our earlier observation that, for every $\lambda > 0$, the method is unstable.

Let us now look at the backward-centered difference method for the advection equation given by

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^{j+1} - u_{n-1}^{j+1}}{2h} = 0.$$

Numerical Methods for PDE: Hyperbolic PDE

For the forward-centered method for advection equation, the eigenvalues are

$$-\frac{\lambda}{2}e^{2\pi imh} + 1 + \frac{\lambda}{2}e^{-2\pi imh} = 1 - i\lambda \sin(2\pi mh).$$

As every eigenvalue has magnitude greater than one, the method is indeed unconditionally unstable.

Similarly, for forward-forward difference method, the eigenvalues are

$$-\lambda e^{2\pi imh} + 1 + \lambda$$

that lie on the circle centered at $1 + \lambda$ of radius $|\lambda|$. This confirms our earlier observation that, for every $\lambda > 0$, the method is unstable.

Let us now look at the backward-centered difference method for the advection equation given by

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^{j+1} - u_{n-1}^{j+1}}{2h} = 0.$$

The method can be rewritten as

$$\frac{\lambda}{2}u_{n+1}^{j+1} + u_n^{j+1} - \frac{\lambda}{2}u_{n-1}^{j+1} = u_n^j$$

Numerical Methods for PDE: Hyperbolic PDE

For the forward-centered method for advection equation, the eigenvalues are

$$-\frac{\lambda}{2}e^{2\pi imh} + 1 + \frac{\lambda}{2}e^{-2\pi imh} = 1 - i\lambda \sin(2\pi mh).$$

As every eigenvalue has magnitude greater than one, the method is indeed unconditionally unstable.

Similarly, for forward-forward difference method, the eigenvalues are

$$-\lambda e^{2\pi imh} + 1 + \lambda$$

that lie on the circle centered at $1 + \lambda$ of radius $|\lambda|$. This confirms our earlier observation that, for every $\lambda > 0$, the method is unstable.

Let us now look at the backward-centered difference method for the advection equation given by

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^{j+1} - u_{n-1}^{j+1}}{2h} = 0.$$

The method can be rewritten as

$$\frac{\lambda}{2}u_{n+1}^{j+1} + u_n^{j+1} - \frac{\lambda}{2}u_{n-1}^{j+1} = u_n^j,$$

that is,

$$u^{j+1} = \left(\frac{\lambda}{2}S_h^+ + I - \frac{\lambda}{2}S_h^- \right)^{-1} u^j.$$

Numerical Methods for PDE: Hyperbolic PDE

Let us now look at the backward-centered difference method for the advection equation given by

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^{j+1} - u_{n-1}^{j+1}}{2h} = 0.$$

The method can be rewritten as

$$\frac{\lambda}{2} u_{n+1}^{j+1} + u_n^{j+1} - \frac{\lambda}{2} u_{n-1}^{j+1} = u_n^j,$$

that is,

$$u^{j+1} = \left(\frac{\lambda}{2} S_h^+ + I - \frac{\lambda}{2} S_h^- \right)^{-1} u^j.$$

Numerical Methods for PDE: Hyperbolic PDE

Let us now look at the backward-centered difference method for the advection equation given by

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^{j+1} - u_{n-1}^{j+1}}{2h} = 0.$$

The method can be rewritten as

$$\frac{\lambda}{2} u_{n+1}^{j+1} + u_n^{j+1} - \frac{\lambda}{2} u_{n-1}^{j+1} = u_n^j,$$

that is,

$$u^{j+1} = \left(\frac{\lambda}{2} S_h^+ + I - \frac{\lambda}{2} S_h^- \right)^{-1} u^j.$$

As all eigenvalues

$$\left(\frac{\lambda}{2} e^{2\pi i m h} + 1 - \frac{\lambda}{2} e^{-2\pi i m h} \right)^{-1} = \frac{1}{1 + i\lambda \sin(2\pi m h)}$$

are less than one in magnitude, the method is unconditionally stable.

Numerical Methods for PDE: Hyperbolic PDE

Let us now look at the backward-centered difference method for the advection equation given by

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^{j+1} - u_{n-1}^{j+1}}{2h} = 0.$$

The method can be rewritten as

$$\frac{\lambda}{2} u_{n+1}^{j+1} + u_n^{j+1} - \frac{\lambda}{2} u_{n-1}^{j+1} = u_n^j,$$

that is,

$$u^{j+1} = \left(\frac{\lambda}{2} S_h^+ + I - \frac{\lambda}{2} S_h^- \right)^{-1} u^j.$$

As all eigenvalues

$$\left(\frac{\lambda}{2} e^{2\pi i m h} + 1 - \frac{\lambda}{2} e^{-2\pi i m h} \right)^{-1} = \frac{1}{1 + i\lambda \sin(2\pi m h)}$$

are less than one in magnitude, the method is unconditionally stable.

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.



Numerical Methods for PDE: Hyperbolic PDE

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.

The scheme is given by

$$\frac{u_n^{j+1} - (u_{n+1}^j + u_{n-1}^j)/2}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0$$

Numerical Methods for PDE: Hyperbolic PDE

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.

The scheme is given by

$$\frac{u_n^{j+1} - (u_{n+1}^j + u_{n-1}^j)/2}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0,$$

or, in the operator form, reads

$$u^{j+1} = \left(\frac{1}{2} - \frac{\lambda}{2}\right) S_h^+ u^j + \left(\frac{1}{2} + \frac{\lambda}{2}\right) S_h^- u^j.$$

Numerical Methods for PDE: Hyperbolic PDE

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.

The scheme is given by

$$\frac{u_n^{j+1} - (u_{n+1}^j + u_{n-1}^j)/2}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0,$$

or, in the operator form, reads

$$u^{j+1} = \left(\frac{1}{2} - \frac{\lambda}{2}\right) S_h^+ u^j + \left(\frac{1}{2} + \frac{\lambda}{2}\right) S_h^- u^j.$$

The eigenvalues are

$$\left(\frac{1}{2} - \frac{\lambda}{2}\right) e^{2\pi i m h} + \left(\frac{1}{2} + \frac{\lambda}{2}\right) e^{-2\pi i m h} = \cos(2\pi m h) - i\lambda \sin(2\pi m h)$$

Numerical Methods for PDE: Hyperbolic PDE

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.

The scheme is given by

$$\frac{u_n^{j+1} - (u_{n+1}^j + u_{n-1}^j)/2}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0,$$

or, in the operator form, reads

$$u^{j+1} = \left(\frac{1}{2} - \frac{\lambda}{2}\right) S_h^+ u^j + \left(\frac{1}{2} + \frac{\lambda}{2}\right) S_h^- u^j.$$

The eigenvalues are

$$\left(\frac{1}{2} - \frac{\lambda}{2}\right) e^{2\pi i m h} + \left(\frac{1}{2} + \frac{\lambda}{2}\right) e^{-2\pi i m h} = \cos(2\pi m h) - i\lambda \sin(2\pi m h)$$

and, therefore, the method is stable if $|\lambda| \leq 1$.

Numerical Methods for PDE: Hyperbolic PDE

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.

The scheme is given by

$$\frac{u_n^{j+1} - (u_{n+1}^j + u_{n-1}^j)/2}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0,$$

or, in the operator form, reads

$$u^{j+1} = \left(\frac{1}{2} - \frac{\lambda}{2}\right) S_h^+ u^j + \left(\frac{1}{2} + \frac{\lambda}{2}\right) S_h^- u^j.$$

The eigenvalues are

$$\left(\frac{1}{2} - \frac{\lambda}{2}\right) e^{2\pi i m h} + \left(\frac{1}{2} + \frac{\lambda}{2}\right) e^{-2\pi i m h} = \cos(2\pi m h) - i\lambda \sin(2\pi m h)$$

and, therefore, the method is stable if $|\lambda| \leq 1$.

Remark

Note that the Lax-Friedrichs scheme can be rewritten as

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} - \frac{ch}{2\lambda} \frac{u_{n+1}^j - 2u_n^j + u_{n-1}^j}{h^2} = 0.$$

Numerical Methods for PDE: Hyperbolic PDE

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.

The scheme is given by

$$\frac{u_n^{j+1} - (u_{n+1}^j + u_{n-1}^j)/2}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0,$$

or, in the operator form, reads

$$u^{j+1} = \left(\frac{1}{2} - \frac{\lambda}{2}\right) S_h^+ u^j + \left(\frac{1}{2} + \frac{\lambda}{2}\right) S_h^- u^j.$$

The eigenvalues are

$$\left(\frac{1}{2} - \frac{\lambda}{2}\right) e^{2\pi i m h} + \left(\frac{1}{2} + \frac{\lambda}{2}\right) e^{-2\pi i m h} = \cos(2\pi m h) - i\lambda \sin(2\pi m h)$$

and, therefore, the method is stable if $|\lambda| \leq 1$.

Remark

Note that the Lax-Friedrichs scheme can be rewritten as

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} - \frac{ch}{2\lambda} \frac{u_{n+1}^j - 2u_n^j + u_{n-1}^j}{h^2} = 0.$$

Thus the method suggests discretization of the equation $\partial u / \partial t + c \partial u / \partial x - (ch/2\lambda) \partial^2 u / \partial x^2 = 0$. This is an advection-diffusion equation.

Numerical Methods for PDE: Hyperbolic PDE

As the final example, we look at the Lax-Friedrichs method which can be seen as a (conditionally) stable variant of the forward-centered difference scheme and maintains $O(k + h^2)$ accuracy.

The scheme is given by

$$\frac{u_n^{j+1} - (u_{n+1}^j + u_{n-1}^j)/2}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} = 0,$$

or, in the operator form, reads

$$u^{j+1} = \left(\frac{1}{2} - \frac{\lambda}{2}\right) S_h^+ u^j + \left(\frac{1}{2} + \frac{\lambda}{2}\right) S_h^- u^j.$$

The eigenvalues are

$$\left(\frac{1}{2} - \frac{\lambda}{2}\right) e^{2\pi i m h} + \left(\frac{1}{2} + \frac{\lambda}{2}\right) e^{-2\pi i m h} = \cos(2\pi m h) - i\lambda \sin(2\pi m h)$$

and, therefore, the method is stable if $|\lambda| \leq 1$.

Remark

Note that the Lax-Friedrichs scheme can be rewritten as

$$\frac{u_n^{j+1} - u_n^j}{k} + c \frac{u_{n+1}^j - u_{n-1}^j}{2h} - \frac{ch}{2\lambda} \frac{u_{n+1}^j - 2u_n^j + u_{n-1}^j}{h^2} = 0.$$

Thus the method suggests discretization of the equation $\partial u / \partial t + c \partial u / \partial x - (ch/2\lambda) \partial^2 u / \partial x^2 = 0$. This is an advection-diffusion equation. The method can be viewed as a forward-centered scheme with small diffusion added.