

# ***Boundary Value Problems: Shooting Method***



*There are a number of problems that can arise with the shooting method.*

# Boundary Value Problems: Shooting Method



*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge.*

# Boundary Value Problems: Shooting Method



*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge. For this reason, we may want to use the [modified Newton method](#).*

# Boundary Value Problems: Shooting Method

*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge. For this reason, we may want to use the [modified Newton method](#):*

$$s_{m+1} = s_m + \gamma d_m, \quad d_m = -\frac{h(s_m)}{h'(s_m)}$$

*and choose  $\gamma > 0$  to minimize*

$$|h(s_m + \gamma d_m)|$$

# Boundary Value Problems: Shooting Method



*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge. For this reason, we may want to use the **modified Newton method**:*

$$s_{m+1} = s_m + \gamma d_m, \quad d_m = -\frac{h(s_m)}{h'(s_m)}$$

*and choose  $\gamma > 0$  to minimize*

$$|h(s_m + \gamma d_m)| = |y(1; s_m + \gamma d_m) - (e + e^{-1})^{-1}|.$$

# Boundary Value Problems: Shooting Method



*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge. For this reason, we may want to use the **modified Newton method**:*

$$s_{m+1} = s_m + \gamma d_m, \quad d_m = -\frac{h(s_m)}{h'(s_m)}$$

*and choose  $\gamma > 0$  to minimize*

$$|h(s_m + \gamma d_m)| = |y(1; s_m + \gamma d_m) - (e + e^{-1})^{-1}|.$$

*Note that the choice  $\gamma = 1$  yields Newton's method.*

# Boundary Value Problems: Shooting Method



*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge. For this reason, we may want to use the modified Newton method:*

$$s_{m+1} = s_m + \gamma d_m, \quad d_m = -\frac{h(s_m)}{h'(s_m)}$$

*and choose  $\gamma > 0$  to minimize*

$$|h(s_m + \gamma d_m)| = |y(1; s_m + \gamma d_m) - (e + e^{-1})^{-1}|.$$

*Note that the choice  $\gamma = 1$  yields Newton's method.*

*A second problem is that the choice  $y^h(t; s)$  may be very sensitive to  $h$ ,  $s$  and other characteristics of the boundary value problem.*

# Boundary Value Problems: Shooting Method

*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge. For this reason, we may want to use the modified Newton method:*

$$s_{m+1} = s_m + \gamma d_m, \quad d_m = -\frac{h(s_m)}{h'(s_m)}$$

*and choose  $\gamma > 0$  to minimize*

$$|h(s_m + \gamma d_m)| = |y(1; s_m + \gamma d_m) - (e + e^{-1})^{-1}|.$$

*Note that the choice  $\gamma = 1$  yields Newton's method.*

*A second problem is that the choice  $y^h(t; s)$  may be very sensitive to  $h$ ,  $s$  and other characteristics of the boundary value problem.*

*For example,*

$$y'' = \lambda^2 y, \quad y(0) = 0, \quad y'(0) = s.$$

*Then*

$$y(t) =$$



# Boundary Value Problems: Shooting Method

*There are a number of problems that can arise with the shooting method.*

*There is no general guess  $s_0$  for the Newton iterations, and with a poor choice, the iteration may diverge. For this reason, we may want to use the modified Newton method:*

$$s_{m+1} = s_m + \gamma d_m, \quad d_m = -\frac{h(s_m)}{h'(s_m)}$$

*and choose  $\gamma > 0$  to minimize*

$$|h(s_m + \gamma d_m)| = |y(1; s_m + \gamma d_m) - (e + e^{-1})^{-1}|.$$

*Note that the choice  $\gamma = 1$  yields Newton's method.*

*A second problem is that the choice  $y^h(t; s)$  may be very sensitive to  $h$ ,  $s$  and other characteristics of the boundary value problem.*

*For example,*

$$y'' = \lambda^2 y, \quad y(0) = 0, \quad y'(0) = s.$$

*Then*

$$y(t) = \frac{s}{\lambda} (e^{\lambda t} - e^{-\lambda t}).$$

# *Numerical Analysis & Scientific Computing II*

## *Lesson 3*

# *Boundary Value Problems for ODEs*

*3.1 Well-posedness*

*3.2 Shooting Method*

***3.3 Finite Difference Method***



*Akash Anand*  
MATH, IIT KANPUR

# Boundary Value Problems: Finite Difference Method



*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

# Boundary Value Problems: Finite Difference Method



*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

*In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.*

# Boundary Value Problems: Finite Difference Method

*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

*In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.*

*Consider a scalar two-point boundary value problem*

$$u'' = f(t, u, u'), \quad a < t < b,$$

*with boundary conditions*

$$u(a) = \alpha, \quad u(b) = \beta.$$

# Boundary Value Problems: Finite Difference Method

*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

*In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.*

*Consider a scalar two-point boundary value problem*

$$u'' = f(t, u, u'), \quad a < t < b,$$

*with boundary conditions*

$$u(a) = \alpha, \quad u(b) = \beta.$$

*Introduce the mesh points  $t_i = a + ih, i = 0, \dots, n + 1$ , where  $h = (b - a)/(n + 1)$ .*

# Boundary Value Problems: Finite Difference Method

*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

*In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.*

*Consider a scalar two-point boundary value problem*

$$u'' = f(t, u, u'), \quad a < t < b,$$

*with boundary conditions*

$$u(a) = \alpha, \quad u(b) = \beta.$$

*Introduce the mesh points  $t_i = a + ih, i = 0, \dots, n + 1$ , where  $h = (b - a)/(n + 1)$ .*



# Boundary Value Problems: Finite Difference Method

*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

*In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.*

*Consider a scalar two-point boundary value problem*

$$u'' = f(t, u, u'), \quad a < t < b,$$

*with boundary conditions*

$$u(a) = \alpha, \quad u(b) = \beta.$$

*Introduce the mesh points  $t_i = a + ih, i = 0, \dots, n + 1$ , where  $h = (b - a)/(n + 1)$ .*

*We seek the approximations  $u_i$  to  $u(t_i)$ , that is,  $u_i \approx u(t_i)$ .*





# Boundary Value Problems: Finite Difference Method

*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

*In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.*

*Consider a scalar two-point boundary value problem*

$$u'' = f(t, u, u'), \quad a < t < b,$$

*with boundary conditions*

$$u(a) = \alpha, \quad u(b) = \beta.$$

*Introduce the mesh points  $t_i = a + ih, i = 0, \dots, n + 1$ , where  $h = (b - a)/(n + 1)$ .*

*We seek the approximations  $u_i$  to  $u(t_i)$ , that is,  $u_i \approx u(t_i)$ .*

*We replace the derivatives appearing in the ODE by finite difference approximations*

$$u'(t_i) \approx \frac{u_{i+1} - u_{i-1}}{2h}, \quad u''(t_i) \approx \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2}.$$



# Boundary Value Problems: Finite Difference Method

*In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.*

*Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.*

*In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.*

*Consider a scalar two-point boundary value problem*

$$u'' = f(t, u, u'), \quad a < t < b,$$

*with boundary conditions*

$$u(a) = \alpha, \quad u(b) = \beta.$$

*Introduce the mesh points  $t_i = a + ih, i = 0, \dots, n + 1$ , where  $h = (b - a)/(n + 1)$ .*

*We seek the approximations  $u_i$  to  $u(t_i)$ , that is,  $u_i \approx u(t_i)$ .*

*We replace the derivatives appearing in the ODE by finite difference approximations*

$$u'(t_i) \approx \frac{u_{i+1} - u_{i-1}}{2h}, \quad u''(t_i) \approx \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2}.$$

*This yields a system of algebraic equations*

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), \quad i = 1, \dots, n,$$

*to be solved for the unknowns  $u_i, i = 1, \dots, n$ .*

# Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns  $u_i, i = 1, \dots, n$ .

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \dots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix}$$

# Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns  $u_i, i = 1, \dots, n$ .

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \cdots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

# Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns  $u_i, i = 1, \dots, n$ .

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \dots & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

which is denoted as

$$\frac{1}{h^2} Au = F(u) + g.$$

# Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns  $u_i, i = 1, \dots, n$ .

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \dots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

which is denoted as

$$\frac{1}{h^2} Au = F(u) + g.$$

Thus, the Newton's method for solving the system of algebraic equations is given by

$$u^{(m+1)} = u^{(m)} - \left[ \frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[ \frac{1}{h^2} Au^{(m)} - F(u^{(m)}) - g \right]$$

where the Jacobian matrix is given by  $[F(u)]_{ij} = [\partial f(t_i, u_i, (u_{i+1} - u_{i-1})/(2h))/\partial u_j]$ .