

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.4 Implicit method

2.5 Stiffness

2.6 Linear Multistep Methods



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Linear Multistep Methods






Can we make the method higher order?

Initial Value Problems: Linear Multistep Methods



Can we make the method higher order?






Initial Value Problems: Linear Multistep Methods



Can we make the method higher order?


Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.



Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?


Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.



Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?

Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.




Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?

Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.


$$y_{n+1} = y_n + \int_{t_n}^{t_{n+1}} f(s, y(s)) ds \approx y_n + \int_{t_n}^{t_{n+1}} p(s) ds$$



Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?

Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.




Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?

Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.

$$\begin{aligned} y_{n+1} &= y_n + \int_{t_n}^{t_{n+1}} f(s, y(s)) ds \approx y_n + \int_{t_n}^{t_{n+1}} p(s) ds \\ &= y_n + b_{-1}f(t_{n+1}, y_{n+1}) + b_0f(t_n, y_n) + b_1f(t_{n-1}, y_{n-1}) \end{aligned}$$

Adams—Moulton
method




Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?

Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.

$$\begin{aligned} y_{n+1} &= y_n + \int_{t_n}^{t_{n+1}} f(s, y(s)) ds \approx y_n + \int_{t_n}^{t_{n+1}} p(s) ds \\ &= y_n + b_{-1}f(t_{n+1}, y_{n+1}) + b_0f(t_n, y_n) + b_1f(t_{n-1}, y_{n-1}) \end{aligned}$$

Adams—Moulton
method




An implicit
method

Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?


Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.



Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?


Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.



Initial Value Problems: Linear Multistep Methods

Can we make the method higher order?

Following the idea that we used to derive trapezoidal method, we could use higher order polynomial approximations.





Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.



Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.

For $k \geq 1$, that is, for a 2 or more step method, how do we start the time marching? Note that we need to know y_0, \dots, y_k , to compute y_{k+1} and we typically only know y_0 !

Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.

For $k \geq 1$, that is, for a 2 or more step method, how do we start the time marching? Note that we need to know y_0, \dots, y_k , to compute y_{k+1} and we typically only know y_0 !

... we use some other method such as a single step method.

Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.

For $k \geq 1$, that is, for a 2 or more step method, how do we start the time marching? Note that we need to know y_0, \dots, y_k , to compute y_{k+1} and we typically only know y_0 !

... we use some other method such as a single step method.

Examples:

1. $y_{n+1} = y_n + hf(t_n, y_n)$

... an explicit one step method

2. $y_{n+1} = y_n + hf(t_{n+1}, y_{n+1})$

... an implicit one step method

3. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_{n+1}))/2$

... an implicit one step method

Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.

For $k \geq 1$, that is, for a 2 or more step method, how do we start the time marching? Note that we need to know y_0, \dots, y_k , to compute y_{k+1} and we typically only know y_0 !

... we use some other method such as a single step method.

Examples:

1. $y_{n+1} = y_n + hf(t_n, y_n)$... an explicit one step method
2. $y_{n+1} = y_n + hf(t_{n+1}, y_{n+1})$... an implicit one step method
3. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_{n+1}))/2$... an implicit one step method
4. $y_{n+1} = y_n + b_0f(t_n, y_n) + b_1f(t_{n-1}, y_{n-1}) + b_2f(t_{n-2}, y_{n-2})$... an explicit **three** step method

Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.

For $k \geq 1$, that is, for a 2 or more step method, how do we start the time marching? Note that we need to know y_0, \dots, y_k , to compute y_{k+1} and we typically only know y_0 !

... we use some other method such as a single step method.

Examples:

1. $y_{n+1} = y_n + hf(t_n, y_n)$... an explicit one step method
2. $y_{n+1} = y_n + hf(t_{n+1}, y_{n+1})$... an implicit one step method
3. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_{n+1}))/2$... an implicit one step method
4. $y_{n+1} = y_n + b_0f(t_n, y_n) + b_1f(t_{n-1}, y_{n-1}) + b_2f(t_{n-2}, y_{n-2})$... an explicit **three** step method
5. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_n + hf(t_n, y_n)))/2$... an explicit one step method

Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.

For $k \geq 1$, that is, for a 2 or more step method, how do we start the time marching? Note that we need to know y_0, \dots, y_k , to compute y_{k+1} and we typically only know y_0 !

... we use some other method such as a single step method.

Improved Euler Method

Examples:

1. $y_{n+1} = y_n + hf(t_n, y_n)$... an explicit one step method
2. $y_{n+1} = y_n + hf(t_{n+1}, y_{n+1})$... an implicit one step method
3. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_{n+1}))$... an implicit one step method
4. $y_{n+1} = y_n + b_0f(t_n, y_n) + b_1f(t_{n-1}, y_{n-1}) + b_2f(t_{n-2}, y_{n-2})$... an explicit **three** step method
5. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_n + hf(t_n, y_n)))/2$... an explicit one step method

Initial Value Problems: Linear Multistep Methods

We consider methods that take constant step size h and determine y_{n+1} using the values from several preceding steps:

$$y_{n+1} = \Phi(f, t_n, y_{n+1}, y_n, y_{n-1}, \dots, y_{n-k}, h).$$

Here y_{n+1} depends on $k + 1$ previous values, so this is called a $(k + 1)$ -step method.

For $k \geq 1$, that is, for a 2 or more step method, how do we start the time marching? Note that we need to know y_0, \dots, y_k , to compute y_{k+1} and we typically only know y_0 !

... we use some other method such as a single step method.

Examples:

1. $y_{n+1} = y_n + hf(t_n, y_n)$
 2. $y_{n+1} = y_n + hf(t_{n+1}, y_{n+1})$
 3. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_{n+1}))$
 4. $y_{n+1} = y_n + b_0f(t_n, y_n) + b_1f(t_{n-1}, y_{n-1}) + b_2f(t_{n-2}, y_{n-2})$
 5. $y_{n+1} = y_n + h(f(t_n, y_n) + f(t_{n+1}, y_n + hf(t_n, y_n)))/2$
- Improved Euler Method*
 ... an explicit one step method
 ... an implicit one step method
 ... an implicit one step method
 ... an explicit **three** step method
 ... an explicit one step method
- Φ is linear in $y_n, f(t_n, y_n), f(t_{n+1}, y_{n+1}),$ etc.*
- non-linear Φ*



Initial Value Problems: Linear Multistep Methods

We consider linear multistep methods with constant step size, which by definition, are methods of the form

$$y_{n+1} = -a_0 y_n - a_1 y_{n-1} - \cdots - a_k y_{n-k} + h[b_{-1} f_{n+1} + b_0 f_n + \cdots + b_k f_{n-k}]$$

where f_n denotes $f(t_n, y_n)$ (for brevity) and a_j, b_j are constants which must be given and determine the specific method.



Initial Value Problems: Linear Multistep Methods

We consider linear multistep methods with constant step size, which by definition, are methods of the form

$$y_{n+1} = -a_0 y_n - a_1 y_{n-1} - \cdots - a_k y_{n-k} + h[b_{-1} f_{n+1} + b_0 f_n + \cdots + b_k f_{n-k}]$$

where f_n denotes $f(t_n, y_n)$ (for brevity) and a_j, b_j are constants which must be given and determine the specific method.

For explicit linear multistep method, $b_{-1} = 0$.

Initial Value Problems: Linear Multistep Methods

We consider linear multistep methods with constant step size, which by definition, are methods of the form

$$y_{n+1} = -a_0 y_n - a_1 y_{n-1} - \cdots - a_k y_{n-k} + h[b_{-1} f_{n+1} + b_0 f_n + \cdots + b_k f_{n-k}]$$

where f_n denotes $f(t_n, y_n)$ (for brevity) and a_j, b_j are constants which must be given and determine the specific method.

For explicit linear multistep method, $b_{-1} = 0$.

It is also convenient to define $a_{-1} = 1$, so that the method can be written more concisely as

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f_{n-j}.$$

Initial Value Problems: Linear Multistep Methods

We consider linear multistep methods with constant step size, which by definition, are methods of the form

$$y_{n+1} = -a_0 y_n - a_1 y_{n-1} - \cdots - a_k y_{n-k} + h[b_{-1} f_{n+1} + b_0 f_n + \cdots + b_k f_{n-k}]$$

where f_n denotes $f(t_n, y_n)$ (for brevity) and a_j, b_j are constants which must be given and determine the specific method.

For explicit linear multistep method, $b_{-1} = 0$.

It is also convenient to define $a_{-1} = 1$, so that the method can be written more concisely as

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f_{n-j}.$$

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Initial Value Problems: Linear Multistep Methods

We consider linear multistep methods with constant step size, which by definition, are methods of the form

$$y_{n+1} = -a_0 y_n - a_1 y_{n-1} - \cdots - a_k y_{n-k} + h[b_{-1} f_{n+1} + b_0 f_n + \cdots + b_k f_{n-k}]$$

where f_n denotes $f(t_n, y_n)$ (for brevity) and a_j, b_j are constants which must be given and determine the specific method.

For explicit linear multistep method, $b_{-1} = 0$.

It is also convenient to define $a_{-1} = 1$, so that the method can be written more concisely as

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f_{n-j}.$$

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Proof.

Initial Value Problems: Linear Multistep Methods

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Proof.

Define

$$F(z) = - \sum_{j=0}^k a_j y_{n-j} + h \sum_{j=0}^k b_j f(t_{n-j}, y_{n-j}) + h b_{-1} f(t_{n+1}, z).$$

Initial Value Problems: Linear Multistep Methods

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Proof.

Define

$$F(z) = - \sum_{j=0}^k a_j y_{n-j} + h \sum_{j=0}^k b_j f(t_{n-j}, y_{n-j}) + h b_{-1} f(t_{n+1}, z).$$

Now, $F(z)$ is Lipschitz with Lipschitz constant less than or equal to $h|b_{-1}|L$ (why?).

Initial Value Problems: Linear Multistep Methods

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Proof.

Define

$$F(z) = - \sum_{j=0}^k a_j y_{n-j} + h \sum_{j=0}^k b_j f(t_{n-j}, y_{n-j}) + h b_{-1} f(t_{n+1}, z).$$

Now, $F(z)$ is Lipschitz with Lipschitz constant less than or equal to $h|b_{-1}|L$ (why?). By hypothesis ($h < h_0$), the Lipschitz constant is strictly less than 1, that is, $F(z)$ is a contraction.

Initial Value Problems: Linear Multistep Methods

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Proof.

Define

$$F(z) = - \sum_{j=0}^k a_j y_{n-j} + h \sum_{j=0}^k b_j f(t_{n-j}, y_{n-j}) + h b_{-1} f(t_{n+1}, z).$$

Now, $F(z)$ is Lipschitz with Lipschitz constant less than or equal to $h|b_{-1}|L$ (why?). By hypothesis ($h < h_0$), the Lipschitz constant is strictly less than 1, that is, $F(z)$ is a contraction. The contraction mapping theorem then guarantees a unique fixed point, say y_{n+1} .

Initial Value Problems: Linear Multistep Methods

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Proof.

Define

$$F(z) = - \sum_{j=0}^k a_j y_{n-j} + h \sum_{j=0}^k b_j f(t_{n-j}, y_{n-j}) + h b_{-1} f(t_{n+1}, z).$$

Now, $F(z)$ is Lipschitz with Lipschitz constant less than or equal to $h|b_{-1}|L$ (why?). By hypothesis ($h < h_0$), the Lipschitz constant is strictly less than 1, that is, $F(z)$ is a contraction. The contraction mapping theorem then guarantees a unique fixed point, say y_{n+1} . Thus, we have $y_{n+1} = F(y_{n+1})$, as desired.

Initial Value Problems: Linear Multistep Methods

Theorem

Let $h_0 = 1/(|b_{-1}|L)$ where L is the Lipschitz constant for f . Then for any $h < h_0$ and any $y_n, y_{n-1}, \dots, y_{n-k}$, there is a unique y_{n+1} such that

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j}).$$

Proof.

Define

$$F(z) = - \sum_{j=0}^k a_j y_{n-j} + h \sum_{j=0}^k b_j f(t_{n-j}, y_{n-j}) + h b_{-1} f(t_{n+1}, z).$$

Now, $F(z)$ is Lipschitz with Lipschitz constant less than or equal to $h|b_{-1}|L$ (why?). By hypothesis ($h < h_0$), the Lipschitz constant is strictly less than 1, that is, $F(z)$ is a contraction. The contraction mapping theorem then guarantees a unique fixed point, say y_{n+1} . Thus, we have $y_{n+1} = F(y_{n+1})$, as desired.

Remark

The contraction mapping theorem also implies that the solution can be computed by fixed point iteration as is often done in practice. Moreover, only a fixed (small) number of iterations are made (introducing an additional error).

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.4 Implicit method

2.5 Stiffness

2.6 Linear Multistep Methods

- Adams methods



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Linear Multistep Methods



Examples


Adams Bashford methods -

Initial Value Problems: Linear Multistep Methods

Examples

Adams Bashford methods -

-- 2-step method




Initial Value Problems: Linear Multistep Methods

Examples

Adams Bashford methods -

-- 2-step method




Initial Value Problems: Linear Multistep Methods

Examples

Adams Bashford methods -

-- 2-step method




Initial Value Problems: Linear Multistep Methods

Examples

Adams Bashford methods -

-- 2-step method

$$\int_{t_n}^{t_{n+1}} p(t) dt$$



Initial Value Problems: Linear Multistep Methods


Examples

Adams Bashford methods -

-- 2-step method

$$\int_{t_n}^{t_{n+1}} p(t) dt =$$

$$\int_{t_n}^{t_{n+1}} \left(f_n \frac{t-t_{n-1}}{h} + f_{n-1} \frac{t_n-t}{h} \right) dt$$



Initial Value Problems: Linear Multistep Methods

Examples


Adams Bashford methods -

-- 2-step method

$$\int_{t_n}^{t_{n+1}} p(t) dt =$$

$$\int_{t_n}^{t_{n+1}} \left(f_n \frac{t-t_{n-1}}{h} + f_{n-1} \frac{t_n-t}{h} \right) dt =$$

$$f_n \left(\frac{3h}{2} \right) + f_{n-1} \left(-\frac{h}{2} \right)$$



Initial Value Problems: Linear Multistep Methods

Examples


Adams Bashford methods -

-- 2-step method

$$\begin{aligned} \int_{t_n}^{t_{n+1}} p(t) dt &= \\ \int_{t_n}^{t_{n+1}} \left(f_n \frac{t-t_{n-1}}{h} + f_{n-1} \frac{t_n-t}{h} \right) dt &= \\ f_n \left(\frac{3h}{2} \right) + f_{n-1} \left(-\frac{h}{2} \right) \end{aligned}$$

Thus,

$$y_{n+1} = y_n + \frac{h}{2} (3f_n - f_{n-1})$$



Initial Value Problems: Linear Multistep Methods

Examples

Adams Bashford methods -


-- 2-step method

$$\begin{aligned} \int_{t_n}^{t_{n+1}} p(t) dt &= \\ \int_{t_n}^{t_{n+1}} \left(f_n \frac{t-t_{n-1}}{h} + f_{n-1} \frac{t_n-t}{h} \right) dt &= \\ f_n \left(\frac{3h}{2} \right) + f_{n-1} \left(-\frac{h}{2} \right) \end{aligned}$$

Thus,

$$y_{n+1} = y_n + \frac{h}{2} (3f_n - f_{n-1})$$

-- General (k+1) step method



Initial Value Problems: Linear Multistep Methods

Examples

Adams Bashford methods -

-- 2-step method


$$\begin{aligned} \int_{t_n}^{t_{n+1}} p(t) dt &= \\ \int_{t_n}^{t_{n+1}} \left(f_n \frac{t-t_{n-1}}{h} + f_{n-1} \frac{t_n-t}{h} \right) dt &= \\ f_n \left(\frac{3h}{2} \right) + f_{n-1} \left(-\frac{h}{2} \right) \end{aligned}$$

Thus,

$$y_{n+1} = y_n + \frac{h}{2} (3f_n - f_{n-1})$$

-- General (k+1) step method

$$p(t) = \sum_{j=0}^k l_j^{(k)}(t) f_{n-j}, \quad \text{where} \quad l_j^{(k)}(t) = \prod_{i=0, i \neq j}^k (t - t_{n-i}) / (t_{n-j} - t_{n-i})$$



Initial Value Problems: Linear Multistep Methods

Examples

Adams Bashford methods -

-- 2-step method

$$\begin{aligned} \int_{t_n}^{t_{n+1}} p(t) dt &= \\ \int_{t_n}^{t_{n+1}} \left(f_n \frac{t-t_{n-1}}{h} + f_{n-1} \frac{t_n-t}{h} \right) dt &= \\ f_n \left(\frac{3h}{2} \right) + f_{n-1} \left(-\frac{h}{2} \right) \end{aligned}$$

Thus,

$$y_{n+1} = y_n + \frac{h}{2} (3f_n - f_{n-1})$$

-- General (k+1) step method


$$p(t) = \sum_{j=0}^k l_j^{(k)}(t) f_{n-j}, \quad \text{where}$$

$$l_j^{(k)}(t) = \prod_{i=0, i \neq j}^k (t - t_{n-i}) / (t_{n-j} - t_{n-i})$$

Thus,

$$y_{n+1} = y_n + \sum_{j=0}^k b_j f_{n-j}, \quad \text{with}$$

$$b_j = \int_{t_n}^{t_{n+1}} l_j^{(k)}(t) dt.$$



Initial Value Problems: Linear Multistep Methods



Examples


Adams Moulton methods -

Initial Value Problems: Linear Multistep Methods

Examples

Adams Moulton methods -

-- 2-step method




Initial Value Problems: Linear Multistep Methods

Examples

Adams Moulton methods -

-- 2-step method

$$\begin{aligned} p(t) = & f_{n+1} \frac{(t - t_n)(t - t_{n-1})}{2h^2} \\ & - f_n \frac{(t - t_{n+1})(t - t_{n-1})}{h^2} \\ & + f_{n-1} \frac{(t - t_{n+1})(t - t_n)}{2h^2} \end{aligned}$$



Initial Value Problems: Linear Multistep Methods


Examples

Adams Moulton methods -

-- 2-step method

$$\begin{aligned}
 p(t) = & f_{n+1} \frac{(t - t_n)(t - t_{n-1})}{2h^2} \\
 & - f_n \frac{(t - t_{n+1})(t - t_{n-1})}{h^2} \\
 & + f_{n-1} \frac{(t - t_{n+1})(t - t_n)}{2h^2}
 \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} p(t) dt$$



Initial Value Problems: Linear Multistep Methods


Examples

Adams Moulton methods -

-- 2-step method

$$\begin{aligned}
 p(t) = & f_{n+1} \frac{(t - t_n)(t - t_{n-1})}{2h^2} \\
 & - f_n \frac{(t - t_{n+1})(t - t_{n-1})}{h^2} \\
 & + f_{n-1} \frac{(t - t_{n+1})(t - t_n)}{2h^2}
 \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} p(t) dt = f_{n+1} \left(\frac{5h}{12} \right) - f_n \left(-\frac{2h}{3} \right) + f_{n-1} \left(-\frac{h}{12} \right)$$



Initial Value Problems: Linear Multistep Methods

Examples

Adams Moulton methods -


-- 2-step method

$$\begin{aligned}
 p(t) = & f_{n+1} \frac{(t - t_n)(t - t_{n-1})}{2h^2} \\
 & - f_n \frac{(t - t_{n+1})(t - t_{n-1})}{h^2} \\
 & + f_{n-1} \frac{(t - t_{n+1})(t - t_n)}{2h^2}
 \end{aligned}$$

$$\int_{t_n}^{t_{n+1}} p(t) dt = f_{n+1} \left(\frac{5h}{12} \right) - f_n \left(-\frac{2h}{3} \right) + f_{n-1} \left(-\frac{h}{12} \right)$$

Thus,

$$y_{n+1} = y_n + \frac{h}{12} (5f_{n+1} + 8f_n - f_{n-1})$$




Initial Value Problems: Linear Multistep Methods

Examples

Adams Moulton methods -

-- 2-step method

$$y_{n+1} = y_n + \frac{h}{12} (5f_{n+1} + 8f_n - f_{n-1})$$



-- General (k+1)-step method

$$p(t) = \sum_{j=-1}^k l_j^{(k)}(t) f_{n-j}, \quad \text{where} \quad l_j^{(k)}(t) = \prod_{i=-1, i \neq j}^k (t - t_{n-i}) / (t_{n-j} - t_{n-i})$$

Thus,

$$y_{n+1} = y_n + \sum_{j=-1}^k b_j f_{n-j}, \quad \text{with} \quad b_j = \int_{t_n}^{t_{n+1}} l_j^{(k)}(t) dt.$$

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.4 Implicit method

2.5 Stiffness

2.6 Linear Multistep Methods

- Consistency and Order



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Linear Multistep Methods

Consistency and Order

For the linear multistep method

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j})$$

define the local error as

$$\ell_{n+1}(y, h) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh)$$

for any $y \in C^1$, and $h > 0$.

Initial Value Problems: Linear Multistep Methods

Consistency and Order

For the linear multistep method

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j})$$

define the local error as

$$\ell_{n+1}(y, h) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh)$$

for any $y \in C^1$, and $h > 0$.

The linear multistep method is **consistent** if

$$\lim_{h \rightarrow 0} \max_{k \leq n < N} \left| \frac{\ell_{n+1}(y, h)}{h} \right| = 0$$

for all $y \in C^1$.