

# *Numerical Analysis & Scientific Computing II*

## *Lesson 5*

# *Integral Equations*



*Akash Anand*  
MATH, IIT KANPUR

# *Numerical Analysis & Scientific Computing II*

## *Lesson 5*

# *Integral Equations*


### *5.1 Some solutions of boundary value problems for PDEs via integral equations*



# Integral Equations: Some examples



*Some solutions of integral Fredholm integral equations*




(Akash Anand)

# Integral Equations: Some examples



*Some solutions of integral Fredholm integral equations in wave scattering*




(Ambuj Pandey,  
Akash Anand)

# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Awanish Tiwari,  
Ambuj Pandey,  
Akash Anand)



# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Awanish Tiwari,  
Ambuj Pandey,  
Akash Anand)

# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Awanish Tiwari,  
Ambuj Pandey,  
Akash Anand)

# Integral Equations: Some examples



*Some solutions of integral Fredholm integral equations in wave scattering*




(Stephen Lintner,  
Oscar Bruno)



# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Akash Anand)

# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Akash Anand)

# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Stephen Lintner,  
Oscar Bruno)

# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Akash Anand)

# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*




(Akash Anand)



# Integral Equations: Some examples



*Some solutions of integral Fredholm integral equations in wave scattering*




(Akash Anand)

# ***Integral Equations: Some examples***



*Some solutions of integral Fredholm integral equations in wave scattering*





(Akash Anand)

# Integral Equations: Some examples



*Some solutions of integral Fredholm integral equations in wave scattering*



(Akash Anand)

# *Numerical Analysis & Scientific Computing II*

## *Lesson 5*

# *Integral Equations*

*5.1 Some solutions of boundary value problems for PDEs via integral equations*

**5.2 An Introduction**



*Akash Anand*  
MATH, IIT KANPUR

# ***Integral Equations: An Introduction***



*We have already encountered integral equations in this course.*





# ***Integral Equations: An Introduction***

*We have already encountered integral equations in this course.*

*Recall that the initial value problem*

$$y' = f(t, y), y(t_0) = y_0,$$

*is equivalent to finding  $y$  satisfying*

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds.$$

# Integral Equations: An Introduction

*We have already encountered integral equations in this course.*

*Recall that the initial value problem*

$$y' = f(t, y), y(t_0) = y_0,$$

*is equivalent to finding  $y$  satisfying*

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds.$$

*This is a particular case of the more general **Volterra integral equations***

$$\int_{t_0}^t K(t, s, y(s)) ds = g(t), \quad t \geq t_0.$$

*where  $K$  and  $g$  are given functions.*

# Integral Equations: An Introduction

We have already encountered integral equations in this course.

Recall that the initial value problem

$$y' = f(t, y), y(t_0) = y_0,$$

is equivalent to finding  $y$  satisfying

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds.$$

This is a particular case of the more general **Volterra integral equations**

$$\int_{t_0}^t K(t, s, y(s)) ds = g(t), \quad t \geq t_0.$$

where  $K$  and  $g$  are given functions. The general **linear Volterra equation of the first kind** is of the form

$$\int_{t_0}^t K(t, s) y(s) ds = g(t), \quad t \geq t_0.$$

# Integral Equations: An Introduction

*We have already encountered integral equations in this course.*

*Recall that the initial value problem*

$$y' = f(t, y), y(t_0) = y_0,$$

*is equivalent to finding  $y$  satisfying*

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds.$$

*This is a particular case of the more general **Volterra integral equations***

$$\int_{t_0}^t K(t, s, y(s)) ds = g(t), \quad t \geq t_0.$$

*where  $K$  and  $g$  are given functions. The general **linear Volterra equation of the first kind** is of the form*

$$\int_{t_0}^t K(t, s) y(s) ds = g(t), \quad t \geq t_0.$$

*The difficulty with these equations, linear or non-linear, is that they are ill-conditioned.*

# Integral Equations: An Introduction

We have already encountered integral equations in this course.

Recall that the initial value problem

$$y' = f(t, y), y(t_0) = y_0,$$

is equivalent to finding  $y$  satisfying

$$y(t) = y_0 + \int_{t_0}^t f(s, y(s)) ds.$$

This is a particular case of the more general **Volterra integral equations**

$$\int_{t_0}^t K(t, s, y(s)) ds = g(t), \quad t \geq t_0.$$

where  $K$  and  $g$  are given functions. The general **linear Volterra equation of the first kind** is of the form

$$\int_{t_0}^t K(t, s) y(s) ds = g(t), \quad t \geq t_0.$$

The difficulty with these equations, linear or non-linear, is that they are ill-conditioned.

The numerical solution of these equations are closely related to the initial value problem. We will, however, focus on a different type of integral equations known as Fredholm integral equations, in particular, of the second kind.



# Integral Equations: An Introduction



The general form of such an integral equation is

$$u(t) - \int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

where  $\Omega$  is a bounded set in  $\mathbb{R}^m$ ,  $m \geq 1$ .

# Integral Equations: An Introduction

The general form of such an integral equation is

$$u(t) - \int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

where  $\Omega$  is a bounded set in  $\mathbb{R}^m$ ,  $m \geq 1$ . The general form of *Fredholm integral equation of the first kind* is

$$\int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

# Integral Equations: An Introduction

The general form of such an integral equation is

$$u(t) - \int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

where  $\Omega$  is a bounded set in  $\mathbb{R}^m$ ,  $m \geq 1$ . The general form of *Fredholm integral equation of the first kind* is

$$\int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

The Fredholm integral equations are widely studied and applied in connection with solving boundary value problems for elliptic PDEs (they are also used in connection with other PDEs).

# Integral Equations: An Introduction

The general form of such an integral equation is

$$u(t) - \int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

where  $\Omega$  is a bounded set in  $\mathbb{R}^m$ ,  $m \geq 1$ . The general form of *Fredholm integral equation of the first kind* is

$$\int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

The Fredholm integral equations are widely studied and applied in connection with solving boundary value problems for elliptic PDEs (they are also used in connection with other PDEs).

## Example

Consider solving the problem

$$\begin{aligned} \Delta u(x) &= 0, & x \in \Omega, \\ u(x) &= g(x), & x \in \Gamma, \end{aligned}$$

where  $\Omega$  is a bounded region in  $\mathbb{R}^3$  with non-empty interior and  $\Gamma$  is the boundary of  $\Omega$ .

# Integral Equations: An Introduction

The general form of such an integral equation is

$$u(t) - \int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

where  $\Omega$  is a bounded set in  $\mathbb{R}^m$ ,  $m \geq 1$ . The general form of *Fredholm integral equation of the first kind* is

$$\int_{\Omega} K(t, s)u(s)ds = f(t), \quad t \in \Omega.$$

The Fredholm integral equations are widely studied and applied in connection with solving boundary value problems for elliptic PDEs (they are also used in connection with other PDEs).

## Example

Consider solving the problem

$$\begin{aligned} \Delta u(x) &= 0, & x \in \Omega, \\ u(x) &= g(x), & x \in \Gamma, \end{aligned}$$

where  $\Omega$  is a bounded region in  $\mathbb{R}^3$  with non-empty interior and  $\Gamma$  is the boundary of  $\Omega$ . If we seek  $u$  in the form of a *single layer potential*, that is

$$u(x) = \int_{\Gamma} \frac{1}{|x - y|} \rho(y) dy, \quad x \in \Omega,$$

where  $\rho(y)$  is called a *single layer density* function and it is the unknown in the equation.



# Integral Equations: An Introduction

## Example

Consider solving the problem

$$\begin{aligned}\Delta u(x) &= 0, & x \in \Omega, \\ u(x) &= g(x), & x \in \Gamma,\end{aligned}$$

where  $\Omega$  is a bounded region in  $\mathbb{R}^3$  with non-empty interior and  $\Gamma$  is the boundary of  $\Omega$ . If we seek  $u$  in the form of a *single layer potential*, that is

$$u(x) = \int_{\Gamma} \frac{1}{|x - y|} \rho(y) dy, \quad x \in \Omega,$$

where  $\rho(y)$  is called a *single layer density* function and it is the unknown in the equation.

Using the boundary condition (and the fact from “Potential Theory” that single layer potentials are continuous everywhere), it is straightforward to see that  $\rho$  satisfies

$$\int_{\Gamma} \frac{1}{|x - y|} \rho(y) dy = g(x), \quad x \in \Gamma,$$

a Fredholm integral equation of the first kind.

# Integral Equations: An Introduction

## Example

Consider solving the problem

$$\begin{aligned}\Delta u(x) &= 0, & x \in \Omega, \\ u(x) &= g(x), & x \in \Gamma,\end{aligned}$$

where  $\Omega$  is a bounded region in  $\mathbb{R}^3$  with non-empty interior and  $\Gamma$  is the boundary of  $\Omega$ . If we seek  $u$  in the form of a *single layer potential*, that is

$$u(x) = \int_{\Gamma} \frac{1}{|x - y|} \rho(y) dy, \quad x \in \Omega,$$

where  $\rho(y)$  is called a *single layer density* function and it is the unknown in the equation.

Using the boundary condition (and the fact from “Potential Theory” that single layer potentials are continuous everywhere), it is straightforward to see that  $\rho$  satisfies

$$\int_{\Gamma} \frac{1}{|x - y|} \rho(y) dy = g(x), \quad x \in \Gamma,$$

a Fredholm integral equation of the first kind. If we see the solution in the form of a *double layer potential*

$$u(x) = \int_{\Gamma} \frac{\partial}{\partial \nu(y)} \left( \frac{1}{|x - y|} \right) \mu(y) dy, \quad x \in \Omega,$$

then  $\mu(y)$  satisfies a Fredholm integral equation of the second kind.

# Integral Equations: An Introduction

## Example

Consider solving the problem

$$\begin{aligned}\Delta u(x) &= 0, & x \in \Omega, \\ u(x) &= g(x), & x \in \Gamma,\end{aligned}$$

where  $\Omega$  is a bounded region in  $\mathbb{R}^3$  with non-empty interior and  $\Gamma$  is the boundary of  $\Omega$ . If we see the solution in the form of a **double layer potential**

$$u(x) = \int_{\Gamma} \frac{\partial}{\partial \nu(y)} \left( \frac{1}{|x - y|} \right) \mu(y) dy, \quad x \in \Omega,$$

then  $\mu(y)$  satisfies a Fredholm integral equation of the second kind.

# Integral Equations: An Introduction

## Example

Consider solving the problem

$$\begin{aligned}\Delta u(x) &= 0, & x \in \Omega, \\ u(x) &= g(x), & x \in \Gamma,\end{aligned}$$

where  $\Omega$  is a bounded region in  $\mathbb{R}^3$  with non-empty interior and  $\Gamma$  is the boundary of  $\Omega$ . If we see the solution in the form of a **double layer potential**

$$u(x) = \int_{\Gamma} \frac{\partial}{\partial \nu(y)} \left( \frac{1}{|x - y|} \right) \mu(y) dy, \quad x \in \Omega,$$

then  $\mu(y)$  satisfies a Fredholm integral equation of the second kind. Indeed, the **double layer density** function  $\mu(y)$  satisfies

$$\frac{1}{2} \mu(x) - \int_{\Gamma} \frac{\partial}{\partial \nu(y)} \left( \frac{1}{|x - y|} \right) \mu(y) dy = -g(x), \quad x \in \Gamma,$$

(another fact from “Potential Theory” known as jump relation for double layer potential).



# Integral Equations: An Introduction

We say that a kernel  $K: \Omega \times \Omega \rightarrow \mathbb{C}$  is **weakly singular** if  $K$  is defined and continuous for all  $x, y \in \Omega \subseteq \mathbb{R}^m$ ,  $x \neq y$ , and there exist positive constants  $M$  and  $\alpha \in (0, m]$  such that

$$|K(x, y)| \leq M|x - y|^{-(m-\alpha)}$$

for all  $x, y \in \Omega$ ,  $x \neq y$ .

# Integral Equations: An Introduction

We say that a kernel  $K: \Omega \times \Omega \rightarrow \mathbb{C}$  is **weakly singular** if  $K$  is defined and continuous for all  $x, y \in \Omega \subseteq \mathbb{R}^m$ ,  $x \neq y$ , and there exist positive constants  $M$  and  $\alpha \in (0, m]$  such that

$$|K(x, y)| \leq M|x - y|^{-(m-\alpha)}$$

for all  $x, y \in \Omega$ ,  $x \neq y$ .

## Remark

One can show that, the integral operator

$$(Au)(x) = \int_{\Omega} K(x, y)u(y)dy,$$

with weakly singular kernel  $K$  maps continuous functions to continuous function, that is, the operator  $A: C(\Omega) \rightarrow C(\Omega)$  is well-defined.

# Integral Equations: An Introduction

We say that a kernel  $K: \Omega \times \Omega \rightarrow \mathbb{C}$  is **weakly singular** if  $K$  is defined and continuous for all  $x, y \in \Omega \subseteq \mathbb{R}^m$ ,  $x \neq y$ , and there exist positive constants  $M$  and  $\alpha \in (0, m]$  such that

$$|K(x, y)| \leq M|x - y|^{-(m-\alpha)}$$

for all  $x, y \in \Omega$ ,  $x \neq y$ .

## Remark

One can show that, the integral operator

$$(Au)(x) = \int_{\Omega} K(x, y)u(y)dy,$$

with weakly singular kernel  $K$  maps continuous functions to continuous function, that is, the operator  $A: C(\Omega) \rightarrow C(\Omega)$  is well-defined.

Moreover, it is known that, for such integral operators, the **Fredholm alternative** holds, that is,

$$(I - A)u = f$$

has a unique solution for every  $f \in C(\Omega)$  if and only if the homogeneous equation  $(I - A)v = 0$  has only the trivial solution  $v = 0$ .