

Numerical Analysis & Scientific Computing II

Lesson 5

Integral Equations

5.1 Some solutions of boundary value problems for PDEs via integral equations

5.2 An Introduction

5.3 Numerical Methods



Akash Anand
MATH, IIT KANPUR

Integral Equations: An Introduction

There are three main ideas for numerical solution of the second kind Fredholm integral equation

$$(I - A)u = f$$

with the linear integral operator

$$(Au)(x) = \int_{\Omega} K(x, y)u(y)dy .$$

Approximate the integral operator by

- *approximating the kernel $K(x, y)$.*
- *approximating the solution $u(x)$.*
- *approximating the integral $\int_{\Omega} f(y)dy$ by a quadrature.*

Numerical Analysis & Scientific Computing II

Lesson 5

Integral Equations

5.2 An Introduction

5.3 Numerical Methods

- Degenerate Kernel Method



Integral Equations: Numerical Methods



The first class of methods, called the *degenerate kernel method*, is based on the approximation to the kernel K in the integral operator.



Integral Equations: Numerical Methods

The first class of methods, called the *degenerate kernel method*, is based on the approximation to the kernel K in the integral operator.

More precisely, the kernel is approximated by a degenerate kernel of the form

$$K(x, y) \approx K_n(x, y) = \sum_{j=1}^n a_j(x) b_j(y).$$

Integral Equations: Numerical Methods

The first class of methods, called the *degenerate kernel method*, is based on the approximation to the kernel K in the integral operator.

More precisely, the kernel is approximated by a degenerate kernel of the form

$$K(x, y) \approx K_n(x, y) = \sum_{j=1}^n a_j(x) b_j(y).$$

The solution of the integral equation of the second kind, $u - Au = f$, is then obtained as

$$u_n(x) - \sum_{j=1}^n \langle b_j, u_n \rangle a_j(x) = f(x),$$
$$\langle f, g \rangle = \int_{\Omega} f(y) g(y) dy.$$

Integral Equations: Numerical Methods

The first class of methods, called the *degenerate kernel method*, is based on the approximation to the kernel K in the integral operator.

More precisely, the kernel is approximated by a degenerate kernel of the form

$$K(x, y) \approx K_n(x, y) = \sum_{j=1}^n a_j(x) b_j(y).$$

The solution of the integral equation of the second kind, $u - Au = f$, is then obtained as

$$u_n(x) - \sum_{j=1}^n \langle b_j, u_n \rangle a_j(x) = f(x),$$
$$\langle f, g \rangle = \int_{\Omega} f(y) g(y) dy.$$

Thus, the approximate solution has the form

$$u_n = f + \sum_{k=1}^n \gamma_k a_k,$$

Integral Equations: Numerical Methods

The first class of methods, called the **degenerate kernel method**, is based on the approximation to the kernel K in the integral operator.

More precisely, the kernel is approximated by a degenerate kernel of the form

$$K(x, y) \approx K_n(x, y) = \sum_{j=1}^n a_j(x) b_j(y).$$

The solution of the integral equation of the second kind, $u - Au = f$, is then obtained as

$$u_n(x) - \sum_{j=1}^n \langle b_j, u_n \rangle a_j(x) = f(x),$$

$$\langle f, g \rangle = \int_{\Omega} f(y) g(y) dy.$$

Thus, the approximate solution has the form

$$u_n = f + \sum_{k=1}^n \gamma_k a_k,$$

where the coefficients $\gamma_1, \gamma_2, \dots, \gamma_n$ satisfy the linear system

$$\gamma_j - \sum_{k=1}^n \langle a_k, b_j \rangle \gamma_k = \langle f, b_j \rangle, \quad j = 1, 2, \dots, n.$$

Integral Equations: Numerical Methods



How do we construct such an approximation for the kernel? How do we know that this approximation is accurate?

Integral Equations: Numerical Methods



How do we construct such an approximation for the kernel? How do we know that this approximation is accurate?

The idea is to construct K_n such that $\|A_n - A\| \rightarrow 0$ as $n \rightarrow \infty$ where

$$(A_n u)(x) = \int_{\Omega} K_n(x, y) u(y) dy.$$

Integral Equations: Numerical Methods



How do we construct such an approximation for the kernel? How do we know that this approximation is accurate?

The idea is to construct K_n such that $\|A_n - A\| \rightarrow 0$ as $n \rightarrow \infty$ where

$$(A_n u)(x) = \int_{\Omega} K_n(x, y) u(y) dy.$$

Note that

$$\|A_n - A\|_{\infty} = \max_{x \in \Omega} \int_{\Omega} |K_n(x, y) - K(x, y)| dy.$$

Integral Equations: Numerical Methods

How do we construct such an approximation for the kernel? How do we know that this approximation is accurate?

The idea is to construct K_n such that $\|A_n - A\| \rightarrow 0$ as $n \rightarrow \infty$ where

$$(A_n u)(x) = \int_{\Omega} K_n(x, y) u(y) dy.$$

Note that

$$\|A_n - A\|_{\infty} = \max_{x \in \Omega} \int_{\Omega} |K_n(x, y) - K(x, y)| dy.$$

Theorem

Let X and Y be Banach spaces and let $A: X \rightarrow Y$ be a bounded linear operator with a bounded operator $A^{-1}: Y \rightarrow X$. Assume the sequence $A_n: X \rightarrow Y$ of bounded linear operators to be norm convergent, that is, $\|A_n - A\| \rightarrow 0$ as $n \rightarrow \infty$. Then, for sufficiently large n , more precisely, for all n with $\|A^{-1}(A_n - A)\| < 1$, the inverse operators $A_n^{-1}: Y \rightarrow X$ exist and are bounded by

$$\|A_n^{-1}\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1}(A_n - A)\|}.$$

For all solutions of the equations $A\varphi = f$ and $A_n\varphi_n = f_n$, we have the error estimate

$$\|\varphi_n - \varphi\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1}(A_n - A)\|} \{\|(A_n - A)\varphi\| + \|f_n - f\|\}.$$

Numerical Analysis & Scientific Computing II

Lesson 5

Integral Equations

5.2 An Introduction

5.3 Numerical Methods

- Degenerate Kernel Method**
- via interpolation**



Integral Equations: Numerical Methods

Theorem

Let X and Y be Banach spaces and let $A: X \rightarrow Y$ be a bounded linear operator with a bounded operator $A^{-1}: Y \rightarrow X$. Assume the sequence $A_n: X \rightarrow Y$ of bounded linear operators to be norm convergent, that is, $\|A_n - A\| \rightarrow 0$ as $n \rightarrow \infty$. Then, for sufficiently large n , more precisely, for all n with $\|A^{-1}(A_n - A)\| < 1$, the inverse operators $A_n^{-1}: Y \rightarrow X$ exist and are bounded by

$$\|A_n^{-1}\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1}(A_n - A)\|}.$$

For all solutions of the equations $A\varphi = f$ and $A_n\varphi_n = f_n$, we have the error estimate

$$\|\varphi_n - \varphi\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1}(A_n - A)\|} \{\|(A_n - A)\varphi\| + \|f_n - f\|\}.$$

Consider the domain to be an interval, that is, $\Omega = (a, b)$ and let K be continuous.

Integral Equations: Numerical Methods

Theorem

Let X and Y be Banach spaces and let $A: X \rightarrow Y$ be a bounded linear operator with a bounded operator $A^{-1}: Y \rightarrow X$. Assume the sequence $A_n: X \rightarrow Y$ of bounded linear operators to be norm convergent, that is, $\|A_n - A\| \rightarrow 0$ as $n \rightarrow \infty$. Then, for sufficiently large n , more precisely, for all n with $\|A^{-1}(A_n - A)\| < 1$, the inverse operators $A_n^{-1}: Y \rightarrow X$ exist and are bounded by

$$\|A_n^{-1}\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1}(A_n - A)\|}.$$

For all solutions of the equations $A\varphi = f$ and $A_n\varphi_n = f_n$, we have the error estimate

$$\|\varphi_n - \varphi\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1}(A_n - A)\|} \{\|(A_n - A)\varphi\| + \|f_n - f\|\}.$$

Consider the domain to be an interval, that is, $\Omega = (a, b)$ and let K be continuous.

One idea that works when K is continuous is interpolation -- approximate K by interpolating in x with respect to the points x_1, x_2, \dots, x_n in $[a, b]$ for each $y \in [a, b]$, we have


$$K_n(x, y) = \sum_{j=1}^n K(x_j, y) l_j^{(n)}(x).$$

Integral Equations: Numerical Methods

Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$



Integral Equations: Numerical Methods


Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the resulting linear system $(I - M)\gamma = F$ is given by

$$m_{ij} = \langle a_j, b_i \rangle = \int_a^b a_j(x) b_i(x) dx$$



Integral Equations: Numerical Methods


Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the resulting linear system $(I - M)\gamma = F$ is given by

$$m_{ij} = \langle a_j, b_i \rangle = \int_a^b a_j(x) b_i(x) dx = \int_a^b l_j^{(n)}(x) K(x_i, x) dx$$



Integral Equations: Numerical Methods


Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$

Then, the resulting linear system $(I - M)\gamma = F$ is given by

$$m_{ij} = \langle a_j, b_i \rangle = \int_a^b a_j(x) b_i(x) dx = \int_a^b l_j^{(n)}(x) K(x_i, x) dx \approx \sum_{k=1}^n K(x_i, x_k) \int_a^b l_j^{(n)}(x) l_k^{(n)}(x) dx$$




Integral Equations: Numerical Methods

Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$



Then, the resulting linear system $(I - M)\gamma = F$ is given by

$$m_{ij} = \langle a_j, b_i \rangle = \int_a^b a_j(x) b_i(x) dx = \int_a^b l_j^{(n)}(x) K(x_i, x) dx \approx \sum_{k=1}^n K(x_i, x_k) \int_a^b l_j^{(n)}(x) l_k^{(n)}(x) dx = \sum_{k=1}^n K(x_i, x_k) w_{jk},$$

where


$$W = \frac{h}{6} \begin{bmatrix} 2 & 1 & & & \\ 1 & 4 & 1 & & \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ & & 1 & 4 & 1 \\ & & & 1 & 2 \end{bmatrix}.$$

Integral Equations: Numerical Methods

Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$



Then, the resulting linear system $(I - M)\gamma = F$ is given by

$$m_{ij} = \langle a_j, b_i \rangle = \int_a^b a_j(x) b_i(x) dx = \int_a^b l_j^{(n)}(x) K(x_i, x) dx \approx \sum_{k=1}^n K(x_i, x_k) \int_a^b l_j^{(n)}(x) l_k^{(n)}(x) dx = \sum_{k=1}^n K(x_i, x_k) w_{jk},$$

where

$$W = \frac{h}{6} \begin{bmatrix} 2 & 1 & & & \\ 1 & 4 & 1 & & \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ & 1 & 4 & 1 & \\ & & 1 & 2 & \end{bmatrix}.$$

Similarly,


$$F_i = \langle f, b_i \rangle = \int_a^b f(x) b_i(x) dx$$

Integral Equations: Numerical Methods

Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$



Then, the resulting linear system $(I - M)\gamma = F$ is given by

$$m_{ij} = \langle a_j, b_i \rangle = \int_a^b a_j(x) b_i(x) dx = \int_a^b l_j^{(n)}(x) K(x_i, x) dx \approx \sum_{k=1}^n K(x_i, x_k) \int_a^b l_j^{(n)}(x) l_k^{(n)}(x) dx = \sum_{k=1}^n K(x_i, x_k) w_{jk},$$

where

$$W = \frac{h}{6} \begin{bmatrix} 2 & 1 & & & \\ 1 & 4 & 1 & & \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ & & 1 & 4 & 1 \\ & & & 1 & 2 \end{bmatrix}.$$

Similarly,


$$F_i = \langle f, b_i \rangle = \int_a^b f(x) b_i(x) dx \approx \sum_{j,k=1}^n b_i(x_j) f(x_k) \int_a^b l_j^{(n)}(x) l_k^{(n)}(x) dx$$

Integral Equations: Numerical Methods

Example

Equidistant piecewise linear interpolation:

$$l_j^{(n)}(x) = \begin{cases} (x - x_{j-1})/h, & x \in [x_{j-1}, x_j], j \geq 1, \\ (x_{j+1} - x)/h, & x \in [x_j, x_{j+1}], j \leq n-1, \\ 0, & \text{otherwise.} \end{cases}$$



Then, the resulting linear system $(I - M)\gamma = F$ is given by

$$m_{ij} = \langle a_j, b_i \rangle = \int_a^b a_j(x) b_i(x) dx = \int_a^b l_j^{(n)}(x) K(x_i, x) dx \approx \sum_{k=1}^n K(x_i, x_k) \int_a^b l_j^{(n)}(x) l_k^{(n)}(x) dx = \sum_{k=1}^n K(x_i, x_k) w_{jk},$$

where

$$W = \frac{h}{6} \begin{bmatrix} 2 & 1 & & & \\ 1 & 4 & 1 & & \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ & & 1 & 4 & 1 \\ & & & 1 & 2 \end{bmatrix}.$$

Similarly,

$$F_i = \langle f, b_i \rangle = \int_a^b f(x) b_i(x) dx \approx \sum_{j,k=1}^n b_i(x_j) f(x_k) \int_a^b l_j^{(n)}(x) l_k^{(n)}(x) dx = \sum_{j,k=1}^n b_i(x_j) f(x_k) w_{jk}.$$