

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE

- *Finite Difference Method*
- **Finite Element Method**



Numerical Methods for PDE: 2nd Order Elliptic PDE



Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Numerical Methods for PDE: 2nd Order Elliptic PDE



Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f \, v$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f \, v$$

Conversely, if u satisfies the second equation for all integrable functions v , then u satisfies the Poisson's equation. In fact, it is sufficient that the equation is satisfied for all C^∞ functions with compact support inside Ω .

Numerical Methods for PDE: 2nd Order Elliptic PDE

Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f v$$

Conversely, if u satisfies the second equation for all integrable functions v , then u satisfies the Poisson's equation. In fact, it is sufficient that the equation is satisfied for all C^∞ functions with compact support inside Ω .

In particular, if v is a C^1 function on Ω which vanishes on Γ , we have

$$-\int_{\Omega} \nabla u \cdot \nabla v = \int_{\Omega} f v.$$

This is the weak formulation of the boundary value problem.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f \, v$$

Conversely, if u satisfies the second equation for all integrable functions v , then u satisfies the Poisson's equation. In fact, it is sufficient that the equation is satisfied for all C^∞ functions with compact support inside Ω .

In particular, if v is a C^1 function on Ω which vanishes on Γ , we have

$$-\int_{\Omega} \nabla u \cdot \nabla v = \int_{\Omega} f \, v.$$

This is the weak formulation of the boundary value problem.

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space.



Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w)$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w)$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

that is

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\|$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\| \leq (1 + C/\gamma) \|u - w\| \text{ for all } w \in V_h.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\| \leq (1 + C/\gamma) \|u - w\| \text{ for all } w \in V_h. \text{ Therefore,}$$

$$\|u_h - u\| \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\| \leq (1 + C/\gamma) \|u - w\| \text{ for all } w \in V_h. \text{ Therefore,}$$

$$\|u_h - u\| \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|.$$

We, therefore, see that the Galerkin approximation error is bounded by a constant multiple of the best approximation error for u by functions in V_h !

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE

- *Finite Difference Method*
- **Finite Element Method**
- **Construction of FEM Approximation Spaces**



Numerical Methods for PDE: 2nd Order Elliptic PDE



Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Numerical Methods for PDE: 2nd Order Elliptic PDE



Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.


Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.





Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.





Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.





Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.





Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.



Numerical Methods for PDE: 2nd Order Elliptic PDE



Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure.



Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition



$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) =$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition



$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$



How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$

The functions φ_{mn} form a basis for subspace of piecewise linear functions with respect to the given partition/triangulation.



Numerical Methods for PDE: 2nd Order Elliptic PDE


Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$

The functions φ_{mn} form a basis for subspace of piecewise linear functions with respect to the given partition/triangulation.



Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$


Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$

The functions φ_{mn} form a basis for subspace of piecewise linear functions with respect to the given partition/triangulation.



Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

We see that matrix on the left hand side of the linear matrix (called stiffness matrix) for the piecewise linear finite elements for the Laplace operator on the unit square using a uniform mesh is exactly the matrix of the 5-point Laplacian.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

We see that matrix on the left hand side of the linear matrix (called stiffness matrix) for the piecewise linear finite elements for the Laplace operator on the unit square using a uniform mesh is exactly the matrix of the 5-point Laplacian.

Note that (*exercise*)

$$\int_{\Omega} \varphi_{mn} = h^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

We see that matrix on the left hand side of the linear matrix (called stiffness matrix) for the piecewise linear finite elements for the Laplace operator on the unit square using a uniform mesh is exactly the matrix of the 5-point Laplacian.

Note that (*exercise*)

$$\int_{\Omega} \varphi_{mn} = h^2$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N,$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Therefore, at vertices, we see that the finite element method converges with order 2 (*why?*).

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N,$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Therefore, at vertices, we see that the finite element method converges with order 2 (*why?*).

Recall that, from the general error analysis, we have

$$\|u_h - u\|_{H^1} \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|_{H^1}.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N,$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Therefore, at vertices, we see that the finite element method converges with order 2 (*why?*).

Recall that, from the general error analysis, we have

$$\|u_h - u\|_{H^1} \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|_{H^1}.$$

Exercise: Find C and γ for the piecewise linear finite elements.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Recall that, from the general error analysis, we have

$$\|u_h - u\|_{H^1} \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|_{H^1}.$$

Exercise: Find C and γ for the piecewise linear finite elements.

From the approximation theory, we have the following result on the best approximation error.

Theorem

Let there be given a family of triangulations $\{\mathcal{T}_h\}$ of a polygonal domain Ω and let $h = \max_{T \in \mathcal{T}_h} \text{diam}(T)$. Let r be a positive integer. For each h let $P_h: C(\Omega) \rightarrow M_0^r(\mathcal{T}_h)$ denote the nodal interpolant, where $M_0^r(\mathcal{T}_h)$ is the space of continuous functions which restrict to polynomials of degree at most r when restricted to any triangle $T \in \mathcal{T}_h$. Then, there is a constant c such that

$$\begin{aligned} \|u - P_h u\|_{L^\infty(\Omega)} &\leq ch^{r+1} \|u^{(r+1)}\|_{L^\infty(\Omega)}, & u \in C^{r+1}(\overline{\Omega}), \\ \|u - P_h u\|_{L^2(\Omega)} &\leq ch^{r+1} \|u^{(r+1)}\|_{L^2(\Omega)}, & u \in H^{r+1}(\Omega). \end{aligned}$$

Moreover, if the family of triangulations are shape regular (the minimal angle of each triangulation is bounded below uniformly), then there is a constant C such that

$$\begin{aligned} \|\nabla(u - P_h u)\|_{L^\infty(\Omega)} &\leq Ch^r \|u^{(r+1)}\|_{L^\infty(\Omega)}, & u \in C^{r+1}(\overline{\Omega}), \\ \|\nabla(u - P_h u)\|_{L^2(\Omega)} &\leq Ch^r \|u^{(r+1)}\|_{L^2(\Omega)}, & u \in H^{r+1}(\Omega). \end{aligned}$$