

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Well-posedness



*In many practical problems involving ODEs, instead of initial values, specifying additional at more than one points may be more relevant. In such case, we say that the problem is a **Boundary Value Problem (BVP)** for ODE.*

For example, if you want to throw a projectile from location A and want it to hit location B, you would need to solve the equation of motion together with conditions at A and B.

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.1 Well-posedness



Boundary Value Problems: Well-posedness

*In many practical problems involving ODEs, instead of initial values, specifying additional at more than one points may be more relevant. In such case, we say that the problem is a **Boundary Value Problem (BVP)** for ODE.*

For example, if you want to throw a projectile from location A and want it to hit location B, you would need to solve the equation of motion together with conditions at A and B.

A general first-order two-point BVP for an ODE has the form

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0$$

where $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ and $g: \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$.

Boundary Value Problems: Well-posedness

In many practical problems involving ODEs, instead of initial values, specifying additional at more than one points may be more relevant. In such case, we say that the problem is a **Boundary Value Problem (BVP)** for ODE.

For example, if you want to throw a projectile from location A and want it to hit location B, you would need to solve the equation of motion together with conditions at A and B.

A general first-order two-point BVP for an ODE has the form

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0$$

where $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ and $g: \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$.

The boundary condition is said to be **separated** if any given component of g involves solution values only at a or b , but not both.

Boundary Value Problems: Well-posedness

In many practical problems involving ODEs, instead of initial values, specifying additional at more than one points may be more relevant. In such case, we say that the problem is a **Boundary Value Problem (BVP)** for ODE.

For example, if you want to throw a projectile from location A and want it to hit location B, you would need to solve the equation of motion together with conditions at A and B.

A general first-order two-point BVP for an ODE has the form

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0$$

where $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ and $g: \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$.

The boundary condition is said to be **separated** if any given component of g involves solution values only at a or b , but not both.

The boundary condition is said to be **linear** if they have the form

$$B_a y(a) + B_b y(b) = c,$$

where $B_a, B_b \in \mathbb{R}^{n \times n}$ and $c \in \mathbb{R}^n$.

Boundary Value Problems: Well-posedness

In many practical problems involving ODEs, instead of initial values, specifying additional at more than one points may be more relevant. In such case, we say that the problem is a **Boundary Value Problem (BVP)** for ODE.

For example, if you want to throw a projectile from location A and want it to hit location B, you would need to solve the equation of motion together with conditions at A and B.

A general first-order two-point BVP for an ODE has the form

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0$$

where $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ and $g: \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$.

The boundary condition is said to be **separated** if any given component of g involves solution values only at a or b , but not both.

The boundary condition is said to be **linear** if they have the form

$$B_a y(a) + B_b y(b) = c,$$

where $B_a, B_b \in \mathbb{R}^{n \times n}$ and $c \in \mathbb{R}^n$.

If the boundary conditions are **both separated and linear**, then for each i , $1 \leq i \leq n$, either the i th row B_a or the i th row of B_b contains only zero entries.

Boundary Value Problems: Well-posedness

In many practical problems involving ODEs, instead of initial values, specifying additional at more than one points may be more relevant. In such case, we say that the problem is a **Boundary Value Problem (BVP)** for ODE.

For example, if you want to throw a projectile from location A and want it to hit location B, you would need to solve the equation of motion together with conditions at A and B.

A general first-order two-point BVP for an ODE has the form

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0$$

where $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ and $g: \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$.

The boundary condition is said to be **separated** if any given component of g involves solution values only at a or b , but not both.

The boundary condition is said to be **linear** if they have the form

$$B_a y(a) + B_b y(b) = c,$$

where $B_a, B_b \in \mathbb{R}^{n \times n}$ and $c \in \mathbb{R}^n$.

If the boundary conditions are **both separated and linear**, then for each i , $1 \leq i \leq n$, either the i th row B_a or the i th row of B_b contains only zero entries.

The BVP is said to be **linear** if both the ODE and the boundary conditions are linear.



Boundary Value Problems: Well-posedness

Example

Consider the two-point BVP for the second-order scalar ODE

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Boundary Value Problems: Well-posedness

Example

Consider the two-point BVP for the second-order scalar ODE

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

This problem is equivalent to the first-order system of ODEs

$$\begin{bmatrix} y_1' \\ y_2' \end{bmatrix} = \begin{bmatrix} y_2 \\ f(t, y_1, y_2) \end{bmatrix}, \quad a < t < b,$$

with separated linear boundary conditions

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} y_1(a) \\ y_2(a) \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} y_1(b) \\ y_2(b) \end{bmatrix} = \begin{bmatrix} \alpha \\ \beta \end{bmatrix}.$$

Boundary Value Problems: Well-posedness



Existence, Uniqueness and Conditioning

Unlike the IVPs, with the BVP, there is no single point at which complete state information is given, and hence no point at which local existence of a solution can be established.

Boundary Value Problems: Well-posedness



Existence, Uniqueness and Conditioning

Unlike the IVPs, with the BVP, there is no single point at which complete state information is given, and hence no point at which local existence of a solution can be established.

Similarly, there is no point with sufficient data to initiate the time stepping procedure for obtaining a numerical solution.

Boundary Value Problems: Well-posedness



Existence, Uniqueness and Conditioning

Unlike the IVPs, with the BVP, there is no single point at which complete state information is given, and hence no point at which local existence of a solution can be established.

Similarly, there is no point with sufficient data to initiate the time stepping procedure for obtaining a numerical solution.

It is, therefore, much more difficult to establish existence and uniqueness of solution for BVPs than for IVPs.

Boundary Value Problems: Well-posedness



Existence, Uniqueness and Conditioning

Unlike the IVPs, with the BVP, there is no single point at which complete state information is given, and hence no point at which local existence of a solution can be established.

Similarly, there is no point with sufficient data to initiate the time stepping procedure for obtaining a numerical solution.

It is, therefore, much more difficult to establish existence and uniqueness of solution for BVPs than for IVPs.

In full generality, the existence and uniqueness of solution does not hold.

Boundary Value Problems: Well-posedness

Existence, Uniqueness and Conditioning

Unlike the IVPs, with the BVP, there is no single point at which complete state information is given, and hence no point at which local existence of a solution can be established.

Similarly, there is no point with sufficient data to initiate the time stepping procedure for obtaining a numerical solution.

It is, therefore, much more difficult to establish existence and uniqueness of solution for BVPs than for IVPs.

In full generality, the existence and uniqueness of solution does not hold.

Example

Consider the two-point BVP

$$u'' = -u, \quad 0 < t < b,$$

with boundary conditions

$$u(0) = 0, \quad u(b) = \beta.$$

Boundary Value Problems: Well-posedness

Existence, Uniqueness and Conditioning

Unlike the IVPs, with the BVP, there is no single point at which complete state information is given, and hence no point at which local existence of a solution can be established.

Similarly, there is no point with sufficient data to initiate the time stepping procedure for obtaining a numerical solution.

It is, therefore, much more difficult to establish existence and uniqueness of solution for BVPs than for IVPs.

In full generality, the existence and uniqueness of solution does not hold.

Example

Consider the two-point BVP

$$u'' = -u, \quad 0 < t < b,$$

with boundary conditions

$$u(0) = 0, \quad u(b) = \beta.$$

The general solution of the ODE satisfying $u(0) = 0$ is $u(t) = c \sin t$ for a constant c . If $b = m\pi, m \in \mathbb{Z}$, then $c \sin b = 0$ for any c , so there are infinitely many solutions of the BVP if $\beta = 0$, but there is no solution $\beta \neq 0$.

Boundary Value Problems: Well-posedness



Consider the general two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

Boundary Value Problems: Well-posedness



Consider the general two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

Let $y(t; x)$ denote the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$. This solution is a solution to the BVP if $g(x, y(b; x)) = 0$.

Boundary Value Problems: Well-posedness

Consider the general two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

Let $y(t; x)$ denote the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$. This solution is a solution to the BVP if $g(x, y(b; x)) = 0$.

The solvability of the BVP therefore depends on the existence and uniqueness of solutions to the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$. We have seen (in the first course) that this is not always true, and therefore, can not expect a general theorem for existence and uniqueness of solutions for BVP.

Boundary Value Problems: Well-posedness

Consider the general two-point BVP

$$y' = f(t, y), \quad a < t < b,$$

with boundary conditions

$$g(y(a), y(b)) = 0.$$

Let $y(t; x)$ denote the solution to the IVP $y' = f(t, y)$, $y(a) = x$, $x \in \mathbb{R}^n$. This solution is a solution to the BVP if $g(x, y(b; x)) = 0$.

The solvability of the BVP therefore depends on the existence and uniqueness of solutions to the system of nonlinear algebraic equations $h(x) = 0$, where $h(x) = g(x, y(b; x))$. We have seen (in the first course) that this is not always true, and therefore, can not expect a general theorem for existence and uniqueness of solutions for BVP.

Such results are available only in certain specialized and simplified conditions.