

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.2 Stability

2.3 Euler's method

2.4 Implicit method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Implicit Methods



Euler's method is an *explicit method* in that it uses only information at time t_k to advance to solution to time t_{k+1} .

Initial Value Problems: Implicit Methods



Euler's method is an **explicit method** in that it uses only information at time t_k to advance to solution to time t_{k+1} .

This is convenient but comes at the cost of having a rather limited region of stability.

Initial Value Problems: Implicit Methods



Euler's method is an **explicit method** in that it uses only information at time t_k to advance to solution to time t_{k+1} .

This is convenient but comes at the cost of having a rather limited region of stability.

A larger stability region can be obtained by using information at t_{k+1} !

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.2 Stability

2.3 Euler's method

2.4 Implicit method

- Backward Euler method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Implicit Methods

Euler's method is an **explicit method** in that it uses only information at time t_k to advance to solution to time t_{k+1} .

This is convenient but comes at the cost of having a rather limited region of stability.

A larger stability region can be obtained by using information at t_{k+1} !

For the IVP $y' = f(t, y)$, $y(t_0) = y_0$, we have

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$

Initial Value Problems: Implicit Methods

Euler's method is an **explicit method** in that it uses only information at time t_k to advance to solution to time t_{k+1} .

This is convenient but comes at the cost of having a rather limited region of stability.


A larger stability region can be obtained by using information at t_{k+1} !

For the IVP $y' = f(t, y)$, $y(t_0) = y_0$, we have

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$

Using the right end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(b)$$



Initial Value Problems: Implicit Methods

Euler's method is an **explicit method** in that it uses only information at time t_k to advance to solution to time t_{k+1} .

This is convenient but comes at the cost of having a rather limited region of stability.

A larger stability region can be obtained by using information at t_{k+1} !

For the IVP $y' = f(t, y)$, $y(t_0) = y_0$, we have


$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$

Using the right end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(b)$$

we get the method

$$y_{k+1} = y_k + h_k f(t_{k+1}, y_{k+1})$$



Initial Value Problems: Implicit Methods

Euler's method is an **explicit method** in that it uses only information at time t_k to advance to solution to time t_{k+1} .

This is convenient but comes at the cost of having a rather limited region of stability.

A larger stability region can be obtained by using information at t_{k+1} !

For the IVP $y' = f(t, y)$, $y(t_0) = y_0$, we have

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


Using the right end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(b)$$

we get the method

$$y_{k+1} = y_k + h_k f(t_{k+1}, y_{k+1})$$

Background
Euler method



Initial Value Problems: Implicit Methods

Euler's method is an **explicit method** in that it uses only information at time t_k to advance to solution to time t_{k+1} .

This is convenient but comes at the cost of having a rather limited region of stability.

A larger stability region can be obtained by using information at t_{k+1} !

For the IVP $y' = f(t, y)$, $y(t_0) = y_0$, we have

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


Using the right end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(b)$$

we get the method

$$y_{k+1} = y_k + h_k f(t_{k+1}, y_{k+1})$$

Background
Euler method



When y_{k+1} appears on the right-hand side of the method

$$y_{k+1} = \Phi(f, y_{k+1}, y_k, h_k)$$

then the method is called an **implicit method**.

Initial Value Problems: Implicit Methods



Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

Initial Value Problems: Implicit Methods



Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1)$$



Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$



Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$

This is a nonlinear equation that can be solved using some of the techniques you learnt in the first course.



Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$

This is a nonlinear equation that can be solved using some of the techniques you learnt in the first course. For example, we could use Newton's method to iteratively solve for y_1 :

Initial Value Problems: Implicit Methods

Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$

This is a nonlinear equation that can be solved using some of the techniques you learnt in the first course. For example, we could use Newton's method to iteratively solve for y_1 :

$$y_1^{(n+1)} = y_1^{(n)} - \frac{0.5(y_1^{(n)})^3 + y_1^{(n)} - 1}{1.5(y_1^{(n)})^2 + 1},$$

Initial Value Problems: Implicit Methods

Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$

This is a nonlinear equation that can be solved using some of the techniques you learnt in the first course. For example, we could use Newton's method to iteratively solve for y_1 :

$$y_1^{(n+1)} = y_1^{(n)} - \frac{0.5(y_1^{(n)})^3 + y_1^{(n)} - 1}{1.5(y_1^{(n)})^2 + 1}, \quad y_1^{(0)} = ?$$

Initial Value Problems: Implicit Methods

Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$

This is a nonlinear equation that can be solved using some of the techniques you learnt in the first course. For example, we could use Newton's method to iteratively solve for y_1 :

$$y_1^{(n+1)} = y_1^{(n)} - \frac{0.5(y_1^{(n)})^3 + y_1^{(n)} - 1}{1.5(y_1^{(n)})^2 + 1}, \quad y_1^{(0)} = y_0$$

Initial Value Problems: Implicit Methods


Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$

This is a nonlinear equation that can be solved using some of the techniques you learnt in the first course. For example, we could use Newton's method to iteratively solve for y_1 :

$$y_1^{(n+1)} = y_1^{(n)} - \frac{0.5(y_1^{(n)})^3 + y_1^{(n)} - 1}{1.5(y_1^{(n)})^2 + 1}, \quad y_1^{(0)} = y_0$$



Initial Value Problems: Implicit Methods





Example

Let us solve $y' = -y^3$, $y(0) = 1$ using the backward Euler method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = y_0 + hf(t_1, y_1) = 1 - 0.5y_1^3$$

This is a nonlinear equation that can be solved using some of the techniques you learnt in the first course. For example, we could use Newton's method to iteratively solve for y_1 :

$$y_1^{(n+1)} = y_1^{(n)} - \frac{0.5(y_1^{(n)})^3 + y_1^{(n)} - 1}{1.5(y_1^{(n)})^2 + 1}, \quad y_1^{(0)} = y_0 - 0.5y_0^3 = 0.5$$



Initial Value Problems: Implicit Methods



Why can we always solve the resulting non-linear system $y_{k+1} = y_k + hf(t_{k+1}, y_{k+1})$?

Initial Value Problems: Implicit Methods



Why can we always solve the resulting non-linear system $y_{k+1} = y_k + hf(t_{k+1}, y_{k+1})$?

Define

$$F(z) = y_k + hf(t_{k+1}, z)$$

Initial Value Problems: Implicit Methods



Why can we always solve the resulting non-linear system $y_{k+1} = y_k + hf(t_{k+1}, y_{k+1})$?

Define

$$F(z) = y_k + hf(t_{k+1}, z)$$

then

$$|F(z_2) - F(z_1)| = |(y_k + hf(t_{k+1}, z_2)) - (y_k + hf(t_{k+1}, z_1))|$$

Initial Value Problems: Implicit Methods



Why can we always solve the resulting non-linear system $y_{k+1} = y_k + hf(t_{k+1}, y_{k+1})$?

Define

$$F(z) = y_k + hf(t_{k+1}, z)$$

then

$$|F(z_2) - F(z_1)| = |(y_k + hf(t_{k+1}, z_2)) - (y_k + hf(t_{k+1}, z_1))|$$

$$|F(z_2) - F(z_1)| \leq h|f(t_{k+1}, z_2) - f(t_{k+1}, z_1)| \leq hL|z_2 - z_1|$$

Initial Value Problems: Implicit Methods



Why can we always solve the resulting non-linear system $y_{k+1} = y_k + hf(t_{k+1}, y_{k+1})$?

Define

$$F(z) = y_k + hf(t_{k+1}, z)$$

then

$$|F(z_2) - F(z_1)| = |(y_k + hf(t_{k+1}, z_2)) - (y_k + hf(t_{k+1}, z_1))|$$

$$|F(z_2) - F(z_1)| \leq h|f(t_{k+1}, z_2) - f(t_{k+1}, z_1)| \leq hL|z_2 - z_1|$$

Therefore, if $h < 1/L$, then F is a contraction. Thus, it has a fixed point, say y_{k+1} .

Initial Value Problems: Implicit Methods



Why can we always solve the resulting non-linear system $y_{k+1} = y_k + hf(t_{k+1}, y_{k+1})$?

Define

$$F(z) = y_k + hf(t_{k+1}, z)$$

then

$$|F(z_2) - F(z_1)| = |(y_k + hf(t_{k+1}, z_2)) - (y_k + hf(t_{k+1}, z_1))|$$

$$|F(z_2) - F(z_1)| \leq h|f(t_{k+1}, z_2) - f(t_{k+1}, z_1)| \leq hL|z_2 - z_1|$$

Therefore, if $h < 1/L$, then F is a contraction. Thus, it has a fixed point, say y_{k+1} . That is,

$$y_{k+1} = F(y_{k+1}) = y_k + hf(t_{k+1}, y_{k+1}).$$

Initial Value Problems: Implicit Methods

Why can we always solve the resulting non-linear system $y_{k+1} = y_k + hf(t_{k+1}, y_{k+1})$?

Define

$$F(z) = y_k + hf(t_{k+1}, z)$$

then

$$|F(z_2) - F(z_1)| = |(y_k + hf(t_{k+1}, z_2)) - (y_k + hf(t_{k+1}, z_1))|$$

$$|F(z_2) - F(z_1)| \leq h|f(t_{k+1}, z_2) - f(t_{k+1}, z_1)| \leq hL|z_2 - z_1|$$

Therefore, if $h < 1/L$, then F is a contraction. Thus, it has a fixed point, say y_{k+1} . That is,

$$y_{k+1} = F(y_{k+1}) = y_k + hf(t_{k+1}, y_{k+1}).$$

Remark:

In the previous example, we could have used the fixed point iteration $y_{k+1}^{(n+1)} = y_k + hf(t_{k+1}, y_{k+1}^{(n)})$ in place of Newton's iterations

$$y_{k+1}^{(n+1)} = y_{k+1}^{(n)} - \frac{0.5(y_{k+1}^{(n)})^3 + y_{k+1}^{(n)} - 1}{1.5(y_{k+1}^{(n)})^2 + 1}.$$

Initial Value Problems: Implicit Methods



To understand the stability, let us again consider the ODE $y' = \lambda y$.

Initial Value Problems: Implicit Methods



To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

Initial Value Problems: Implicit Methods



To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

Initial Value Problems: Implicit Methods



To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1 - h\lambda} \right) y_k$$

Initial Value Problems: Implicit Methods



To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$

$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$

Initial Value Problems: Implicit Methods



To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$

$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$

For stability of the backward Euler method, we need

$$\left|\frac{1}{1-h\lambda}\right| \leq 1 \quad \text{or} \quad |1 - \lambda h| \geq 1.$$

Initial Value Problems: Implicit Methods

To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$

$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$

For stability of the backward Euler method, we need

$$\left|\frac{1}{1-h\lambda}\right| \leq 1 \quad \text{or} \quad |1 - \lambda h| \geq 1.$$

Note that this condition holds for all $h > 0$ where $\text{Re}(\lambda) < 0$.

Initial Value Problems: Implicit Methods



To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$


$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$

$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$

For stability of the backward Euler method, we need

$$\left|\frac{1}{1-h\lambda}\right| \leq 1 \quad \text{or} \quad |1 - \lambda h| \geq 1.$$

Note that this condition holds for all $h > 0$ where $\text{Re}(\lambda) < 0$.



Initial Value Problems: Implicit Methods

To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$


$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$

For stability of the backward Euler method, we need

$$\left|\frac{1}{1-h\lambda}\right| \leq 1 \quad \text{or} \quad |1 - \lambda h| \geq 1.$$

Note that this condition holds for all $h > 0$ where $\text{Re}(\lambda) < 0$.

Compare with
Euler's method



Initial Value Problems: Implicit Methods

To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$


$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$

For stability of the backward Euler method, we need

$$\left|\frac{1}{1-h\lambda}\right| \leq 1 \quad \text{or} \quad |1 - \lambda h| \geq 1.$$

Note that this condition holds for all $h > 0$ where $\text{Re}(\lambda) < 0$.

Unconditionally
stable!



Compare with
Euler's method

Initial Value Problems: Implicit Methods

To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method

$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$

$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$


For stability of the backward Euler method, we need

$$\left|\frac{1}{1-h\lambda}\right| \leq 1 \quad \text{or} \quad |1 - \lambda h| \geq 1.$$


Note that this condition holds for all $h > 0$ where $\text{Re}(\lambda) < 0$.

Unconditionally
stable!

...
means that the
method is stable in
the entire left half
of the complex
plane.



Compare with
Euler's method



Initial Value Problems: Implicit Methods

To understand the stability, let us again consider the ODE $y' = \lambda y$. Applying backward Euler method


$$y_{k+1} = y_k + h(\lambda y_{k+1})$$

$$(1 - h\lambda)y_{k+1} = y_k$$

$$y_{k+1} = \left(\frac{1}{1-h\lambda}\right) y_k$$

$$y_k = \left(\frac{1}{1-h\lambda}\right)^k y_0$$

Unconditionally
stable!



For stability of the backward Euler method, we need

$$\left|\frac{1}{1-h\lambda}\right| \leq 1 \quad \text{or} \quad |1 - \lambda h| \geq 1.$$

Note that this condition holds for all $h > 0$ where $\text{Re}(\lambda) < 0$.

For the general ODE $y' = f(t, y)$, we have


Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$e_{k+1} = y_{k+1} - y(t_{k+1})$$

Unconditionally
stable!




Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned} e_{k+1} &= y_{k+1} - y(t_{k+1}) \\ &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \end{aligned}$$

Unconditionally
stable!




Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned} e_{k+1} &= y_{k+1} - y(t_{k+1}) \\ &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\ &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_k, y(t_k)) \\ &\quad + [y(t_k) + h_k f(t_k, y(t_k)) - y(t_{k+1})] \end{aligned}$$

Unconditionally
stable!




Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned} e_{k+1} &= y_{k+1} - y(t_{k+1}) \\ &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\ &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\ &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\ &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \end{aligned}$$

Unconditionally
stable!




Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!




Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1}$$


Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have


$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

Initial Value Problems: Implicit Methods

For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.


Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.


Is backward Euler method more accurate?

Initial Value Problems: Implicit Methods

For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.

Is backward Euler method more accurate?


$$\ell_{k+1} = y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})$$

Initial Value Problems: Implicit Methods

For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.

Is backward Euler method more accurate?


$$\ell_{k+1} = y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1}) = h_k f(t_{k+1}, y(t_{k+1})) + [y(t_k) - y(t_{k+1})]$$

Initial Value Problems: Implicit Methods

For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.


Is backward Euler method more accurate?

$$\begin{aligned}
 \ell_{k+1} &= y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1}) = h_k f(t_{k+1}, y(t_{k+1})) + [y(t_k) - y(t_{k+1})] \\
 &= h_k f(t_{k+1}, y(t_{k+1})) + [(t_k - t_{k+1})y'(t_{k+1}) + O((t_k - t_{k+1})^2)]
 \end{aligned}$$

For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.

Is backward Euler method more accurate?

$$\begin{aligned}
 \ell_{k+1} &= y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1}) = h_k f(t_{k+1}, y(t_{k+1})) + [y(t_k) - y(t_{k+1})] \\
 &= h_k f(t_{k+1}, y(t_{k+1})) + [(t_k - t_{k+1})y'(t_{k+1}) + O((t_k - t_{k+1})^2)] \\
 &= h_k f(t_{k+1}, y(t_{k+1})) + [-h_k f(t_{k+1}, y(t_{k+1})) + O(h_k^2)]
 \end{aligned}$$


Initial Value Problems: Implicit Methods



For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.

Is backward Euler method more accurate?


$$\begin{aligned}
 \ell_{k+1} &= y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1}) = h_k f(t_{k+1}, y(t_{k+1})) + [y(t_k) - y(t_{k+1})] \\
 &= h_k f(t_{k+1}, y(t_{k+1})) + [(t_k - t_{k+1})y'(t_{k+1}) + O((t_k - t_{k+1})^2)] \\
 &= h_k f(t_{k+1}, y(t_{k+1})) + [-h_k f(t_{k+1}, y(t_{k+1})) + O(h_k^2)] = O(h_k^2)
 \end{aligned}$$

Initial Value Problems: Implicit Methods

For the general ODE $y' = f(t, y)$, we have

$$\begin{aligned}
 e_{k+1} &= y_{k+1} - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_{k+1}) \\
 &= y_k + h_k f(t_{k+1}, y_{k+1}) - y(t_k) - h_k f(t_{k+1}, y(t_{k+1})) \\
 &\quad + [y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1})] \\
 &= e_k + h_k [f(t_{k+1}, y_{k+1}) - f(t_{k+1}, y(t_{k+1}))] + \ell_{k+1} \\
 &= e_k + h_k [f'(t_{k+1}, \alpha y_{k+1} + (1 - \alpha)y(t_{k+1}))] e_{k+1} + \ell_{k+1}
 \end{aligned}$$

Unconditionally
stable!



We, therefore, have

$$(I - h_k f') e_{k+1} = e_k + \ell_{k+1} \quad \text{or} \quad e_{k+1} = (I - h_k f')^{-1} e_k + (I - h_k f')^{-1} \ell_{k+1}$$

For stability, we need $\rho(I - h_k f')^{-1} \leq 1$, i.e., eigenvalues of $h_k f'$ must lie outside the unit circle centered at 1.

Is backward Euler method more accurate?

$$\begin{aligned}
 \ell_{k+1} &= y(t_k) + h_k f(t_{k+1}, y(t_{k+1})) - y(t_{k+1}) = h_k f(t_{k+1}, y(t_{k+1})) + [y(t_k) - y(t_{k+1})] \\
 &= h_k f(t_{k+1}, y(t_{k+1})) + [(t_k - t_{k+1})y'(t_{k+1}) + O((t_k - t_{k+1})^2)] \\
 &= h_k f(t_{k+1}, y(t_{k+1})) + [-h_k f(t_{k+1}, y(t_{k+1})) + O(h_k^2)] = O(h_k^2)
 \end{aligned}$$

first-order
accurate!