

# *Numerical Analysis & Scientific Computing II*

## *Module 2*

# *Initial Value Problems*

*2.4 Implicit method*

*2.5 Stiffness*

**2.6 Linear Multistep Methods**

**- Consistency and Order**



*Akash Anand*  
MATH, IIT KANPUR

# Initial Value Problems: Linear Multistep Methods

## Consistency and Order

For the linear multistep method

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j})$$

define the local error as

$$\ell_{n+1}(y, h) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh)$$

for any  $y \in C^1$ , and  $h > 0$ .

# Initial Value Problems: Linear Multistep Methods

## Consistency and Order

For the linear multistep method

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j})$$

define the local error as

$$\ell_{n+1}(y, h) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh)$$

for any  $y \in C^1$ , and  $h > 0$ .

The linear multistep method is **consistent** if

$$\lim_{h \rightarrow 0} \max_{k \leq n < N} \left| \frac{\ell_{n+1}(y, h)}{h} \right| = 0$$

for all  $y \in C^1$ .

# Initial Value Problems: Linear Multistep Methods

## Consistency and Order

For the linear multistep method

$$\sum_{j=-1}^k a_j y_{n-j} = h \sum_{j=-1}^k b_j f(t_{n-j}, y_{n-j})$$

define the local error as

$$\ell_{n+1}(y, h) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh)$$

for any  $y \in C^1$ , and  $h > 0$ .

The linear multistep method is **consistent** if

$$\lim_{h \rightarrow 0} \max_{k \leq n < N} \left| \frac{\ell_{n+1}(y, h)}{h} \right| = 0$$

for all  $y \in C^1$ .

The method has **order  $p$**  if for all  $y \in C^{p+1}$  there exists constants  $C, h_0 > 0$  such that

$$\max_{k \leq n < N} \left| \frac{\ell_{n+1}(y, h)}{h} \right| \leq Ch^p$$

whenever  $h < h_0$ .

# Initial Value Problems: Linear Multistep Methods



## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

# Initial Value Problems: Linear Multistep Methods

## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

## Theorem

*A linear multistep is consistent if and only if*

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

*The method is of order  $p$  if and only if*

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

# Initial Value Problems: Linear Multistep Methods

## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

## Theorem

*A linear multistep is consistent if and only if*

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

Consistency conditions

*The method is of order  $p$  if and only if*

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

Order conditions

# Initial Value Problems: Linear Multistep Methods

## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

## Theorem

*A linear multistep is consistent if and only if*

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

Consistency conditions

*The method is of order  $p$  if and only if*

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

Order conditions

## Proof.



# Initial Value Problems: Linear Multistep Methods

## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

## Theorem

*A linear multistep is consistent if and only if*

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

Consistency conditions

*The method is of order  $p$  if and only if*

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

Order conditions

## Proof.

*Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .*

# Initial Value Problems: Linear Multistep Methods

## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

## Theorem

*A linear multistep is consistent if and only if*

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k ja_j + \sum_{j=-1}^k b_j = 0.$$

Consistency conditions

*The method is of order  $p$  if and only if*

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

Order conditions

## Proof.

Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .

Note that  $\xi_j \rightarrow t_n$  as  $h \rightarrow 0$ .

# Initial Value Problems: Linear Multistep Methods

## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

## Theorem

*A linear multistep is consistent if and only if*

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k ja_j + \sum_{j=-1}^k b_j = 0.$$

Consistency conditions

*The method is of order  $p$  if and only if*

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

Order conditions

## Proof.

Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .

Note that  $\xi_j \rightarrow t_n$  as  $h \rightarrow 0$ . Now,

$$\ell_{n+1}(y, h) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh)$$

# Initial Value Problems: Linear Multistep Methods

## Note

*It is not true that every method of order  $p$  converges with order  $p$ . It may not even converge at all!*

## Theorem

*A linear multistep is consistent if and only if*

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k ja_j + \sum_{j=-1}^k b_j = 0.$$

Consistency conditions

*The method is of order  $p$  if and only if*

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

Order conditions

## Proof.

Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .

Note that  $\xi_j \rightarrow t_n$  as  $h \rightarrow 0$ . Now,

$$\ell_{n+1}(y, h) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j [y(t_n) - jhy'(\xi_j)]$$

# Initial Value Problems: Linear Multistep Methods

## Proof.

Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .

Note that  $\xi_j \rightarrow t_n$  as  $h \rightarrow 0$ . Now,

$$\begin{aligned}\ell_{n+1}(y, h) &= h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j [y(t_n) - jhy'(\xi_j)] \\ &= -y(t_n)C_0 + hy'(t_n)C_1 + R,\end{aligned}$$

where

$$C_0 = \sum_{j=-1}^k a_j, \quad C_1 = \sum_{j=-1}^k ja_j + \sum_{j=-1}^k b_j,$$

$$R = h \sum_{j=-1}^k b_j [y'(t_n - jh) - y'(t_n)] + h \sum_{j=-1}^k ja_j [y'(\xi_j) - y'(t_n)]$$

# Initial Value Problems: Linear Multistep Methods

## Proof.

Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .

Note that  $\xi_j \rightarrow t_n$  as  $h \rightarrow 0$ . Now,

$$\begin{aligned}\ell_{n+1}(y, h) &= h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j [y(t_n) - jhy'(\xi_j)] \\ &= -y(t_n)C_0 + hy'(t_n)C_1 + R,\end{aligned}$$

where

$$C_0 = \sum_{j=-1}^k a_j, \quad C_1 = \sum_{j=-1}^k ja_j + \sum_{j=-1}^k b_j,$$

$$R = h \sum_{j=-1}^k b_j [y'(t_n - jh) - y'(t_n)] + h \sum_{j=-1}^k ja_j [y'(\xi_j) - y'(t_n)]$$

By uniform continuity of  $y'$ , we see that  $R/h \rightarrow 0$  as  $h \rightarrow 0$ .

# Initial Value Problems: Linear Multistep Methods

## Proof.

Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .

Note that  $\xi_j \rightarrow t_n$  as  $h \rightarrow 0$ . Now,

$$\begin{aligned}\ell_{n+1}(y, h) &= h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j [y(t_n) - jhy'(\xi_j)] \\ &= -y(t_n)C_0 + hy'(t_n)C_1 + R,\end{aligned}$$

where

$$\begin{aligned}C_0 &= \sum_{j=-1}^k a_j, \quad C_1 = \sum_{j=-1}^k ja_j + \sum_{j=-1}^k b_j, \\ R &= h \sum_{j=-1}^k b_j [y'(t_n - jh) - y'(t_n)] + h \sum_{j=-1}^k ja_j [y'(\xi_j) - y'(t_n)]\end{aligned}$$

By uniform continuity of  $y'$ , we see that  $R/h \rightarrow 0$  as  $h \rightarrow 0$ . Therefore,  $\ell_{n+1}(y, h)/h \rightarrow 0$  if and only if ...

# Initial Value Problems: Linear Multistep Methods

## Proof.

Using Taylor's theorem, we have  $y(t_n - jh) = y(t_n) - jhy'(\xi_j)$  for some  $\xi_j \in (t_n - kh, t_n + h)$ ,  $j = -1, \dots, k$ .

Note that  $\xi_j \rightarrow t_n$  as  $h \rightarrow 0$ . Now,

$$\begin{aligned}\ell_{n+1}(y, h) &= h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j y(t_n - jh) = h \sum_{j=-1}^k b_j y'(t_n - jh) - \sum_{j=-1}^k a_j [y(t_n) - jhy'(\xi_j)] \\ &= -y(t_n)C_0 + hy'(t_n)C_1 + R,\end{aligned}$$

where

$$C_0 = \sum_{j=-1}^k a_j, \quad C_1 = \sum_{j=-1}^k ja_j + \sum_{j=-1}^k b_j,$$

$$R = h \sum_{j=-1}^k b_j [y'(t_n - jh) - y'(t_n)] + h \sum_{j=-1}^k ja_j [y'(\xi_j) - y'(t_n)]$$

By uniform continuity of  $y'$ , we see that  $R/h \rightarrow 0$  as  $h \rightarrow 0$ . Therefore,  $\ell_{n+1}(y, h)/h \rightarrow 0$  if and only if  $C_0 = 0$  and  $C_1 = 0$ , that is consistency conditions are satisfied.



**Proof.** ...

Similarly, if  $y \in C^{p+1}$  we have

$$y(t_n - jh) = \sum_{m=0}^p \frac{(-j)^m}{m!} h^m y^{(m)}(t_n) + \frac{(-j)^{p+1}}{(p+1)!} h^{p+1} y^{(p+1)}(\xi_j)$$
$$y'(t_n - jh) = \sum_{m=1}^p \frac{(-j)^{m-1}}{(m-1)!} h^{m-1} y^{(m)}(t_n) + \frac{(-j)^p}{p!} h^p y^{(p+1)}(\zeta_j)$$

for some  $\xi_j, \zeta_j \in (t_n - kh, t_n + h), j = -1, \dots, k$ .

# Initial Value Problems: Linear Multistep Methods

**Proof.** ...

Similarly, if  $y \in C^{p+1}$  we have

$$y(t_n - jh) = \sum_{m=0}^p \frac{(-j)^m}{m!} h^m y^{(m)}(t_n) + \frac{(-j)^{p+1}}{(p+1)!} h^{p+1} y^{(p+1)}(\xi_j)$$

$$y'(t_n - jh) = \sum_{m=1}^p \frac{(-j)^{m-1}}{(m-1)!} h^{m-1} y^{(m)}(t_n) + \frac{(-j)^p}{p!} h^p y^{(p+1)}(\zeta_j)$$

for some  $\xi_j, \zeta_j \in (t_n - kh, t_n + h), j = -1, \dots, k$ . This yields

$$\ell_{n+1}(y, h) = \sum_{m=0}^p h^m y^{(m)}(t_n) C_m + R$$

where

$$C_m = \frac{1}{m!} \left[ m \sum_{j=-1}^k (-j)^{m-1} b_j - \sum_{j=-1}^k (-j)^m a_j \right], \quad R = h^{p+1} \sum_{j=-1}^k \left[ b_j \frac{(-j)^p}{p!} y^{(p+1)}(\zeta_j) - a_j \frac{(-j)^{p+1}}{(p+1)!} y^{(p+1)}(\xi_j) \right].$$

# Initial Value Problems: Linear Multistep Methods

**Proof.** ...

Similarly, if  $y \in C^{p+1}$  we have

$$y(t_n - jh) = \sum_{m=0}^p \frac{(-j)^m}{m!} h^m y^{(m)}(t_n) + \frac{(-j)^{p+1}}{(p+1)!} h^{p+1} y^{(p+1)}(\xi_j)$$

$$y'(t_n - jh) = \sum_{m=1}^p \frac{(-j)^{m-1}}{(m-1)!} h^{m-1} y^{(m)}(t_n) + \frac{(-j)^p}{p!} h^p y^{(p+1)}(\zeta_j)$$

for some  $\xi_j, \zeta_j \in (t_n - kh, t_n + h), j = -1, \dots, k$ . This yields

$$\ell_{n+1}(y, h) = \sum_{m=0}^p h^m y^{(m)}(t_n) C_m + R$$

where

$$C_m = \frac{1}{m!} \left[ m \sum_{j=-1}^k (-j)^{m-1} b_j - \sum_{j=-1}^k (-j)^m a_j \right], \quad R = h^{p+1} \sum_{j=-1}^k \left[ b_j \frac{(-j)^p}{p!} y^{(p+1)}(\zeta_j) - a_j \frac{(-j)^{p+1}}{(p+1)!} y^{(p+1)}(\xi_j) \right].$$

Since  $R = O(h^{p+1})$ ,  $\ell_{n+1}(y, h)/h = O(h^p)$  if and only if all the  $C_m$  vanish.

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Remarks

1. This theorem is an example of how a complicated analytic condition may sometimes reduce to a simple algebraic criterion.

# Initial Value Problems: Linear Multistep Methods

## Theorem

A linear multistep is consistent if and only if

$$\sum_{j=-1}^k a_j = 0, \quad \sum_{j=-1}^k j a_j + \sum_{j=-1}^k b_j = 0.$$

The method is of order  $p$  if and only if

$$\sum_{j=-1}^k (-j)^m a_j - m \sum_{j=-1}^k (-j)^{m-1} b_j = 0, \quad m = 0, 1, \dots, p.$$

## Remarks

1. This theorem is an example of how a complicated analytic condition may sometimes reduce to a simple algebraic criterion.
2. Such algebraic criteria for multistep methods can be expressed in terms of characteristic polynomials of the method:

$$\rho(z) = \sum_{j=-1}^k a_j z^{k-j}, \quad \sigma(z) = \sum_{j=-1}^k b_j z^{k-j}.$$

For example, the consistency conditions are  $\rho(1) = 0$  and  $\rho'(1) = \sigma(1)$ .