

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.1 Well-posedness

- Linear two point BVP





Boundary Value Problems: Well-posedness

Theorem

Consider the linear two-point BVP

$$y' = A(t)y + r(t), \quad a < t < b,$$

where $A(t)$ and $b(t)$ are continuous, with boundary conditions

$$B_a y(a) + B_b y(b) = c.$$

The BVP has a unique solution if and only if the matrix

$$Q = B_a Y(a) + B_b Y(b)$$

is non-singular where Y is the fundamental solution matrix for the ODE whose i th column $y_i(t)$ is the solution to the homogeneous ODE $y' = A(t)y$ with initial condition $y(a) = e_i$, where e_i is the i th column of the identity matrix; these columns are called solution modes.

Boundary Value Problems: Well-posedness

Theorem

Consider the linear two-point BVP

$$y' = A(t)y + r(t), \quad a < t < b,$$

where $A(t)$ and $b(t)$ are continuous, with boundary conditions

$$B_a y(a) + B_b y(b) = c.$$

The BVP has a unique solution if and only if the matrix

$$Q = B_a Y(a) + B_b Y(b)$$

is non-singular where Y is the fundamental solution matrix for the ODE whose i th column $y_i(t)$ is the solution to the homogeneous ODE $y' = A(t)y$ with initial condition $y(a) = e_i$, where e_i is the i th column of the identity matrix; these columns are called solution modes.

Proof.

Assume that the matrix Q is invertible.

Boundary Value Problems: Well-posedness

Theorem

Consider the linear two-point BVP

$$y' = A(t)y + r(t), \quad a < t < b,$$

where $A(t)$ and $b(t)$ are continuous, with boundary conditions

$$B_a y(a) + B_b y(b) = c.$$

The BVP has a unique solution if and only if the matrix

$$Q = B_a Y(a) + B_b Y(b)$$

is non-singular where Y is the fundamental solution matrix for the ODE whose i th column $y_i(t)$ is the solution to the homogeneous ODE $y' = A(t)y$ with initial condition $y(a) = e_i$, where e_i is the i th column of the identity matrix; these columns are called solution modes.

Proof.

Assume that the matrix Q is invertible.

Uniqueness of solution follows from the fact that, if $y_1(t)$ and $y_2(t)$ are two solutions to the BVP, then $y(t) = y_1(t) - y_2(t)$ satisfies

$$y'(t) = A(t)y(t), \quad B_a y(a) + B_b y(b) = 0,$$

Boundary Value Problems: Well-posedness

Theorem

Consider the linear two-point BVP

$$y' = A(t)y + r(t), \quad a < t < b,$$

where $A(t)$ and $b(t)$ are continuous, with boundary conditions

$$B_a y(a) + B_b y(b) = c.$$

The BVP has a unique solution if and only if the matrix

$$Q = B_a Y(a) + B_b Y(b)$$

is non-singular where Y is the fundamental solution matrix for the ODE whose i th column $y_i(t)$ is the solution to the homogeneous ODE $y' = A(t)y$ with initial condition $y(a) = e_i$, where e_i is the i th column of the identity matrix; these columns are called solution modes.

Proof.

Assume that the matrix Q is invertible.

Uniqueness of solution follows from the fact that, if $y_1(t)$ and $y_2(t)$ are two solutions to the BVP, then $y(t) = y_1(t) - y_2(t)$ satisfies

$$y'(t) = A(t)y(t), \quad B_a y(a) + B_b y(b) = 0,$$

and hence $y(t)$ must have the form $y(t) = Y(t)d$ for some $d \in \mathbb{R}^n$ satisfying, $Qd = 0$.

Boundary Value Problems: Well-posedness

Proof.

Uniqueness of solution follows from the fact that, if $y_1(t)$ and $y_2(t)$ are two solutions to the BVP, then $y(t) = y_1(t) - y_2(t)$ satisfies

$$y'(t) = A(t)y(t), B_a y(a) + B_b y(b) = 0,$$

and hence $y(t)$ must have the form $y(t) = Y(t)d$ for some $d \in \mathbb{R}^n$ satisfying, $Qd = 0$.

Now, one can see that the unique solution to the BVP is given by

$$y(t) = Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(t) \int_a^t Y^{-1}(s)r(s)ds$$

by directly verifying that it satisfies the ODE and the boundary condition.

Boundary Value Problems: Well-posedness

Proof.

Uniqueness of solution follows from the fact that, if $y_1(t)$ and $y_2(t)$ are two solutions to the BVP, then $y(t) = y_1(t) - y_2(t)$ satisfies

$$y'(t) = A(t)y(t), \quad B_a y(a) + B_b y(b) = 0,$$

and hence $y(t)$ must have the form $y(t) = Y(t)d$ for some $d \in \mathbb{R}^n$ satisfying, $Qd = 0$.

Now, one can see that the unique solution to the BVP is given by

$$y(t) = Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(t) \int_a^t Y^{-1}(s)r(s)ds$$

by directly verifying that it satisfies the ODE and the boundary condition. Indeed,

$$y'(t) = A(t)Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + A(t)Y(t) \int_a^t Y^{-1}(s)r(s)ds + Y(t)Y^{-1}(t)r(t)$$

Boundary Value Problems: Well-posedness

Proof.

Uniqueness of solution follows from the fact that, if $y_1(t)$ and $y_2(t)$ are two solutions to the BVP, then $y(t) = y_1(t) - y_2(t)$ satisfies

$$y'(t) = A(t)y(t), \quad B_a y(a) + B_b y(b) = 0,$$

and hence $y(t)$ must have the form $y(t) = Y(t)d$ for some $d \in \mathbb{R}^n$ satisfying, $Qd = 0$.

Now, one can see that the unique solution to the BVP is given by

$$y(t) = Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(t) \int_a^t Y^{-1}(s)r(s)ds$$

by directly verifying that it satisfies the ODE and the boundary condition. Indeed,

$$y'(t) = A(t)Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + A(t)Y(t) \int_a^t Y^{-1}(s)r(s)ds + Y(t)Y^{-1}(t)r(t)$$

$$y(a) = Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right),$$

$$y(b) = Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(b) \int_a^b Y^{-1}(s)r(s)ds.$$

Boundary Value Problems: Well-posedness

Proof.

Now, one can see that the unique solution to the BVP is given by

$$y(t) = Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(t) \int_a^t Y^{-1}(s)r(s)ds$$

by directly verifying that it satisfies the ODE and the boundary condition. Indeed,

$$y'(t) = A(t)Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + A(t)Y(t) \int_a^t Y^{-1}(s)r(s)ds + Y(t)Y^{-1}(t)r(t),$$

$$y(a) = Y(a)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right),$$

$$y(b) = Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(b) \int_a^b Y^{-1}(s)r(s)ds.$$

Therefore,

$$B_a y(a) + B_b y(b)$$

Boundary Value Problems: Well-posedness

Proof.

Now, one can see that the unique solution to the BVP is given by

$$y(t) = Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(t) \int_a^t Y^{-1}(s)r(s)ds$$

by directly verifying that it satisfies the ODE and the boundary condition. Indeed,

$$y'(t) = A(t)Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + A(t)Y(t) \int_a^t Y^{-1}(s)r(s)ds + Y(t)Y^{-1}(t)r(t),$$

$$y(a) = Y(a)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right),$$

$$y(b) = Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(b) \int_a^b Y^{-1}(s)r(s)ds.$$

Therefore,

$$\begin{aligned} & B_a y(a) + B_b y(b) \\ &= B_a Y(a)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + B_b Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) \\ &+ B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \end{aligned}$$

Boundary Value Problems: Well-posedness

Proof.

Now, one can see that the unique solution to the BVP is given by

$$y(t) = Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(t) \int_a^t Y^{-1}(s)r(s)ds$$

by directly verifying that it satisfies the ODE and the boundary condition. Indeed,

$$y'(t) = A(t)Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + A(t)Y(t) \int_a^t Y^{-1}(s)r(s)ds + Y(t)Y^{-1}(t)r(t),$$

$$y(a) = Y(a)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right),$$

$$y(b) = Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(b) \int_a^b Y^{-1}(s)r(s)ds.$$

Therefore,

$$\begin{aligned} & B_a y(a) + B_b y(b) \\ &= B_a Y(a)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + B_b Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) \\ &+ B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \\ &= (B_a Y(a)Q^{-1} + B_b Y(b)Q^{-1})c - (B_a Y(a)Q^{-1} + B_b Y(b)Q^{-1} - I)B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \end{aligned}$$

Boundary Value Problems: Well-posedness

Proof.

Now, one can see that the unique solution to the BVP is given by

$$y(t) = Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(t) \int_a^t Y^{-1}(s)r(s)ds$$

by directly verifying that it satisfies the ODE and the boundary condition. Indeed,

$$y'(t) = A(t)Y(t)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + A(t)Y(t) \int_a^t Y^{-1}(s)r(s)ds + Y(t)Y^{-1}(t)r(t),$$

$$y(a) = Y(a)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right),$$

$$y(b) = Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + Y(b) \int_a^b Y^{-1}(s)r(s)ds.$$

Therefore,

$$\begin{aligned} & B_a y(a) + B_b y(b) \\ &= B_a Y(a)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) + B_b Y(b)Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \right) \\ &+ B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds \\ &= (B_a Y(a)Q^{-1} + B_b Y(b)Q^{-1})c - (B_a Y(a)Q^{-1} + B_b Y(b)Q^{-1} - I)B_b Y(b) \int_a^b Y^{-1}(s)r(s)ds = c. \end{aligned}$$

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP as

$$y'(t) = y'_0(t) + A(t)Yd$$

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP as

$$y'(t) = y'_0(t) + A(t)Yd = (A(t)y_0 + r(t)) + A(t)Yd$$

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP as

$$y'(t) = y'_0(t) + A(t)Yd = (A(t)y_0 + r(t)) + A(t)Yd = A(t)y + r(t)$$

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP as

$$y'(t) = y'_0(t) + A(t)Yd = (A(t)y_0 + r(t)) + A(t)Yd = A(t)y + r(t)$$

and

$$B_a y(a) + B_b y(b)$$

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP as

$$y'(t) = y'_0(t) + A(t)Yd = (A(t)y_0 + r(t)) + A(t)Yd = A(t)y + r(t)$$

and

$$B_a y(a) + B_b y(b) = B_a y_0(a) + B_b y_0(b) + B_a Yd + B_b Yd$$

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP as

$$y'(t) = y'_0(t) + A(t)Yd = (A(t)y_0 + r(t)) + A(t)Yd = A(t)y + r(t)$$

and

$$B_a y(a) + B_b y(b) = B_a y_0(a) + B_b y_0(b) + B_a Yd + B_b Yd = c + Qd = c.$$

Boundary Value Problems: Well-posedness

Proof.

Therefore,

$$\begin{aligned}
 & B_a y(a) + B_b y(b) \\
 &= B_a Y(a) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) + B_b Y(b) Q^{-1} \left(c - B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \right) \\
 &+ B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds \\
 &= (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1}) c - (B_a Y(a) Q^{-1} + B_b Y(b) Q^{-1} - I) B_b Y(b) \int_a^b Y^{-1}(s) r(s) ds = c.
 \end{aligned}$$

Finally, for the converse, assume that the BVP has a unique solution, say $y_0(t)$, but Q is not invertible.

Then there is $0 \neq d \in \mathbb{R}^n$ such that $Qd = 0$, and therefore, $y(t) = y_0(t) + Yd$ is another solution to the BVP as

$$y'(t) = y'_0(t) + A(t)Yd = (A(t)y_0 + r(t)) + A(t)Yd = A(t)y + r(t)$$

and

$$B_a y(a) + B_b y(b) = B_a y_0(a) + B_b y_0(b) + B_a Yd + B_b Yd = c + Qd = c.$$

A contradiction.

Boundary Value Problems: Well-posedness

If we define $\Phi(t) = Y(t)Q^{-1}$ and the Green's function

$$G(t, s) = \begin{cases} \Phi(t)B_a\Phi(a)\Phi^{-1}(s), & a \leq s \leq t, \\ -\Phi(t)B_b\Phi(b)\Phi^{-1}(s), & t < s \leq b. \end{cases}$$

Then the solution $y(t)$ can be expressed compactly as

$$y(t) = \Phi(t)c + \int_a^b G(t, s)r(s)ds.$$

Boundary Value Problems: Well-posedness

If we define $\Phi(t) = Y(t)Q^{-1}$ and the Green's function

$$G(t, s) = \begin{cases} \Phi(t)B_a\Phi(a)\Phi^{-1}(s), & a \leq s \leq t, \\ -\Phi(t)B_b\Phi(b)\Phi^{-1}(s), & t < s \leq b. \end{cases}$$

Then the solution $y(t)$ can be expressed compactly as

$$y(t) = \Phi(t)c + \int_a^b G(t, s)r(s)ds.$$

Consider the perturbed problem

$$\hat{y}' = A(t)\hat{y} + \hat{r}(t), \quad a < t < b,$$

with boundary conditions

$$B_a\hat{y}(a) + B_b\hat{y}(b) = \hat{c}.$$

Boundary Value Problems: Well-posedness

If we define $\Phi(t) = Y(t)Q^{-1}$ and the Green's function

$$G(t, s) = \begin{cases} \Phi(t)B_a\Phi(a)\Phi^{-1}(s), & a \leq s \leq t, \\ -\Phi(t)B_b\Phi(b)\Phi^{-1}(s), & t < s \leq b. \end{cases}$$

Then the solution $y(t)$ can be expressed compactly as

$$y(t) = \Phi(t)c + \int_a^b G(t, s)r(s)ds.$$

Consider the perturbed problem

$$\hat{y}' = A(t)\hat{y} + \hat{r}(t), \quad a < t < b,$$

with boundary conditions

$$B_a\hat{y}(a) + B_b\hat{y}(b) = \hat{c}.$$

Let $z(t) = \hat{y}(t) - y(t)$, $\Delta r(t) = \hat{r}(t) - r(t)$, and $\Delta c(t) = \hat{c}(t) - c(t)$. Then, $z(t)$ satisfies the BVP

$$z' = A(t)z + \Delta r(t), \quad a < t < b,$$

with boundary conditions

$$B_az(a) + B_bz(b) = \Delta c.$$

Boundary Value Problems: Well-posedness

If we define $\Phi(t) = Y(t)Q^{-1}$ and the Green's function

$$G(t, s) = \begin{cases} \Phi(t)B_a\Phi(a)\Phi^{-1}(s), & a \leq s \leq t, \\ -\Phi(t)B_b\Phi(b)\Phi^{-1}(s), & t < s \leq b. \end{cases}$$

Then the solution $y(t)$ can be expressed compactly as

$$y(t) = \Phi(t)c + \int_a^b G(t, s)r(s)ds.$$

Consider the perturbed problem

$$\hat{y}' = A(t)\hat{y} + \hat{r}(t), \quad a < t < b,$$

with boundary conditions

$$B_a\hat{y}(a) + B_b\hat{y}(b) = \hat{c}.$$

Let $z(t) = \hat{y}(t) - y(t)$, $\Delta r(t) = \hat{r}(t) - r(t)$, and $\Delta c(t) = \hat{c}(t) - c(t)$. Then, $z(t)$ satisfies the BVP

$$z' = A(t)z + \Delta r(t), \quad a < t < b,$$

with boundary conditions

$$B_az(a) + B_bz(b) = \Delta c.$$

Therefore,

$$\|z\| \leq \max\{\|\Phi\|, \|G\|\} \left(|\Delta c| + \int_a^b |\Delta r(s)|ds \right).$$

Boundary Value Problems: Well-posedness

If we define $\Phi(t) = Y(t)Q^{-1}$ and the Green's function

$$G(t, s) = \begin{cases} \Phi(t)B_a\Phi(a)\Phi^{-1}(s), & a \leq s \leq t, \\ -\Phi(t)B_b\Phi(b)\Phi^{-1}(s), & t < s \leq b. \end{cases}$$

Then the solution $y(t)$ can be expressed compactly as

$$y(t) = \Phi(t)c + \int_a^b G(t, s)r(s)ds.$$

Consider the perturbed problem

$$\hat{y}' = A(t)\hat{y} + \hat{r}(t), \quad a < t < b,$$

with boundary conditions

$$B_a\hat{y}(a) + B_b\hat{y}(b) = \hat{c}.$$

Let $z(t) = \hat{y}(t) - y(t)$, $\Delta r(t) = \hat{r}(t) - r(t)$, and $\Delta c(t) = \hat{c}(t) - c(t)$. Then, $z(t)$ satisfies the BVP

$$z' = A(t)z + \Delta r(t), \quad a < t < b,$$

with boundary conditions

$$B_az(a) + B_bz(b) = \Delta c.$$

Therefore,

$$\|z\| \leq \max\{\|\Phi\|, \|G\|\} \left(|\Delta c| + \int_a^b |\Delta r(s)|ds \right).$$

Absolute Condition
Number