

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE

- Finite Difference Method

- More Stability Analysis – Energy Estimate



Numerical Methods for PDE: 2nd Order Elliptic PDE



Stability Analysis using an energy estimate



Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

We denote the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^N \sum_{n=1}^N u(mh, nh) v(mh, nh)$$

with the corresponding norm $\|\cdot\|_h$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

We denote the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^N \sum_{n=1}^N u(mh, nh) v(mh, nh)$$

with the corresponding norm $\|\cdot\|_h$.

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

We denote the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^N \sum_{n=1}^N u(mh, nh) v(mh, nh)$$

with the corresponding norm $\|\cdot\|_h$.

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$|v(mh, nh)|^2 \leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

We denote the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^N \sum_{n=1}^N u(mh, nh) v(mh, nh)$$

with the corresponding norm $\|\cdot\|_h$.

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$|v(mh, nh)|^2 \leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2 \leq \left(\sum_{k=1}^N |v(kh, nh) - v((k-1)h, nh)| \right)^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

We denote the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^N \sum_{n=1}^N u(mh, nh) v(mh, nh)$$

with the corresponding norm $\|\cdot\|_h$.

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$\begin{aligned} |v(mh, nh)|^2 &\leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2 \leq \left(\sum_{k=1}^N |v(kh, nh) - v((k-1)h, nh)| \right)^2 \\ &\leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)| \right)^2 \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

We denote the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^N \sum_{n=1}^N u(mh, nh) v(mh, nh)$$

with the corresponding norm $\|\cdot\|_h$.

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$\begin{aligned} |v(mh, nh)|^2 &\leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2 \leq \left(\sum_{k=1}^N |v(kh, nh) - v((k-1)h, nh)| \right)^2 \\ &\leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)| \right)^2 \leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2 \right) \left(h \sum_{k=1}^N 1^2 \right) \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Stability Analysis using an energy estimate

Let $v \in L(\Omega_h)$ and define the backward difference operator

$$\partial_{x_1} v(mh, nh) = \frac{v(mh, nh) - v((m-1)h, nh)}{h}, 1 \leq m \leq N, 0 \leq n \leq N.$$

We denote the inner product

$$\langle u, v \rangle_h = h^2 \sum_{m=1}^N \sum_{n=1}^N u(mh, nh) v(mh, nh)$$

with the corresponding norm $\|\cdot\|_h$.

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$\begin{aligned} |v(mh, nh)|^2 &\leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2 \leq \left(\sum_{k=1}^N |v(kh, nh) - v((k-1)h, nh)| \right)^2 \\ &\leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)| \right)^2 \leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2 \right) \left(h \sum_{k=1}^N 1^2 \right) = h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2. \end{aligned}$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$\begin{aligned} |v(mh, nh)|^2 &\leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2 \leq \left(\sum_{k=1}^N |v(kh, nh) - v((k-1)h, nh)| \right)^2 \\ &\leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)| \right)^2 \leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2 \right) \left(h \sum_{k=1}^N 1^2 \right) = h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2. \end{aligned}$$

Therefore,

$$\sum_{m=1}^N |v(mh, nh)|^2 \leq \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$\begin{aligned} |v(mh, nh)|^2 &\leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2 \leq \left(\sum_{k=1}^N |v(kh, nh) - v((k-1)h, nh)| \right)^2 \\ &\leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)| \right)^2 \leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2 \right) \left(h \sum_{k=1}^N 1^2 \right) = h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2. \end{aligned}$$

Therefore,

$$h \sum_{m=1}^N |v(mh, nh)|^2 \leq h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Lemma

If $v \in L(\Omega_h)$, then $\|v\|_h \leq \|\partial_{x_1} v\|_h$.

Proof:

For $1 \leq m \leq N, 0 \leq n \leq N$,

$$\begin{aligned} |v(mh, nh)|^2 &\leq \left(\sum_{k=1}^m |v(kh, nh) - v((k-1)h, nh)| \right)^2 \leq \left(\sum_{k=1}^N |v(kh, nh) - v((k-1)h, nh)| \right)^2 \\ &\leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)| \right)^2 \leq \left(h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2 \right) \left(h \sum_{k=1}^N 1^2 \right) = h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2. \end{aligned}$$

Therefore,

$$h \sum_{m=1}^N |v(mh, nh)|^2 \leq h \sum_{k=1}^N |\partial_{x_1} v(kh, nh)|^2$$

and

$$h \sum_{m=1}^N \sum_{n=1}^N |v(mh, nh)|^2 \leq h \sum_{k=1}^N \sum_{n=1}^N |\partial_{x_1} v(kh, nh)|^2.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Lemma

If $v, w \in L(\Omega_h)$, then

$$-\langle \Delta_h v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h + \langle \partial_{x_2} v, \partial_{x_2} w \rangle_h.$$

Proof:

For $v_0, v_1, \dots, v_N, w_0, w_1, \dots, w_N \in \mathbb{R}$ with $w_0 = w_N = 0$. Then,

$$\begin{aligned} \sum_{k=1}^N (v_k - v_{k-1})(w_k - w_{k-1}) &= \sum_{k=1}^N v_k w_k + \sum_{k=1}^N v_{k-1} w_{k-1} - \sum_{k=1}^N v_{k-1} w_k - \sum_{k=1}^N v_k w_{k-1} \\ &= 2 \sum_{k=1}^{N-1} v_k w_k - \sum_{k=1}^{N-1} v_{k-1} w_k - \sum_{k=1}^{N-1} v_{k+1} w_k = - \sum_{k=1}^{N-1} (v_{k+1} - 2v_k + v_{k-1}) w_k. \end{aligned}$$

Hence,

$$-h \sum_{k=1}^{N-1} \frac{v((k+1)h, nh) - 2v(kh, nh) + v((k-1)h, nh)}{h^2} w(kh, nh) = h \sum_{k=1}^N \partial_{x_1} v(kh, nh) \partial_{x_1} w(kh, nh)$$

and thus

$$-\langle D_{h,x_1}^2 v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Lemma

If $v, w \in L(\Omega_h)$, then

$$-\langle \Delta_h v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h + \langle \partial_{x_2} v, \partial_{x_2} w \rangle_h.$$

Proof:

Hence,

$$-h \sum_{k=1}^{N-1} \frac{v((k+1)h, nh) - 2v(kh, nh) + v((k-1)h, nh)}{h^2} w(kh, nh) = h \sum_{k=1}^N \partial_{x_1} v(kh, nh) \partial_{x_1} w(kh, nh)$$

and thus

$$-\langle D_{h,x_1}^2 v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h.$$

Similarly

$$-\langle D_{h,x_2}^2 v, w \rangle_h = \langle \partial_{x_2} v, \partial_{x_2} w \rangle_h.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Lemma

If $v, w \in L(\Omega_h)$, then

$$-\langle \Delta_h v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h + \langle \partial_{x_2} v, \partial_{x_2} w \rangle_h.$$

Proof:

Hence,

$$-h \sum_{k=1}^{N-1} \frac{v((k+1)h, nh) - 2v(kh, nh) + v((k-1)h, nh)}{h^2} w(kh, nh) = h \sum_{k=1}^N \partial_{x_1} v(kh, nh) \partial_{x_1} w(kh, nh)$$

and thus

$$-\langle D_{h,x_1}^2 v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h.$$

Similarly

$$-\langle D_{h,x_2}^2 v, w \rangle_h = \langle \partial_{x_2} v, \partial_{x_2} w \rangle_h.$$

Combining the two results, we get the following stability result:

If $v \in L(\Omega_h)$, then

$$\|v\|_h^2 \leq \|\partial_{x_1} v\|_h^2 \leq \|\partial_{x_1} v\|_h^2 + \|\partial_{x_2} v\|_h^2 = -\langle \Delta_h v, v \rangle_h \leq \|\Delta_h v\|_h \|v\|_h,$$

where $\Delta_h v$ is extended to the boundary grid Γ_h as zero while computing $\|\Delta_h v\|_h$.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Lemma

If $v, w \in L(\Omega_h)$, then

$$-\langle \Delta_h v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h + \langle \partial_{x_2} v, \partial_{x_2} w \rangle_h.$$

Proof:

Hence,

$$-h \sum_{k=1}^{N-1} \frac{v((k+1)h, nh) - 2v(kh, nh) + v((k-1)h, nh)}{h^2} w(kh, nh) = h \sum_{k=1}^N \partial_{x_1} v(kh, nh) \partial_{x_1} w(kh, nh)$$

and thus

$$-\langle D_{h,x_1}^2 v, w \rangle_h = \langle \partial_{x_1} v, \partial_{x_1} w \rangle_h.$$

Similarly

$$-\langle D_{h,x_2}^2 v, w \rangle_h = \langle \partial_{x_2} v, \partial_{x_2} w \rangle_h.$$

Combining the two results, we get the following stability result:

If $v \in L(\Omega_h)$, then

$$\|v\|_h^2 \leq \|\partial_{x_1} v\|_h^2 \leq \|\partial_{x_1} v\|_h^2 + \|\partial_{x_2} v\|_h^2 = -\langle \Delta_h v, v \rangle_h \leq \|\Delta_h v\|_h \|v\|_h,$$

where $\Delta_h v$ is extended to the boundary grid Γ_h as zero while computing $\|\Delta_h v\|_h$. Thus, $\|v\|_h \leq \|\Delta_h v\|_h$.

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE

- Finite Difference Method**
- General Domains**



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP


$$\begin{aligned}u'' &= f(t), & a < t < b, \\u(a) &= 0, & u(b) = 0,\end{aligned}$$

to two dimensions is

$$\begin{aligned}\Delta u &:= u_{x_1x_1} + u_{x_2x_2} = f, & \text{in } \Omega, \\u &= g, & \text{on } \Gamma.\end{aligned}$$

For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.

$N = 8$



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned}u'' &= f(t), & a < t < b, \\u(a) &= 0, & u(b) = 0,\end{aligned}$$


to two dimensions is

$$\begin{aligned}\Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\u &= g, & \text{on } \Gamma.\end{aligned}$$

~~For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.~~

Now, let's look at general domains.

$N = 8$



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned}u'' &= f(t), & a < t < b, \\u(a) &= 0, & u(b) = 0,\end{aligned}$$

to two dimensions is


$$\begin{aligned}\Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\u &= g, & \text{on } \Gamma.\end{aligned}$$

~~For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.~~

Now, let's look at general domains.

Again, $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points.

$N = 8$



Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) = 0, \end{aligned}$$

$N = 8$


to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$

~~For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.~~

Now, let's look at general domains.

Again, $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points. The set of points in Ω_h whose all four neighbors are in Ω_h are denoted by Ω_h° .



A mesh point
with all four
nearest
neighbors in
the interior

Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

to two dimensions is


$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$

~~For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.~~

Now, let's look at general domains.

Again, $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points. The set of points in Ω_h whose all four neighbors are in Ω_h are denoted by Ω_h° .

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$


~~For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.~~

Now, let's look at general domains.

Again, $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points. The set of points in Ω_h whose all four neighbors are in Ω_h are denoted by Ω_h° .

Also, $\Gamma_h = \Gamma \cap \mathbb{R}_h^2$ and $\bar{\Omega}_h = \Omega \cup \Gamma_h$.

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) = 0, \end{aligned}$$

to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$

~~For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.~~


Now, let's look at general domains.

Again, $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points. The set of points in Ω_h whose all four neighbors are in Ω_h are denoted by Ω_h° .

Also, $\Gamma_h = \Gamma \cap \mathbb{R}_h^2$ and $\bar{\Omega}_h = \Omega \cup \Gamma_h$.

On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

A natural generalization to the two-point BVP

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

to two dimensions is

$$\begin{aligned} \Delta u &:= u_{x_1 x_1} + u_{x_2 x_2} = f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma. \end{aligned}$$

~~For simplicity, we will first consider a very simple domain $\Omega = (0,1) \times (0,1)$.~~

Now, let's look at general domains.


Again, $\Omega_h = \Omega \cap \mathbb{R}_h^2$, the set of interior mesh points. The set of points in Ω_h whose all four neighbors are in Ω_h are denoted by Ω_h° .

Also, $\Gamma_h = \Gamma \cap \mathbb{R}_h^2$ and $\bar{\Omega}_h = \Omega \cup \Gamma_h$.

On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

For $(x_1, x_2) \in \Omega_h \setminus \Omega_h^\circ$, let $(x_1 + h_1, x_2)$, $(x_1, x_2 + h_2)$, $(x_1 - h_1, x_2)$, and $(x_1, x_2 - h_4)$ be the nearest neighbors (with $0 \leq h_k \leq h$), and let v_1, v_2, v_3 and v_4 denote the values of v at these four points. And finally, let $v_0 = v(x_1, x_2)$.

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE


On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

For $(x_1, x_2) \in \Omega_h \setminus \Omega_h^\circ$, let $(x_1 + h_1, x_2)$, $(x_1, x_2 + h_2)$, $(x_1 - h_1, x_2)$, and $(x_1, x_2 - h_2)$ be the nearest neighbors (with $0 \leq h_k \leq h$), and let v_1, v_2, v_3 and v_4 denote the values of v at these four points. And finally, let $v_0 = v(x_1, x_2)$.

To motivate the derivation, consider the following approximation of $v''(0)$:

$$v''(0) \approx \alpha_- v(h_-) + \alpha_0 v(0) + \alpha_+ v(h_+)$$

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE


On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

For $(x_1, x_2) \in \Omega_h \setminus \Omega_h^\circ$, let $(x_1 + h_1, x_2)$, $(x_1, x_2 + h_2)$, $(x_1 - h_1, x_2)$, and $(x_1, x_2 - h_2)$ be the nearest neighbors (with $0 \leq h_k \leq h$), and let v_1, v_2, v_3 and v_4 denote the values of v at these four points. And finally, let $v_0 = v(x_1, x_2)$.

To motivate the derivation, consider the following approximation of $v''(0)$:

$$\begin{aligned} v''(0) &\approx \alpha_- v(h_-) + \alpha_0 v(0) + \alpha_+ v(h_+) \\ &\approx (\alpha_- + \alpha_0 + \alpha_+) v(0) + (\alpha_+ h_+ - \alpha_- h_-) v'(0) \\ &\quad + \frac{1}{2} (\alpha_- h_-^2 + \alpha_+ h_+^2) v'' + \dots \end{aligned}$$

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

For $(x_1, x_2) \in \Omega_h \setminus \Omega_h^\circ$, let $(x_1 + h_1, x_2)$, $(x_1, x_2 + h_2)$, $(x_1 - h_1, x_2)$, and $(x_1, x_2 - h_2)$ be the nearest neighbors (with $0 \leq h_k \leq h$), and let v_1, v_2, v_3 and v_4 denote the values of v at these four points. And finally, let $v_0 = v(x_1, x_2)$.

To motivate the derivation, consider the following approximation of $v''(0)$:

$$\begin{aligned} v''(0) &\approx \alpha_- v(h_-) + \alpha_0 v(0) + \alpha_+ v(h_+) \\ &\approx (\alpha_- + \alpha_0 + \alpha_+) v(0) + (\alpha_+ h_+ - \alpha_- h_-) v'(0) \\ &\quad + \frac{1}{2} (\alpha_- h_-^2 + \alpha_+ h_+^2) v'' + \dots \end{aligned}$$


Thus, to obtain a consistent approximation, we must have

$$\alpha_- + \alpha_0 + \alpha_+ = 0$$

$$\alpha_+ h_+ - \alpha_- h_- = 0$$

$$\alpha_- h_-^2 + \alpha_+ h_+^2 = 2$$

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

For $(x_1, x_2) \in \Omega_h \setminus \Omega_h^\circ$, let $(x_1 + h_1, x_2)$, $(x_1, x_2 + h_2)$, $(x_1 - h_1, x_2)$, and $(x_1, x_2 - h_2)$ be the nearest neighbors (with $0 \leq h_k \leq h$), and let v_1, v_2, v_3 and v_4 denote the values of v at these four points. And finally, let $v_0 = v(x_1, x_2)$.

To motivate the derivation, consider the following approximation of $v''(0)$:

$$\begin{aligned} v''(0) &\approx \alpha_- v(h_-) + \alpha_0 v(0) + \alpha_+ v(h_+) \\ &\approx (\alpha_- + \alpha_0 + \alpha_+) v(0) + (\alpha_+ h_+ - \alpha_- h_-) v'(0) \\ &\quad + \frac{1}{2} (\alpha_- h_-^2 + \alpha_+ h_+^2) v'' + \dots \end{aligned}$$

Thus, to obtain a consistent approximation, we must have

$$\alpha_- + \alpha_0 + \alpha_+ = 0$$


$$\alpha_+ h_+ - \alpha_- h_- = 0$$

$$\alpha_- h_-^2 + \alpha_+ h_+^2 = 2$$

which yields

$$\alpha_- = \frac{2}{h_-(h_- + h_+)}, \alpha_+ = \frac{2}{h_+(h_- + h_+)}, \alpha_0 = -\frac{2}{h_- h_+}.$$

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

For $(x_1, x_2) \in \Omega_h \setminus \Omega_h^\circ$, let $(x_1 + h_1, x_2)$, $(x_1, x_2 + h_2)$, $(x_1 - h_1, x_2)$, and $(x_1, x_2 - h_4)$ be the nearest neighbors (with $0 \leq h_k \leq h$), and let v_1, v_2, v_3 and v_4 denote the values of v at these four points. And finally, let $v_0 = v(x_1, x_2)$.

To motivate the derivation, consider the following approximation of $v''(0)$:


$$v''(0) \approx \frac{\alpha_-}{2} v(h_-) + \alpha_0 v(0) + \frac{\alpha_+}{2} v(h_+)$$

$$\alpha_- = \frac{2}{h_-(h_- + h_+)}, \alpha_+ = \frac{2}{h_+(h_- + h_+)}, \alpha_0 = -\frac{2}{h_- h_+}.$$

This calculation leads us to the Shortley-Weller formula for $\Delta_h v$:

$$\begin{aligned} \Delta_h v(x_1, x_2) = & \frac{2}{h_1(h_1+h_3)} v_1 + \frac{2}{h_2(h_2+h_4)} v_2 + \frac{2}{h_3(h_1+h_3)} v_3 \\ & + \frac{2}{h_4(h_2+h_4)} v_4 - \left(\frac{2}{h_1 h_3} + \frac{2}{h_2 h_4} \right) v_0 \end{aligned}$$

$N = 8$



A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

On Ω_h° , $\Delta_h v$ is defined as the usual 5-point Laplacian.

For $(x_1, x_2) \in \Omega_h \setminus \Omega_h^\circ$, let $(x_1 + h_1, x_2)$, $(x_1, x_2 + h_2)$, $(x_1 - h_1, x_2)$, and $(x_1, x_2 - h_2)$ be the nearest neighbors (with $0 \leq h_k \leq h$), and let v_1, v_2, v_3 and v_4 denote the values of v at these four points. And finally, let $v_0 = v(x_1, x_2)$.

To motivate the derivation, consider the following approximation of $v''(0)$:

$$v''(0) \approx \frac{\alpha_-}{2} v(h_-) + \alpha_0 v(0) + \frac{\alpha_+}{2} v(h_+)$$

$$\alpha_- = \frac{2}{h_-(h_- + h_+)}, \alpha_+ = \frac{2}{h_+(h_- + h_+)}, \alpha_0 = -\frac{2}{h_- h_+}.$$

This calculation leads us to the Shortley-Weller formula for $\Delta_h v$:

$$\Delta_h v(x_1, x_2) = \frac{2}{h_1(h_1 + h_3)} v_1 + \frac{2}{h_2(h_2 + h_4)} v_2 + \frac{2}{h_3(h_1 + h_3)} v_3$$


$$+ \frac{2}{h_4(h_2 + h_4)} v_4 - \left(\frac{2}{h_1 h_3} + \frac{2}{h_2 h_4} \right) v_0$$

Using Taylor's theorem with remainder, we can easily see that for $v \in C^3(\bar{\Omega})$,

$$\|\Delta v - \Delta_h v\|_{\infty, \Omega_h} = \frac{2M_3}{3} h,$$

where M_3 is the maximum of the L^∞ norms of the third derivative of v .

$N = 8$




A mesh point
with all four
nearest
neighbors in
the interior

A mesh point with at
least one nearest
neighbor on the
boundary

Numerical Methods for PDE: 2nd Order Elliptic PDE

We can obtain the discrete maximum/minimum principle with virtually the same proof as for the square domain and then a stability result follows as before (*exercise*). In this way, we can obtain an $O(h)$ convergence result.

$N = 8$




Numerical Methods for PDE: 2nd Order Elliptic PDE

We can obtain the discrete maximum/minimum principle with virtually the same proof as for the square domain and then a stability result follows as before (*exercise*). In this way, we can obtain an $O(h)$ convergence result.

$N = 8$

In fact, this result can be improved. Although the truncation error is only $O(h)$, it is $O(h^2)$ all points except those neighboring the boundary and these account for only $O(h^{-1})$ of the total $O(h^{-2})$ points in Ω_h . Moreover, these points are within h of the boundary where the solution is known exactly. For both these reasons, the contribution to the error from these points is smaller than what we observed earlier.



Numerical Methods for PDE: 2nd Order Elliptic PDE

We can obtain the discrete maximum/minimum principle with virtually the same proof as for the square domain and then a stability result follows as before (*exercise*). In this way, we can obtain an $O(h)$ convergence result.

$N = 8$

In fact, this result can be improved. Although the truncation error is only $O(h)$, it is $O(h^2)$ all points except those neighboring the boundary and these account for only $O(h^{-1})$ of the total $O(h^{-2})$ points in Ω_h . Moreover, these points are within h of the boundary where the solution is known exactly. For both these reasons, the contribution to the error from these points is smaller than what we observed earlier.

Theorem

Let u be the solution to

$$\begin{aligned} \Delta u &= f, & \text{in } \Omega, \\ u &= g, & \text{on } \Gamma, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} \Delta_h u_h &= f, & \text{on } \Omega_h, \\ u_h &= g, & \text{on } \Gamma_h. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, \bar{\Omega}_h} \leq \frac{M_4 d^2}{96} h^2 + \frac{2M_3}{3} h^3,$$

where d is the diameter of the smallest disc containing Ω and M_k is the maximum of the L^∞ norms of the k th derivative of v .

