

Boundary Value Problems: Finite Difference Method

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$

Proof:

It is sufficient to show that, if $D_h^2 u_h(t_i) = 0$ for $t_i, i = 1, \dots, n$, and $u_h(a) = 0$, $u_h(b) = 0$, then $u_h \equiv 0$.

Boundary Value Problems: Finite Difference Method

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$

Proof:

It is sufficient to show that, if $D_h^2 u_h(t_i) = 0$ for $t_i, i = 1, \dots, n$, and $u_h(a) = 0$, $u_h(b) = 0$, then $u_h \equiv 0$. But this follows from the discrete maximum principle and discrete minimum principle.

Boundary Value Problems: Finite Difference Method

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$

Proof:

It is sufficient to show that, if $D_h^2 u_h(t_i) = 0$ for $t_i, i = 1, \dots, n$, and $u_h(a) = 0$, $u_h(b) = 0$, then $u_h \equiv 0$. But this follows from the discrete maximum principle and discrete minimum principle.

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

$$\text{Let } w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

therefore,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \max_{1 \leq i \leq n} (u_h(t_i) + w(t_i)\|f\|_{\infty, h}) \leq \max\{u_h(a) + w(a)\|f\|_{\infty, h}, (u_h(b) + w(b)\|f\|_{\infty, h})\}.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

therefore,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \max_{1 \leq i \leq n} (u_h(t_i) + w(t_i)\|f\|_{\infty, h}) \leq \max\{u_h(a) + w(a)\|f\|_{\infty, h}, (u_h(b) + w(b)\|f\|_{\infty, h})\}.$$

Hence,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{\alpha, \beta\}.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

therefore,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \max_{1 \leq i \leq n} (u_h(t_i) + w(t_i)\|f\|_{\infty, h}) \leq \max\{u_h(a) + w(a)\|f\|_{\infty, h}, (u_h(b) + w(b)\|f\|_{\infty, h})\}.$$

Hence,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{\alpha, \beta\}.$$

A similar argument applies to $-u_h$ giving the theorem.

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Proof:

The error estimate follows from the previous theorem applied to the discrete problem

$$\begin{aligned} D_h^2(u_h - u)(t_i) &= D_h^2 u_h(t_i) - D_h^2 u(t_i) = f(t_i) - D_h^2 u(t_i), & i = 1, \dots, n, \\ (u_h - u)(a) &= 0, & (u_h - u)(b) = 0. \end{aligned}$$

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Proof:

The error estimate follows from the previous theorem applied to the discrete problem

$$\begin{aligned} D_h^2(u_h - u)(t_i) &= D_h^2 u_h(t_i) - D_h^2 u(t_i) = f(t_i) - D_h^2 u(t_i), & i = 1, \dots, n, \\ (u_h - u)(a) &= 0, & (u_h - u)(b) = 0. \end{aligned}$$

Corollary

If $u \in C^2([a, b])$, then

$$\lim_{h \rightarrow 0} \|u_h - u\|_{\infty, h} = 0.$$

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Corollary

If $u \in C^2([a, b])$, then

$$\lim_{h \rightarrow 0} \|u_h - u\|_{\infty, h} = 0.$$

If $u \in C^4([a, b])$, then

$$\|u_h - u\|_{\infty, h} \leq \frac{h^2(b-a)^2}{96} \|u^{(4)}\|_{\infty, [a, b]}.$$



Boundary Value Problems: Finite Difference Method

Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.



Boundary Value Problems: Finite Difference Method

Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.

An estimate of the form $\|v\| \leq C_h \|f\|$ whenever $D_h^2 v(t_i) = f(t_i)$, $i = 1, \dots, n$, and $v(a) = 0$, $v(b) = 0$, is a **stability estimate** and if it holds with C_h independent of h , we say that the discretization is **stable**.

Boundary Value Problems: Finite Difference Method

Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.

An estimate of the form $\|v\| \leq C_h \|f\|$ whenever $D_h^2 v(t_i) = f(t_i), i = 1, \dots, n$, and $v(a) = 0, v(b) = 0$, is a **stability estimate** and if it holds with C_h independent of h , we say that the discretization is **stable**.

The previous result can be summarized as “**consistency + stability implies convergence**”.

Boundary Value Problems: Finite Difference Method

Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.

An estimate of the form $\|v\| \leq C_h \|f\|$ whenever $D_h^2 v(t_i) = f(t_i), i = 1, \dots, n$, and $v(a) = 0, v(b) = 0$, is a **stability estimate** and if it holds with C_h independent of h , we say that the discretization is **stable**.

The previous result can be summarized as “**consistency + stability implies convergence**”.

As a final remark, the finite difference method helped us find the solution values at the mesh points, but the solution at non-mesh points are not readily available from the method. If needed, one can obtain the solution at non-mesh points through interpolation or try other approximation approaches ...

Lesson 3

Boundary Value Problems for ODEs

3.2 Shooting Method

3.3 Finite Difference Method

3.4 Variational Methods



Boundary Value Problems: Variational Methods



One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

Boundary Value Problems: Variational Methods



One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Boundary Value Problems: Variational Methods

One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

We seek the solution of the form

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \varphi_i(t),$$

where $\varphi_i(t)$ are basis functions defined on $[a, b]$ and y is an n -vector of parameters to be determined.

Boundary Value Problems: Variational Methods

One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

We seek the solution of the form

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \varphi_i(t),$$

where $\varphi_i(t)$ are basis functions defined on $[a, b]$ and y is an n -vector of parameters to be determined.

Some popular finite dimensional approximation spaces are

- (1) polynomials of degree n or less.*
- (2) trigonometric polynomials of degree n or less.*
- (3) piecewise polynomials of degree n or less.*

Boundary Value Problems: Variational Methods

One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

We seek the solution of the form

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \varphi_i(t),$$

where $\varphi_i(t)$ are basis functions defined on $[a, b]$ and y is an n -vector of parameters to be determined.

Some popular finite dimensional approximation spaces are

- (1) polynomials of degree n or less.*
- (2) trigonometric polynomials of degree n or less.*
- (3) piecewise polynomials of degree n or less.*

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.