

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.2 Shooting Method

3.3 Finite Difference Method

3.4 Variational Methods

- Collocation Method



Akash Anand
MATH, IIT KANPUR



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points.



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$

that is,

$$\sum_{j=1}^n y_j \varphi_j''(t) = f\left(t_i, \sum_{j=1}^n y_j \varphi_j(t_i), \sum_{j=1}^n y_j \varphi_j'(t_i)\right), \quad i = 2, \dots, n - 1.$$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$

that is,

$$\sum_{j=1}^n y_j \varphi_j''(t) = f\left(t_i, \sum_{j=1}^n y_j \varphi_j(t_i), \sum_{j=1}^n y_j \varphi_j'(t_i)\right), \quad i = 2, \dots, n - 1.$$

In addition, we also enforce the boundary condition at the end-points:

$$\sum_{j=1}^n y_j \varphi_j(t_1) = \alpha, \quad \sum_{j=1}^n y_j \varphi_j(t_n) = \beta.$$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$

that is,

$$\sum_{j=1}^n y_j \varphi_j''(t) = f\left(t_i, \sum_{j=1}^n y_j \varphi_j(t_i), \sum_{j=1}^n y_j \varphi_j'(t_i)\right), \quad i = 2, \dots, n - 1.$$

In addition, we also enforce the boundary condition at the end-points:

$$\sum_{j=1}^n y_j \varphi_j(t_1) = \alpha, \quad \sum_{j=1}^n y_j \varphi_j(t_n) = \beta.$$

This yields a system of n algebraic equation in n unknowns. This may be linear or non-linear depending on whether f is linear or non-linear.



Boundary Value Problems: Variational Methods



In particular, for the approximation space consisting of polynomials of degree $n - 1$ or less (i.e., \mathcal{P}_{n-1}), we have

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \ell_i(t) \in \mathcal{P}_{n-1},$$

where the Lagrange basis, given by

$$\ell_i(t) = \frac{\prod_{k=1, k \neq i}^n (t - t_k)}{\prod_{k=1, k \neq i}^n (t_i - t_k)} \in \mathcal{P}_{n-1}$$

is used.

Boundary Value Problems: Variational Methods

In particular, for the approximation space consisting of polynomials of degree $n - 1$ or less (i.e., \mathcal{P}_{n-1}), we have

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \ell_i(t) \in \mathcal{P}_{n-1},$$

where the Lagrange basis, given by

$$\ell_i(t) = \frac{\prod_{k=1, k \neq i}^n (t - t_k)}{\prod_{k=1, k \neq i}^n (t_i - t_k)} \in \mathcal{P}_{n-1}$$

is used. Then the collocation method yields the following system of algebraic equations: $Ay = F(y)$ where

$$Ay = \begin{bmatrix} \ell_1(t_1) & \ell_2(t_1) & \cdots & \ell_n(t_1) \\ \ell_1''(t_2) & \ell_2''(t_2) & & \\ \vdots & \vdots & \ddots & \vdots \\ \ell_1(t_n) & \cdots & \ell_{n-1}''(t_{n-1}) & \ell_n''(t_{n-1}) \\ & & \ell_{n-1}(t_n) & \ell_n(t_n) \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix}, F(y) = \begin{bmatrix} f\left(t_2, \sum_{i=1}^n y_i \ell_i(t_2), \sum_{i=1}^n y_i \ell_i'(t_2)\right) \\ \vdots \\ f\left(t_{n-1}, \sum_{i=1}^n y_i \ell_i(t_{n-1}), \sum_{i=1}^n y_i \ell_i'(t_{n-1})\right) \end{bmatrix}.$$

α β

Boundary Value Problems: Variational Methods

In particular, for the approximation space consisting of polynomials of degree $n - 1$ or less (i.e., \mathcal{P}_{n-1}), we have

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \ell_i(t) \in \mathcal{P}_{n-1},$$

where the Lagrange basis, given by

$$\ell_i(t) = \frac{\prod_{k=1, k \neq i}^n (t - t_k)}{\prod_{k=1, k \neq i}^n (t_i - t_k)} \in \mathcal{P}_{n-1}$$

is used. Then the collocation method yields the following system of algebraic equations: $Ay = F(y)$ where

$$Ay = \begin{bmatrix} \ell_1(t_1) & \ell_2(t_1) & \cdots & \ell_n(t_1) \\ \ell_1''(t_2) & \ell_2''(t_2) & & \\ \vdots & \vdots & \ddots & \vdots \\ & & \ell_{n-1}''(t_{n-1}) & \ell_n''(t_{n-1}) \\ \ell_1(t_n) & \cdots & \ell_{n-1}(t_n) & \ell_n(t_n) \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix}, F(y) = \begin{bmatrix} f\left(t_2, \sum_{i=1}^n y_i \ell_i(t_2), \sum_{i=1}^n y_i \ell_i'(t_2)\right) \\ \vdots \\ f\left(t_{n-1}, \sum_{i=1}^n y_i \ell_i(t_{n-1}), \sum_{i=1}^n y_i \ell_i'(t_{n-1})\right) \end{bmatrix}.$$

α β

The Newton iterations take the form $y^{(m+1)} = y^{(m)} - \left(A - F'(y^{(m)})\right)^{-1} (Ay^{(m)} - F(y^{(m)}))$.



Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) &= 1. \end{aligned}$$



Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2 \left(t - \frac{1}{2} \right) (t - 1) y_1 - 4t(t - 1) y_2 + 2t \left(t - \frac{1}{2} \right) y_3.$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2 \left(t - \frac{1}{2} \right) (t - 1) y_1 - 4t(t - 1) y_2 + 2t \left(t - \frac{1}{2} \right) y_3.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\4y_1 - 8y_2 + 4y_3 &= 6 \left(\frac{1}{2} \right), \\y_3 &= 1.\end{aligned}$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2 \left(t - \frac{1}{2} \right) (t - 1) y_1 - 4t(t - 1) y_2 + 2t \left(t - \frac{1}{2} \right) y_3.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\4y_1 - 8y_2 + 4y_3 &= 6 \left(\frac{1}{2} \right), \\y_3 &= 1.\end{aligned}$$

Thus the approximate solution is

$$u(t) \approx -4t(t - 1) \left(\frac{1}{8} \right) + 2t \left(t - \frac{1}{2} \right)$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2 \left(t - \frac{1}{2} \right) (t - 1) y_1 - 4t(t - 1) y_2 + 2t \left(t - \frac{1}{2} \right) y_3.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\4y_1 - 8y_2 + 4y_3 &= 6 \left(\frac{1}{2} \right), \\y_3 &= 1.\end{aligned}$$

Thus the approximate solution is

$$u(t) \approx -4t(t - 1) \left(\frac{1}{8} \right) + 2t \left(t - \frac{1}{2} \right) = -\frac{1}{2}t + \frac{3}{2}t^2.$$

Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

Alternately, if we use the monomial basis $\{1, t, t^2\}$ for the space of quadratic polynomials, we then have approximate solution of the form

$$v(t, y) = y_1 + y_2 t + y_3 t^2.$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

Alternately, if we use the monomial basis $\{1, t, t^2\}$ for the space of quadratic polynomials, we then have approximate solution of the form

$$v(t, y) = y_1 + y_2 t + y_3 t^2.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\2y_3 &= 6 \left(\frac{1}{2} \right), \\y_1 + y_2 + y_3 &= 1.\end{aligned}$$

Thus, the approximate solution is

$$u(t) \approx -\frac{1}{2}t + \frac{3}{2}t^2.$$